

**NICOTINE POLACRILEX ORIGINAL FLAVOR- nicotine polacrilex gum, chewing**  
**NICOTINE POLACRILEX ORIGINAL FLAVOR- nicotine polacrilex gum, chewing**  
**Publix Supermarkets, Inc.**

-----  
**Nicotine Polacrilex Gum 2 mg and 4 mg Original Flavor - Publix**

### ***Drug Facts***

#### ***Active ingredient (in each chewing piece) - 2 mg***

Nicotine Polacrilex 2 mg (nicotine)

#### ***Active ingredient (in each chewing piece) - 4 mg***

Nicotine Polacrilex 4 mg (nicotine)

### ***Purpose***

Stop smoking aid

### ***Use***

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

### ***Warnings***

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

### **Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

### **Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug

- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.**

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions - 2 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Directions - 4 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Other information - 2 mg**

- **each piece contains:** calcium 115 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Other information - 4 mg**

- **each piece contains:** calcium 110 mg, sodium 11 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

**Inactive ingredients - 2 mg**

acesulfame potassium, butylated hydroxytoluene, carnauba wax, flavors, gum base, sodium bicarbonate, sodium carbonate, sorbitol and talc.

**Inactive ingredients - 4 mg**

acesulfame potassium, butylated hydroxytoluene, carnauba wax, D&C yellow #10 lake, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, sodium bicarbonate, sodium carbonate, sorbitol and talc.

## **Principal display panel 2 mg**

\*Compare to the active ingredient in Nicorette® Gum

SUGAR FREE

nicotine gum

NICOTINE POLACRILEX GUM USP, 2 mg (nicotine)

## **STOP SMOKING AID**

ORIGINAL FLAVOR

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

## **TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per da during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.
  - Not for sale to those under 18 years of age.
  - Proof of age required.
  - Not for sale in vending machines or from any source where proof of age cannot be verified.

\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette Gum.

Blister packaged for your protection. Do not use if individual seals are open or torn.

DISTRIBUTED BY PUBLIX SUPER MARKETS, INC.,

3300 PUBLIX CORPORATE PARKWAY,

LAKELAND, FL 33811, 1-888-267-3037

**Package label 2 mg**



SUGAR FREE

# nicotinegum

NICOTINE POLACRILEX GUM

USP, 2 mg (nicotine)

STOP SMOKING AID



NDC 56062-606-62

SUGAR FREE

# nicotinegum

NICOTINE POLACRILEX GUM

USP, 2 mg (nicotine)

STOP SMOKING AID

### ORIGINAL FLAVOR

- **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**
- **If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg**



ACTUAL SIZE

**2 mg** 110 PIECES OF GUM  
2 mg EACH

\*Compare to the active ingredient in Nicorette® Gum

### TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** – Chew at least **9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** – Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.

DISTRIBUTED BY PUBLIC SUPER MARKETS, INC., 3300 PUBLIC CORPORATE PARKWAY, LAKELAND, FL 33811, 1-888-267-3037  
PUBLIC.com  
PUBLIC GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK

protect from light  
nicotinegum USP, 2 mg (nicotine) contains water, flavors, gum base, sodium benzoate, sodium carboxymethylcellulose, sorbitol, and xanthan gum. ©2011 Public Super Markets, Inc. All rights reserved.

may need to be adjusted.  
Stop use and seek a doctor if:  
• you get symptoms of nicotine overdose such as nausea, vomiting, diarrhea, dizziness, weakness, and rapid heartbeat  
• you have symptoms of an allergic reaction (such as difficulty breathing or rash)  
Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Keep loose pieces of gum in paper and throw away in the trash in case of overdose; get medical help or contact a Poison Control Center (1-800-222-1222) right away.  
**Directions**  
• If you are under 18 years of age, ask a doctor before use  
• Before using this product, read the enclosed User's Guide for complete directions and other important information  
• Begin using the gum on your quit day  
• If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum  
• If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum



Not for sale to those under 18 years of age.  
Proof of age required.  
Not for sale in vending machines or from any source where proof of age cannot be verified.



**Package label 4 mg**



SUGAR FREE

# nicotinegum

NICOTINE POLACRILEX GUM

USP, 4 mg (nicotine)

STOP SMOKING AID



NDC 56062-609-62

SUGAR FREE

# nicotinegum

NICOTINE POLACRILEX GUM

USP, 4 mg (nicotine)

STOP SMOKING AID

### ORIGINAL FLAVOR

- **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.**
- **If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg**



ACTUAL SIZE

# 4 mg

 110 PIECES OF GUM  
4 mg EACH

\*Compare to the active ingredient in Nicorette® Gum

### TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** – Chew at least **9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** – Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.

PUBLIC GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK

DISTRIBUTED BY PUBLIX SUPER MARKETS, INC., 3900 PUBLIC CORPORATE PARKWAY, LAKELAND, FL 33811, 1-888-267-3037

**Directions**

- If you are under 18 years of age, ask a doctor before use
- Before using this product, read the enclosed User's Guide for complete directions and other important information
- Loosely tear the gum from your gum pack
- If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

©2011, TAC



P1D-0235A F007065





## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-608-62	11 in 1 CARTON	01/01/2014	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074507	01/01/2014	

## NICOTINE POLACRILEX ORIGINAL FLAVOR

nicotine polacrilex gum, chewing

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-609
Route of Administration	BUCCAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

### Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SORBITOL (UNII: 506T60A25R)	
TALC (UNII: 7SEV7J4R1U)	

### Product Characteristics

Color	yellow (yellowish)	Score	no score
Shape	SQUARE	Size	14mm
Flavor		Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-609-62	11 in 1 CARTON	01/01/2014	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074707	01/01/2014	

**Labeler** - Publix Supermarkets, Inc. (006922009)

Revised: 12/2022

Publix Supermarkets, Inc.