

**CHILDRENS PLUS MULTI SYMPTOM COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension**  
**Best Choice (Valu Merchandisers Company)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Drug Facts**

**Active ingredients (in each 5 mL)**

**Acetaminophen 160 mg**

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

**Purposes**

**Pain reliever/fever reducer**

Antihistamine

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - sore throat
  - cough
  - sneezing and runny nose
  - headache
  - nasal congestion
  - stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or

followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

### **Ask a doctor before use if the child has**

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

### **Ask a doctor or pharmacist before use if your child is taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dose (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or last more than 3 days
- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adult as well as children even if you do not notice any signs or symptoms.

### **Directions**

- **this product does not contain directions or complete warnings for adult use**

- mL=milliliter
- shake well before using
- do not give more than 5 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose, otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

### Other Information

- store between 20-25°C (68-77°F). Do not refrigerate

### Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### Principal Display Panel

For Ages 6-11 Years

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TYLENOL® PLUS MULTI-SYMPTOM COLD\*

### Children's Plus Multi-Symptom Cold

#### Acetaminophen

- Pain Reliever / Fever Reducer

#### Chlorpheniramine Maleate

- Antihistamine

#### Dextromethorphan HBr

- Cough Suppressant

#### Phenylephrine HCl

- Nasal Decongestant

Grape Flavor

Oral Suspension

FL OZ (mL)

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

PROUDLY DISTRIBUTED BY:

VALU MERCHANDISERS, CO.

5000 KANSAS AVE

KANSAS CITY, KS 66106

www.bestchoicebrand.com

**Package Label**

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>nervousness, dizziness or sleeplessness occurs</li> <li>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul> <p><b>Keep out of reach of children.</b></p> <p><b>Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>this product does not contain directions or complete warnings for adult use</li> <li>do not give more than 5 doses in any 24-hour period</li> <li>measure only with dosing cup provided. Do not use any other dosing device.</li> <li>if needed, repeat dose every 4 hours while symptoms last</li> <li>find the right dose on chart below. If possible, use weight to dose; otherwise, use age.</li> </ul> <table border="1"> <thead> <tr> <th>Weight (lb)</th> <th>Age (yr)</th> <th>Dose (mL)</th> </tr> </thead> <tbody> <tr> <td>48-95</td> <td>6-11</td> <td>10 mL</td> </tr> <tr> <td>36-47</td> <td>4-5</td> <td>do not use unless directed by a doctor</td> </tr> <tr> <td>under 36</td> <td>under 4</td> <td>do not use</td> </tr> </tbody> </table> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store between 20-25°C (68-77°F). Do not refrigerate.</li> </ul> <p><b>Inactive ingredients:</b> carboxymethylcellulose sodium, citric acid, D&amp;C red #33, FD&amp;C blue #1, FD&amp;C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum</p> <p><b>Questions or comments?</b> Call 1-877-753-3835 Monday-Friday 9AM-5PM EST</p>	Weight (lb)	Age (yr)	Dose (mL)	48-95	6-11	10 mL	36-47	4-5	do not use unless directed by a doctor	under 36	under 4	do not use	<p>For Ages 6 to 11 Years</p> <p><b>Best Choice®</b></p> <p>COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TYLENOL® PLUS MULTI-SYMPTOM COLD*</p> <p><b>Children's PLUS MULTI-SYMPTOM COLD</b></p>  <p><b>Acetaminophen</b></p> <ul style="list-style-type: none"> <li>Pain Reliever / Fever Reducer</li> </ul> <p><b>Chlorpheniramine Maleate</b></p> <ul style="list-style-type: none"> <li>Antihistamine</li> </ul> <p><b>Dextromethorphan HBr</b></p> <ul style="list-style-type: none"> <li>Cough Suppressant</li> </ul> <p><b>Phenylephrine HCl</b></p> <ul style="list-style-type: none"> <li>Nasal Decongestant</li> </ul> <p><b>GRAPE FLAVOR</b></p> <p>Oral Suspension 4 FL OZ (118 mL)</p>	<p>For Ages 6 to 11 Years</p> <p><b>Best Choice®</b></p> <p>COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TYLENOL® PLUS MULTI-SYMPTOM COLD*</p> <p><b>Children's PLUS MULTI-SYMPTOM COLD</b></p>  <p><b>Acetaminophen</b></p> <ul style="list-style-type: none"> <li>Pain Reliever / Fever Reducer</li> </ul> <p><b>Chlorpheniramine Maleate</b></p> <ul style="list-style-type: none"> <li>Antihistamine</li> </ul> <p><b>Dextromethorphan HBr</b></p> <ul style="list-style-type: none"> <li>Cough Suppressant</li> </ul> <p><b>Phenylephrine HCl</b></p> <ul style="list-style-type: none"> <li>Nasal Decongestant</li> </ul> <p><b>GRAPE FLAVOR</b></p> <p>Oral Suspension 4 FL OZ (118 mL)</p>	<p><b>Drug Facts</b></p> <table border="1"> <thead> <tr> <th>Active ingredients (in each 5 mL)</th> <th>Purposes</th> </tr> </thead> <tbody> <tr> <td>Acetaminophen 160 mg</td> <td>Pain reliever/fever reducer</td> </tr> <tr> <td>Chlorpheniramine maleate 1 mg</td> <td>Antihistamine</td> </tr> <tr> <td>Dextromethorphan HBr 5 mg</td> <td>Cough suppressant</td> </tr> <tr> <td>Phenylephrine HCl 2.5 mg</td> <td>Nasal decongestant</td> </tr> </tbody> </table> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves the following cold/flu symptoms</li> <li>minor aches and pains</li> <li>sore throat</li> <li>cough</li> <li>sneezing and runny nose</li> <li>headache</li> <li>nasal congestion</li> <li>stuffy nose</li> <li>temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if your child takes:</p> <ul style="list-style-type: none"> <li>more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount</li> <li>with other drugs containing acetaminophen</li> </ul> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>to make a child sleepy</li> <li>with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.</li> <li>if your child is allergic to acetaminophen or any of the inactive ingredients in this product</li> </ul> <p><b>Ask a doctor before use if your child has</b></p> <ul style="list-style-type: none"> <li>liver disease</li> <li>glaucoma</li> <li>thyroid disease</li> <li>diabetes</li> <li>heart disease</li> <li>high blood pressure</li> <li>a breathing problem such as chronic bronchitis</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>persistent or chronic cough such as occurs with asthma</li> </ul> <p><b>Ask a doctor or pharmacist before use if your child is taking</b></p> <ul style="list-style-type: none"> <li>the blood thinning drug warfarin</li> <li>sedatives or tranquilizers</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>do not exceed recommended dose (see Overdose warning)</li> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>sedatives and tranquilizers may increase drowsiness</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>new symptoms occur</li> <li>redness or swelling is present</li> <li>pain, nasal congestion or cough gets worse or lasts more than 5 days</li> <li>fever gets worse or lasts more than 3 days</li> </ul>	Active ingredients (in each 5 mL)	Purposes	Acetaminophen 160 mg	Pain reliever/fever reducer	Chlorpheniramine maleate 1 mg	Antihistamine	Dextromethorphan HBr 5 mg	Cough suppressant	Phenylephrine HCl 2.5 mg	Nasal decongestant
Weight (lb)	Age (yr)	Dose (mL)																							
48-95	6-11	10 mL																							
36-47	4-5	do not use unless directed by a doctor																							
under 36	under 4	do not use																							
Active ingredients (in each 5 mL)	Purposes																								
Acetaminophen 160 mg	Pain reliever/fever reducer																								
Chlorpheniramine maleate 1 mg	Antihistamine																								
Dextromethorphan HBr 5 mg	Cough suppressant																								
Phenylephrine HCl 2.5 mg	Nasal decongestant																								

BEST CHOICE



0 70038 61689 4

PLD-H253C F0004114

Lot No:  
Exp. Date:

**BEST CHOICE Children's Plus Multi-Symptom Cold**

**CHILDRENS PLUS MULTI SYMPTOM COLD**  
acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride

suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63941-304
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: HBR47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-304-04	1 in 1 BOX	05/31/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	05/31/2015	

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)

Revised: 3/2018

Best Choice (Valu Merchandisers Company)