

CD 5-2- iodine solution Brown Dairy Equipment

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

CD 5-2 Commercial Grade Iodine Pre and Post Dip

ACTIVE INGREDIENT: .5% Titratable Iodine

Contains 2% Skin Conditioner

DIRECTIONS FOR USE

PRE-DIPPING

- Immediately prior to milking, remove any excessive soils from teat surfaces.
- Apply CD 5-2 with dip cup or sprayer covering entire surface of teat.*
- Allow for minimum of 15 seconds contact time.
- Wipe each teat with a single service towel to remove teat dip and soil from teat surface.
- Attach milking unit.

* It is a common practice to "fore-strip" teats prior to milking for purposes of milk let down and mastitis detection. The procedure of fore-stripping may be performed prior to or after application of CD 5-2.

POST-DIPPING

- Immediately after removal of milking unit apply CD 5-2 or CD 1-10 with dip cup or sprayer ensuring complete coverage of teat surfaces with teat dip solution.

DO NOT DILUTE TEAT DIP, USE AT FULL STRENGTH. IF TEAT DIP IN THE TEAT CUP BECOMES COLOUDY OR DIRTY IN APPEARANCE, DISCARD AND REPLACE CONTENTS OF DIP CUP WITH FRESH TEAT DIP. DO NOT RETURN CONTAMINATED TEAT DIP TO THE ORIGINAL CONTAINER.

If original container becomes frozen, thaw completely then mix thoroughly before use.

KEEP OUT OF REACH OF CHILDREN.

PRECAUTIONARY STATEMENTS

DANGER

CAN CAUSE EYE DAMAGE

Protect eyes when handling. DO NOT get in eyes or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

Eyes: In case of contact with eyes, flush immediately with water for at least 15 minutes. Call a physician.

Internal: If swallowed, promptly rinse mouth then give a large quantity of milk or water. Avoid alcohol. Contact a physician immediately. Never give anything by mouth to an unconscious person.

FOR ASSISTANCE WITH MEDICAL EMERGENCY, CALL: 1-800-424-9300

REFER TO SAFETY DATA SHEET. (SDS)

HEALTH 2

FLAMMABILITY 0

REACTIVITY 1

PERSONAL PROTECTION A

Manufactured by: Dairy Dynamics 17820 Washington St. Union, IL 60180



CD 5-2

Commercial Grade Iodine Pre and Post Dip

An aid in reducing the spread of mastitis causing organisms

DIRECTIONS FOR USE

PRE-DIPPING:

- Immediately prior to milking, remove any excessive soils from teat surfaces.
- Apply CD 5-2 with dip cup or sprayer covering entire surface of teat.*
- Allow for minimum of 15 seconds contact time.
- Wipe each teat with a single service towel to remove teat dip and soil from teat surface
- Attach milking unit.

* It is a common practice to "fore-strip" teats prior to milking for purposes of milk let down and mastitis detection. The procedure of fore-stripping may be performed prior to or after application of CD 5-2.

POST-DIPPING:

- Immediately after removal of milking unit apply CD 5-2 or CD 1-10 with dip cup or sprayer ensuring complete coverage of teat surfaces with teat dip solution.

DO NOT DILUTE TEAT DIP. USE AT FULL STRENGTH. IF TEAT DIP IN THE TEAT CUP BECOMES CLOUDY OR DIRTY IN APPEARANCE, DISCARD AND REPLACE CONTENTS OF DIP CUP WITH FRESH TEAT DIP. DO NOT RETURN CONTAMINATED TEAT DIP TO THE ORIGINAL CONTAINER.

If original container becomes frozen, thaw completely then mix thoroughly before use.



Manufactured by: Dairy Dynamics 17820 Washington St. Union, IL 60180

TAKE TIME



OBSERVE LABEL DIRECTIONS

NET CONTENTS: 275 GALLONS

**ACTIVE INGREDIENT: .5% Titratable Iodine
Contains 2% Skin Conditioners**

CD 5-2
Teat Dip Formulation-Commercial
5,000 PPM Iodine (.5% Iodine)
Contains 2% Skin Conditioners

**PRECAUTIONARY STATEMENTS
DANGER**

**KEEP OUT OF REACH OF CHILDREN.
CAN CAUSE EYE DAMAGE.**

Protect eyes when handling. DO NOT get in eyes or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

Eyes: In case of contact with eyes, flush immediately with water for at least 15 minutes. Call a physician.

Internal: If swallowed, promptly rinse mouth then give a large quantity of milk or water. Avoid alcohol. Contact a physician immediately. Never give anything by mouth to an unconscious person.

FOR ASSISTANCE WITH MEDICAL EMERGENCY,

CALL: 1-800-424-9300.

REFER TO SAFETY DATA SHEET. (SDS)

HEALTH	2
FLAMMABILITY	0
REACTIVITY	1
PERSONAL PROTECTION	A



XXX-977-CDS

CD 5-2

iodine solution

Product Information

Product Type

OTC ANIMAL DRUG

Item Code (Source)

NDC:62897-977

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	5 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
C12-15 PARETH-9 (UNII: H3Z1Y6WP1R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
PHOSPHORIC ACID (UNII: E4GA8884NN)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62897-977-55	208.2 L in 1 DRUM		
2	NDC:62897-977-75	1041 L in 1 TANK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/06/2021	

Labeler - Brown Dairy Equipment (017030149)

Establishment

Name	Address	ID/FEI	Business Operations
Dairy Dynamics L. L. C.		142010953	manufacture, pack, label, api manufacture

Revised: 12/2021

Brown Dairy Equipment