

**NAPROXEN SODIUM- naproxen sodium tablet**  
**FREDS INC**

-----  
**ALL DAY PAIN RELIEF**  
**Naproxen Sodium Tablets, USP 220 mg**

**Active ingredient**  
**(in each tablet)**

**Active ingredient**  
**(in each caplet)**

Naproxen sodium 220 mg  
(naproxen 200 mg) (NSAID)\*\*  
\*\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Allergy alert**

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of the reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away

## **Directions**

### **for Tablets**

- **do not take more than directed**
  - **the smallest effective dose should be used**
  - drink a full glass of water with each dose
- Adults and children 12 years and older
- take 1 tablet every 8 to 12 hours while symptoms last
  - for the first dose you may take 2 tablets within the first hour
  - do not exceed 2 tablets in any 8- to 12- hour period
  - do not exceed 3 tablets in a 24- hour period

Children under 12 years

- ask a doctor

### **For Caplets**

- **do not take more than directed**
  - **the smallest effective dose should be used**
  - drink a full glass of water with each dose
- Adults and children 12 years and older
- take 1 caplet every 8 to 12 hours while symptoms last
  - for the first dose you may take 2 caplets within the first hour
  - do not exceed 2 caplets in any 8- to 12- hour period
  - do not exceed 3 caplets in a 24- hour period

Children under 12 years

- ask a doctor

## **Other information**

- **each tablets contains:** sodium 20 mg
- store at 20 - 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

## **Inactive ingredients**

FD&C blue #2 aluminum lake, hypromellose, maize starch, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, titanium dioxide.

## **Questions or comments?**

1-877-770-3183 Mon-Fri 9:00 AM to 4:00 PM EST

## **ALL DAY PAIN RELIEF**

***Naproxen Sodium Tablets, USP 220 mg***

***Pain Reliever/Fever Reducer (NSAID)***

***GLUTEN-FREE***

***STRENGTH TO LAST 12 HOURS***

***24 Tablets***

**Drug Facts (continued)**  
**Other information**  
 ■ each tablet contains sodium 20 mg  
 ■ store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).  
**Inactive ingredients** FD&C blue #2 aluminum lake, hydroxypropyl methylcellulose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, polydioxane K-30, sodium starch glycolate, stearic acid, titanium dioxide  
**Questions or comments?**  
 1-877-770-3183 Mon-Fri 9:00 AM to 4:00 PM EST  
 100% Satisfaction Guarantee  
 Questions or comments: 1-888-451-9920 (2722)  
 DISTRIBUTED BY: Fred's, Inc.  
 4300 NearGulfway Rd., Memphis, TN 38118  
 www.freds.com  
 MADE IN INDIA

**KEEP CAUTION FOR REFERENCE**  
**DO NOT USE IF CARTON IS OPEN OR IF FOIL SEAL ON BOTTLE OPENING IS MISSING OR BROKEN.**

**Drug Facts**  
**Active ingredient (in each tablet)**  
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\* - Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Purpose**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis  
 ■ headache  
 ■ muscle aches  
 ■ menstrual cramps  
 ■ toothache  
 ■ the common cold  
 ■ temporarily reduces fever

**Uses**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ minor pain of arthritis  
 ■ headache  
 ■ muscle aches  
 ■ menstrual cramps  
 ■ toothache  
 ■ the common cold  
 ■ temporarily reduces fever

**Warnings**  
**Allergy alert** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin redness ■ rash ■ dizziness  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems

**fred's Pharmacy**

**ALL DAY PAIN RELIEF**

**Naproxen Sodium Tablets, USP 220 mg** **GLUTEN-FREE**

**Pain Reliever/Fever Reducer (NSAID)** **STRENGTH TO LAST 12 HOURS**

**24 Tablets** **Compare to the Active Ingredient of Aleve® Tablets\***

**ACTUAL SIZE**



**COATING FREE AREA**

Lot Exp.

**31**  
000006 0202

**COATING FREE AREA**

**300000202**  
**531000012**

**COATING FREE AREA**

**Drug Facts (continued)**

■ vomit blood  
 ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ you have difficulty swallowing  
 ■ it feels like the pill is stuck in your throat  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed  
 ■ the smallest effective dose should be used  
 ■ drink a full glass of water with each dose

**Adults and children 12 years and older**  
 ■ take 1 tablet every 8 to 12 hours while symptoms last  
 ■ for the first dose, you may take 2 tablets within the first hour  
 ■ do not exceed 2 tablets in any 8- to 12-hour period  
 ■ do not exceed 3 tablets in a 24-hour period  
 ■ ask a doctor

Children under 12 years

**Drug Facts (continued)**

■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if ■ the stomach bleeding warning applies to you**  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ you are taking a diuretic  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers  
 ■ you have asthma

**Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition**  
 ■ taking any other drug

**When using this product ■ take with food or milk if stomach upset occurs**  
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

**Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:**  
 ■ feel faint

**COATING FREE AREA**

**300000202**  
**531000012**


**COATING FREE AREA**

**COATING FREE AREA**

**300000202**  
**531000012**

**COATING FREE AREA**

**ALL DAY PAIN RELIEF**  
**Naproxen Sodium Tablets, USP 220 mg**  
**GLUTEN-FREE**  
**Pain Reliever/Fever Reducer (NSAID)**  
**STRENGTH TO LAST 12 HOURS**  
**24 Caplets +**  
**+ Capsule-Shaped Tablets**



**ALL DAY PAIN RELIEF**  
**Naproxen Sodium Tablets, USP 220 mg**  
**GLUTEN-FREE**  
**Pain Reliever/Fever Reducer (NSAID)**  
**STRENGTH TO LAST 12 HOURS**  
**50 Caplets +**  
**+ Capsule-Shaped Tablets**

**KEEP CARTON FOR REFERENCE**  
**DO NOT USE IF CARTON IS OPEN OR IF FOIL SEAL ON BOTTLE OPENING IS MISSING OR BROKEN.**

**Drug Facts**  
**Active ingredient Purpose**  
 (in each caplet)  
 Naproxen sodium 220 mg  
 (naproxen 220 mg) (NSAID)\*... Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - minor pain of arthritis
  - backache
  - headache
  - the common cold
  - temporarily reduces fever

**Warnings**  
 Always alert Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 

- hives
- facial swelling
- skin redness
- itching
- asthma (wheezing)
- swelling
- skin redness
- rash
- blistering

 If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warnings:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 

- are age 60 or older
- have had stomach ulcers or bleeding problems

 →

**Drug Facts (continued)**  
**Other information**  
 each tablet contains: sodium 20 mg  
 store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients** FD&C blue #2 aluminum lake, hydroxyethylcellulose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone K-30, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
 1-877-770-3183 Mon-Fri 9:00 AM to 4:00 PM EST

**100%** Satisfaction Guarantee  
 Questions or comments? 1-888-291-7422 (2222)

DISTRIBUTED BY: Fred's, Inc.  
 4300 Newwood Rd, Memphis, TN 38118  
 www.freds.com  
 MADE IN INDIA

**fred's Pharmacy**

**ALL DAY PAIN RELIEF**

**Naproxen Sodium Tablets, USP 220 mg**

**Pain Reliever/Fever Reducer (NSAID)**

**STRENGTH TO LAST 12 HOURS**

**GLUTEN-FREE**

**24 Caplets\***  
 \*Capsule-Shaped Tablets

**Compare to the Active Ingredient of:**  
**Alieve® Caplets®**

**ACTUAL SIZE**

**31**  
 000000  
 0000

**COATING FREE AREA**

**COATING FREE AREA**

Lot Exp.

8457912609 5  
 0

\*The product has not been evaluated by the FDA for safety and effectiveness in children. It is not intended for use in children. Always use as directed. Trade name of Fred's Pharmacy, LLC. P-00

**Drug Facts (continued)**

- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- have gas, worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period
- ask a doctor

Children under 12 years →

**COATING FREE AREA**

300000200  
 531000011

**Drug Facts (continued)**

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.
- do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if in the stomach bleeding warning applies to you**

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition taking any other drug**

When using this product do not take with food or milk if stomach upset occurs

- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel heart

 →

**COATING FREE AREA**

300000200  
 531000011

**COATING FREE AREA**

6 0 4 8 8 0



**Drug Facts (continued)**  
 microcrystalline cellulose, polyethylene glycol, povidone K-30, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
 1-877-770-3183 Mon-Fri 9:00 AM to 4:00 PM EST

\*This product is not manufactured or distributed by Bayer HealthCare, LLC., distributor of Aleve. Aleve is registered Trademark of Bayer HealthCare, LLC.

DISTRIBUTED BY: Fred's, Inc.  
 4300 NEW GETWELL RD,  
 MEMPHIS, TN 38118  
 www.fredsinc.com  
 MADE IN INDIA

**100%** satisfaction guaranteed  
 Questions or comments:  
 1-888-331-FRED (3732)

11-00

**KEEP CARTON FOR REFERENCE**  
**DO NOT USE IF CARTON IS OPEN OR IF FOL SEAL ON BOTTLE OPENING IS MISSING OR BROKEN.**

**Drug Facts**  
**Active ingredient Purpose (In each caplet)**  
 Naproxen sodium 220 mg Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - backache
  - headache
  - the common cold
  - temporarily reduces fever

**Warnings**

**Always alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- itch
- blister

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, →

**fred's Pharmacy**

**ALL DAY PAIN RELIEF**

**Naproxen Sodium Tablets, USP 220 mg** **GLUTEN-FREE**

**Pain Reliever/Fever Reducer (NSAID)** **STRENGTH TO LAST 12 HOURS**

**50 Caplets\*** **Compare to the Active Ingredient of Aleve® Caplets\***

**100%** satisfaction guaranteed  
 Questions or comments:  
 1-888-331-FRED (3732)

**ACTUAL SIZE**



**0 84579 12608 8**

**COATING FREE AREA**

Lot Exp.

**Drug Facts (continued)**

- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

**Adults and children 12 years and older**

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period
- ask a doctor

Children under 12 years

**Other information**

- each caplet contains sodium 20 mg
- store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients:** PEG 400, aluminum lake, hypromellose 2910, maize starch, →

**31**  
0000000000

**COATING FREE AREA**

**Drug Facts (continued)**

- naproxen, or others
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition or taking any other drug.**

**When using this product:**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.


**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days

**COATING FREE AREA**

**300000206**  
**531000014**

**ALL DAY PAIN RELIEF**  
**Naproxen Sodium Tablets, USP 220 mg**  
**GLUTEN-FREE**  
**Pain Reliever/Fever Reducer (NSAID)**  
**STRENGTH TO LAST 12 HOURS**  
**100 Caplets +**  
**+ Capsule-Shaped Tablets**



# NAPROXEN SODIUM

naproxen sodium tablet

## Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:55315-992 |
| Route of Administration | ORAL           |                    |               |

Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength |
|---|-------------------|----------|
| NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ) | NAPROXEN SODIUM   | 220 mg   |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)                  |          |
| FD&C BLUE NO. 2 (UNII: L06K8R7DQK)                       |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)                         |          |
| POVIDONE K30 (UNII: U725QWY32X)                          |          |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                      |          |
| STARCH, CORN (UNII: O8232NY3SJ)                          |          |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)            |          |
| STEARIC ACID (UNII: 4ELV7Z65AP)                          |          |

### Product Characteristics

|          |      |              |          |
|----------|------|--------------|----------|
| Color    | blue | Score        | no score |
| Shape    | OVAL | Size         | 12mm     |
| Flavor   |      | Imprint Code | 220      |
| Contains |      |              |          |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:55315-992-24 | 24 in 1 BOTTLE; Type 0: Not a Combination Product  | 08/17/2017           |                    |
| 2 | NDC:55315-992-05 | 50 in 1 BOTTLE; Type 0: Not a Combination Product  | 08/17/2017           |                    |
| 3 | NDC:55315-992-10 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 08/17/2017           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA091353                               | 08/17/2017           |                    |

## NAPROXEN SODIUM

naproxen sodium tablet

### Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:55315-991 |
| Route of Administration | ORAL           |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength |
|---|-------------------|----------|
| NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ) | NAPROXEN SODIUM   | 220 mg   |



## Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                      |          |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)            |          |
| STEARIC ACID (UNII: 4ELV7Z65AP)                          |          |
| FD&C BLUE NO. 2 (UNII: L06K8R7DQK)                       |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)                         |          |
| STARCH, CORN (UNII: O8232NY3SJ)                          |          |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)                  |          |
| POVIDONE K30 (UNII: U725QWY32X)                          |          |

## Product Characteristics

|          |       |              |          |
|----------|-------|--------------|----------|
| Color    | blue  | Score        | no score |
| Shape    | ROUND | Size         | 9mm      |
| Flavor   |       | Imprint Code | I1       |
| Contains |       |              |          |

## Packaging

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:55315-991-24 | 24 in 1 BOTTLE; Type 0: Not a Combination Product | 08/17/2017           |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA091353                               | 08/17/2017           |                    |

**Labeler** - FREDS INC (005866116)

Revised: 7/2017

FREDS INC