

CAREONE SEVERE COLD- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated
American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

American Sales Company Severe Cold Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
- nasal congestion
- headache
- cough
- minor aches and pains
- sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 10 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- **each caplet contains:** sodium 4 mg
- store at 20-25°C (68-77°F)
- do not use if blister unit is broken or torn

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to the active ingredients in Mucinex[®] Fast-Max[®] Cold & Flu

SEVERE COLD

Pain Reliever/Fever Reducer-Acetaminophen

Cough Suppressant-Dextromethorphan HBr

Expectorant-Guaifenesin

Nasal Decongestant-Phenylephrine HCl

Maximum Strength

Relieves Aches, Fever & Sore Throat

Controls Cough

Relieves Nasal & Chest Congestion

Thins & Loosens Mucus

For Ages 12+

All in one Relief

Gluten Free

Actual Size

OUR PHARMACISTS RECOMMEND

20 CAPLETS

CAREone®

NDC 41520-773-01

Compare to the active ingredients
in Mucinex® Fast-Max® Cold & Flu*

SEVERE COLD

Pain Reliever/Fever Reducer-Acetaminophen
Cough Suppressant-Dextromethorphan HBr
Expectorant-Guaifenesin
Nasal Decongestant-Phenylephrine HCl

Maximum Strength**

Relieves Aches, Fever & Sore Throat
Controls Cough
Relieves Nasal & Chest Congestion
Thins & Loosens Mucus

For Ages 12+

All in one Relief***

Gluten Free




Actual Size



20 CAPLETS

922C9 OF SF2



PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Keep outer package for full information.

Lift here to open

DISTRIBUTED BY FOODHOLD U.S.A., LLC
LANDOVER, MD 20785 · 1-877-846-9949
©2019 S&S Brands, LLC
Quality guaranteed or your money back.

*Per 4-hour dose, dose every 4 hours
***Helps to relieve these symptoms day or night.

*This product is not manufactured or distributed by Recitt Benelux, distributor of Mucinex® Fast-Max® Cold & Flu.

3 41520 00063 8

<p>Drug Facts</p> <p>Active Ingredients (in each caplet) Purposes</p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p>Uses</p> <p>Temporarily relieves these common cold and flu symptoms:</p> <ul style="list-style-type: none"> nasal congestion headache cough minor aches and pains sore throat sinus congestion and pressure temporarily reduces fever helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness blisters itching swelling <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease heart disease diabetes 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> high blood pressure thyroid disease trouble urinating due to an enlarged prostate gland persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>When using this product do not use more than directed</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur pain, nasal congestion or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed (see Overdose warning) do not take more than 10 caplets in any 24-hour period adults and children 12 years of age and over: take 2 caplets every 4 hours children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> each caplet contains sodium 4 mg store at 20-25°C (68-77°F) do not use if blister unit is broken or torn <p>Inactive ingredients: croscarmellose sodium, croscopolone, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, talc, titanium dioxide</p> <p>Questions or comments? 1-800-719-0260</p>
---	---

922C9 OF SB2

CAREONE SEVERE COLD

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-773
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	L922
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-773-01	10 in 1 CARTON	04/07/2016	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/07/2016	

Labeler - American Sales Company (809183973)

Revised: 8/2019

American Sales Company