


BENZONATATE - benzonatate capsule
Ascend Laboratories, LLC

Benzonatate Capsules, USP
100 mg, 150 mg and 200 mg

DESCRIPTION

Benzonatate, a non-narcotic oral antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26-nonaoxaocacosan-28-yl p-(butylamino) benzoate; with a molecular weight of 603.7.



Each soft gelatin capsule, for oral administration, contains 100 mg, 150 mg or 200 mg of benzonatate USP. Benzonatate Capsules, USP also contain the following inactive ingredients: D&C Yellow #10, gelatin, glycerin, purified water, methylparaben and propylparaben.

CLINICAL PHARMACOLOGY

Benzonatate acts peripherally by anesthetizing the stretch receptors located in the respiratory passages, lungs, and pleura by dampening their activity and thereby reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect lasts for 3 to 8 hours. Benzonatate has no inhibitory effect on the respiratory center in recommended dosage.

INDICATIONS & USAGE

Benzonatate USP is indicated for the symptomatic relief of cough.

CONTRAINDICATIONS

Hypersensitivity to benzonatate or related compounds.

WARNINGS

Hypersensitivity

Severe hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. Severe reactions have required intervention with vasopressor agents and supportive measures.

Psychiatric Effects

Isolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also been reported in patients taking benzonatate in combination with other prescribed drugs.

Accidental Ingestion and Death in Children

Keep benzonatate capsules out of reach of children. Accidental ingestion of benzonatate resulting in death has been reported in children below age 10. Signs and symptoms of overdose have been reported within 15 to 20 minutes and death has been reported within one hour of ingestion. If accidental ingestion occurs, seek medical attention immediately (see **OVERDOSAGE**).

PRECAUTIONS

Benzonatate is chemically related to anesthetic agents of the para-amino-benzoic acid class (e.g. procaine; tetracaine) and has been associated with adverse CNS effects possibly related to a prior sensitivity to related agents or interaction with concomitant medication.

Information for Patients Swallow benzonatate capsules whole. Do not break, chew, dissolve, cut, or crush Benzonatate Capsules. Release of benzonatate from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. If numbness or tingling of the tongue, mouth, throat, or face occurs, refrain from oral ingestion of food or liquids until the numbness has resolved. If the symptoms worsen or persist, seek medical attention.

Keep benzonatate out of reach of children. Accidental ingestion resulting in death has been reported in children. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. Signs and symptoms may include restlessness, tremors, convulsions, coma and cardiac arrest. If accidental ingestion occurs, seek medical attention immediately. Overdosage resulting in death may occur in adults. Do not exceed a single dose of 200 mg and a total daily dosage of 600 mg. If you miss a dose of benzonatate capsule, skip that dose and take the next dose at the next scheduled time. Do not take 2 doses of benzonatate at one time.

Usage in Pregnancy

PREGNANCY CATEGORY C

Animal reproduction studies have not been conducted with benzonatate. It is also not known whether benzonatate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Benzonatate should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when benzonatate is administered to a nursing woman.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity, mutagenicity, and reproduction studies have not been conducted with benzonatate.

Pediatric Use

Safety and effectiveness in children below the age of 10 have not been established. Accidental ingestion resulting in death has been reported in children below age 10. Keep out of reach of children.

ADVERSE REACTIONS

Potential Adverse Reactions to benzonatate may include:

Hypersensitivity reactions including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

CNS: sedation; headache; dizziness; mental confusion; visual hallucinations.

GI: constipation; nausea; GI upset.

Dermatologic: pruritus; skin eruptions.

Other: nasal congestion; sensation of burning in the eyes; vague “chilly” sensation; numbness of the chest; hypersensitivity. Deliberate or accidental overdose has resulted in death, particularly in children.

OVERDOSAGE

Intentional and unintentional overdose may result in death, particularly in children. The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

Signs and Symptoms

The signs and symptoms of overdose of benzonatate have been reported within 15 to 20 minutes. If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly, which may cause choking and airway compromise. CNS stimulation may cause restlessness and tremors which may proceed to clonic convulsions followed

by profound CNS depression. Convulsions, coma, cerebral edema and cardiac arrest leading to death have been reported within 1 hour of ingestion.

Treatment

In case of overdose, seek medical attention immediately. Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate special attention to protection against aspiration of gastric contents and orally administered materials. Convulsions should be treated with a short-acting barbiturate given intravenously and carefully titrated for the smallest effective dosage. Intensive support of respiration and cardiovascular-renal function is an essential feature of the treatment of severe intoxication from overdosage. Do not use CNS stimulants.

DOSAGE & ADMINISTRATION

Adults and Children over 10 years of age: Usual dose is one 100 mg, 150 mg or 200 mg capsule three times a day as needed for cough. If necessary to control cough, up to 600 mg daily in three divided doses may be given. **Benzonatate should be swallowed whole.** Benzonatate capsules are not to be broken, chewed, dissolved, cut or crushed.

HOW SUPPLIED

Benzonatate Capsules USP, 100 mg: Yellow soft gelatin capsules, imprinted "105", available in bottles of 100's (NDC 67877-573-01), and 500's (NDC 67877-573-05).

Benzonatate Capsules USP, 150 mg: Yellow soft gelatin capsules, imprinted "128", available in bottles of 100 (NDC 67877-574-01).

Benzonatate Capsules USP, 200 mg: Yellow soft gelatin capsules, imprinted "106", available in bottles of 100's (NDC 67877-575-01) and 500's (NDC 67877-575-05). Store at 20°C to 25° C (68°F to 77°F). [See USP Controlled Room Temperature].

PROTECT FROM LIGHT.

Manufactured by:

Softgel Healthcare Pvt. Ltd.,
Survey No. 20/1,
Vandalur-Kelambakkam Road,
Pudupakkam, Chengalpattu,
Tamilnadu - 603 103, India.

Distributed by:

Ascend Laboratories, LLC
Bedminster, NJ 07921

Revised: November 2025

PT3732-01 (2411326)

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL 100 mg

NDC 67877-573-05

Benzonatate Capsules, USP

100 mg

500 Capsules

Each soft gelatin capsule contains:
Benzonatate, USP 100 mg

USUAL DOSAGE: See package insert.

Information for Patients: Swallow benzonatate capsules whole. Do not break, chew, dissolve, cut, or crush benzonatate capsules USP.

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP.

KEEP TIGHTLY CLOSED

Manufactured by: Softgel Healthcare Pvt. Ltd.,
Survey No. 20/1, Vandalur-Kelambakkam Road,
Pudupakkam, Chengalpattu,
Tamilnadu - 603 103, India.

Distributed by: Ascend Laboratories, LLC
Bedminster, NJ 07921

ASCEND
Laboratories, LLC

NDC 67877-573-05

**Benzonatate
Capsules, USP**

100 mg

PROTECT FROM LIGHT

Rx Only 500 Capsules

M.L. No.: TN00002123
PL1423-01 (2620548)

Unvarnished area
52.5 x 27.5 mm (L x H)
Rest label should be
with UV Varnish

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL 200 mg

NDC 67877-575-01

Benzonatate Capsules, USP

200 mg

100 Capsules

Each soft gelatin capsule contains:
Benzonatate, USP 200 mg

USUAL DOSAGE: See package insert.

Information for Patients: Swallow benzonatate capsules whole. Do not break, chew, dissolve, cut, or crush benzonatate capsules USP.

Store at 20°C to 25°C (68°F to 77°F)
[See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP.

KEEP TIGHTLY CLOSED

Manufactured by: Softgel Healthcare Pvt. Ltd.,
Survey No. 20/1, Vandalur-Kelambakkam Road,
Pudupakkam, Chengalpattu,
Tamilnadu - 603 103, India.

Distributed by: Ascend Laboratories, LLC
Bedminster, NJ 07921

ASCEND
Laboratories, LLC

NDC 67877-575-01

**Benzonatate
Capsules, USP**

200 mg

PROTECT FROM LIGHT

Rx Only 100 Capsules

M.L. No.: TN00002123
PL1424-01 (2620549)

Unvarnished area
32.5 x 22 mm (LxH)
Rest label should be
with UV Varnish

BENZONATATE

benzonatate capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67877-573
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZONATATE (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR)	BENZONATATE	100 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	105
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67877-573-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2017	
2	NDC:67877-573-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040627	03/22/2017	

BENZONATATE

benzonatate capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67877-575
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZONATATE (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR)	BENZONATATE	200 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	106
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67877-575-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2017	
2	NDC:67877-575-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040627	03/22/2017	

Labeler - Ascend Laboratories, LLC (141250469)

Establishment

Name	Address	ID/FEI	Business Operations
Softgel Healthcare Pvt. Ltd.		676666501	ANALYSIS(67877-573, 67877-575) , LABEL(67877-573, 67877-575) , MANUFACTURE(67877-573, 67877-575) , PACK(67877-573, 67877-575)

