

JILGYUNGYI- glucose, sodium chloride tablet
JILGYUNGYI CO.,LTD.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ACTIVE INGREDIENT

Active Ingredient: Glucose 50.97%, Sodium Chloride 35.50%

INACTIVE INGREDIENT

Inactive Ingredients:

Butylene glycol, Water, Allantoin, Opuntia ficus-indica extract, Centella asiatica extract, Aloe barbadensis leaf extract, Gingko biloba leaf extract, Mentha piperita(peppermint) leaf extract, Sophora angustifolia root extract, Angelica gigas root extract, 1,2-Hexanediol, Citric acid, Magnesium stearate, Soluble collagen, Sodium citrate

PURPOSE

Purpose: Odor care, Prevent vaginitis

WARNINGS

Warnings:

For vaginal use only

Do not use if you have (are)

- an allergic reaction (rashes, and pimples, redness, swelling, skin irritation, discomfort, burning, vaginal peeling or bleeding, pain in the abdomen or pelvic area, fluttering heartbeat and chest pain, lumbar pain, abnormal vaginal secretion, pollakisuria, abnormal dryness, excessive menstrual bleeding, odor and green colored vaginal secretion, hypotension, faint, and papulopustule, pelvic pain)
- pregnant
- hyper sensitive skin and skin ailment
- scars on vagina
- itching, rash, swelling, redness, discomfort, burning, irritation vaginal peeling or bleeding

When using this product

- do not use while having the monthly menstrual period
- prior to absorption there is a risk the tablet may fall out of the vagina
- often wash towels or underwear and do not wear reuse underwear again prior to washing due to risk of infections
- try to avoid sexual relations when you have vaginal yeast infection, it can be aggravated
- try not to use tampons and feminine cleansers

- avoid contact with eye area
 - do not eat or swallow
 - do not use rubber products such as latex, condoms, and diaphragms together
 - do not use other vaginal tablets while using Jil Gyung Yi
- If pregnant or breast-feeding, do not use.

Store at cool temperature and dry place with a closed container. Avoid direct sunlight

Overdose warning

- If you accidentally take the tablet orally, stop using this medication and see doctor immediately.

Stop use and ask a doctor if you have

- fever (over 38°C), pain in abdomen, odor, lumbar pain, abnormal secretion, nausea and abnormal shoulder pain.

KEEP OUT OF REACH OF CHILDREN

KEEP OUT OF REACH OF CHILDREN

INDICATIONS & USAGE

Uses:

Moisturizes vaginal dryness and helps to improve elasticity

Relieves external itching and alleviates odors before and after the menstruation

DOSAGE & ADMINISTRATION

Directions:

For adults – 2 tablets at the same time

- For a first-timer, keep use 2 days consecutively and after that, use once a week.
- Do not use up to 6 hours prior to any sexual relation.
- Do not use up to 6 months after giving birth.
- Symptoms may vary depending on the age of the patient.

Directions

- Use only adults and children 12 years age and over

1) Wash hands before using this product

2) Open the package and vaginal tablets are placed into the vagina in much the same way as a tampon stand, squat, or lie on your back in a comfortable position then insert the tablets into the vagina as far as it will comfortably go.

3) It is recommended to use the treatment during night time prior to bed.

4) Use of a sanitary pad after application, because there may be vaginal secretion 30min after application. If there are lots of secretion, change a new sanitary pad.

5) Wash the area around the vagina (the vulva) gently next day, bacteria in your vaginal secretion can cause itching.

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - JILGYUNGYI 6 Tablets

Inactive ingredients

Butylene glycol, Water, Allantoin, Opuntia ficus-indica extract, Centella asiatica extract, Aloe barbadensis leaf extract, Ginkgo biloba leaf extract, Mentha piperita (peppermint) leaf extract, Sophora angustifolia root extract, Angelica gigas root extract, 1,2-Hexanediol, Citric acid, Magnesium stearate, Soluble collagen, Sodium citrate

Manufactured by JILGYUNGYI CO., LTD.

3rd Floor, 310, 40, Imhro, Uiwang-si, Gyeonggi-do, Rep. of Korea
11-8, Banpo-daero 8-gil, Seocho-gu, Seoul, Republic of Korea

Jil Gyung Yi
Feminine Cleanser



Warnings

For vaginal use only
Do not use if you have (are):
- an allergic reaction (rashes, and pimples, redness, swelling, skin irritation, discomfort, burning, vaginal peeling or bleeding, pain in the abdomen or pelvic area, fluttering heartbeat and chest pain, lumbar pain, abnormal vaginal secretion, pollakisuria, abnormal dryness, excessive menstrual bleeding, odor and green colored vaginal secretion, hypotension, faint, and papulopustule, pelvic pain)
- Pregnant or breast-feeding
- hyper sensitive skin and skin ailment
- scars on vagina
- itching, rash, swelling, redness, discomfort, burning, irritation vaginal peeling or bleeding
When using this product:
- do not use while having the monthly menstrual period
- prior to absorption there is a risk the tablet may fall out of the vagina
- often wash towels or underwear and do not wear reuse underwear again prior to washing due to risk of infections
- try to avoid sexual relations when you have vaginal yeast infection, it can be aggravated
- try not to use tampons and feminine cleansers
- avoid contact with eye area
- do not eat or swallow
- do not use rubber products such as latex, condoms, and diaphragms together
- do not use other vaginal tablets while using Jil Gyung Yi

Store at cool temperature and dry place with a closed container.
- Avoid direct sunlight

Overdose warning
- If you accidentally take the tablet orally, stop using this medication and see doctor immediately.

Stop use and ask a doctor if you have
- fever (over 38°C), pain in abdomen, odor, lumbar pain, abnormal secretion, nausea and abnormal shoulder pain.

Directions
- For adults, use 2 tablets at the same time
- For a first-timer, keep use 2 days consecutively and after that, use once a week.
- Do not use up to 6 hours prior to any sexual relation.
- Do not use up to 6 months after giving birth.
- Symptoms may vary depending on the age of the patient.

Instruction:
- Use only adults and children 12 years age and over
1) Wash hands before using this product
2) Open the package and vaginal tablets are placed into the vagina in much the same way as a tampon stand, squat, or lie on your back in a comfortable position then insert the tablets into the vagina as far as it will comfortably go.
3) It is recommended to use the treatment during night time prior to bed.
4) Use of a sanitary pad after application, because there may be vaginal secretion 30min after application. If there are lots of secretion, change a new sanitary pad.
5) Wash the area around the vagina (the vulva) gently next day, bacteria in your vaginal secretion can cause itching.



| Batch number |
| E X P |

6 tablets

Relieves external itching and alleviates odors before and after the menstruation

Moisturizes vaginal dryness and helps to improve elasticity

Prevent vaginitis

Odor care

Sodium Chloride 35.50%

Glucose 50.97%

Active ingredients

Purpose

600mg / tablet

Jil Gyung Yi Feminine cleanser

Jil Gyung Yi
Feminine Cleanser

Jil Gyung Yi
Feminine Cleanser

Jil Gyung Yi
Feminine Cleanser

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - JILGYUNGYI 10 Tablets

Inactive ingredients

Butylene glycol, Water, Allantoin, Opuntia ficus-indica extract, Centella asiatica extract, Aloe barbadensis leaf extract, Ginkgo biloba leaf extract, Mentha piperita (peppermint) leaf extract, Sophora angustifolia root extract, Angelica gigas root extract, 1,2-Hexanediol, Citric acid, Magnesium stearate, Soluble collagen, Sodium citrate

Manufactured by JILGYUNGYI CO., LTD.

3rd Floor, 310, 40, Imi-ro, Uiwang-si, Gyeonggi-do, Rep. of Korea
11-8, Banpo-daero 8-gil, Seocho-gu, Seoul, Republic of Korea

JilGyungYi

Feminine Cleanser

Warnings

For vaginal use only

Do not use if you have (are):

- an allergic reaction (rashes, and pimples, redness, swelling, skin irritation, discomfort, burning, vaginal peeling or bleeding, pain in the abdomen or pelvic area, fluttering heartbeat and chest pain, lumbar pain, abnormal vaginal secretion, polyakisuria, abnormal dryness, excessive menstrual bleeding, odor and green colored vaginal secretion, hypotension, faint, and papulopustule, pelvic pain)
- Pregnant or breast-feeding
- hyper sensitive skin and skin ailment
- scars on vagina
- itching, rash, swelling, redness, discomfort, burning, irritation vaginal peeling or bleeding

When using this product:

- do not use while having the monthly menstrual period
- prior to absorption there is a risk the tablet may fall out of the vagina
- often wash towels or underwear and do not wear reuse underwear again prior to washing due to risk of infections
- try to avoid sexual relations when you have vaginal yeast infection, it can be aggravated
- try not to use tampons and feminine cleansers
- avoid contact with eye area
- do not eat or swallow
- do not use rubber products such as latex, condoms, and diaphragms together
- do not use other vaginal tablets while using Jil Gyung Yi

Store at cool temperature and dry place with a closed container.

- Avoid direct sunlight

Overdose warning

- If you accidentally take the tablet orally, stop using this medication and see doctor immediately.

Stop use and ask a doctor if you have

- fever (over 38°C), pain in abdomen, odor, lumbar pain, abnormal secretion, nausea and abnormal shoulder pain.

Directions

- For adults, use 2 tablets at the same time

- For a first-timer, keep use 2 days consecutively and after that, use once a week.

- Do not use up to 6 hours prior to any sexual relation.

- Do not use up to 6 months after giving birth.

- Symptoms may vary depending on the age of the patient.

Instruction:

- Use only adults and children 12 years age and over
- 1) Wash hands before using this product
- 2) Open the package and vaginal tablets are placed into the vagina in much the same way as a tampon stand, squat, or lie on your back in a comfortable position then insert the tablets into the vagina as far as it will comfortably go.
- 3) It is recommended to use the treatment during night time prior to bed.
- 4) Use of a sanitary pad after application, because there may be vaginal secretion 30min after application. If there are lots of secretion, change a new sanitary pad.
- 5) Wash the area around the vagina (the vulva) gently next day, bacteria in your vaginal secretion can cause itching.



| Batch number |
| E X P |

10 tablets

Relieves external itching and helps to improve elasticity
Moisturizes vaginal dryness and helps to improve elasticity
Relieves external itching and alleviates odors before and after the menstruation

Drug Facts
Active ingredients
Glucose 50.97%
Sodium Chloride 35.50%
Purpose
Odor care
Prevent vaginitis

Jil Gyung Yi Feminine cleanser 600mg / tablet

JilGyungYi

Feminine Cleanser

JilGyungYi
Feminine Cleanser

JilGyungYi
Feminine Cleanser

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - JILGYUNGYI 20 Tablets

JilGyungYi
Feminine Cleanser

JilGyungYi

Feminine Cleanser

Jil Gyung Yi Feminine cleanser

600mg / tablet

Drug Facts

Active ingredients

Glucose 50.97%

Sodium Chloride 35.50%

Uses

Moisturizes vaginal dryness and helps to improve elasticity
Relieves external itching and alleviates odors before and after the menstruation

Purpose

Odor care

Prevent vaginitis

20 tablets

JilGyungYi
Feminine Cleanser

| Batch number |

| EXP |



Warnings
For vaginal use only (are):
Do not use if you have (are):
- an allergic reaction (rash, hives, and pimples, redness, swelling, skin irritation, discomfort, burning, vaginal peeling or bleeding, pain in the abdomen or pelvic area, fluttering heartbeat and chest pain, lumbar pain, abnormal vaginal secretion, pollektisura, abnormal dryness, excessive menstrual bleeding, odor and green color);
- frequent or breast-feeding vaginal secretion, hypotension, faint, and papulopustule, pelvic pain);
- hyper sensitive skin and skin ailment
- scars on vagina
- itching, rash, swelling, redness, discomfort, burning, irritation
When using this product:
- do not use while having the monthly menstrual period
- prior to absorption there is a risk the tablet may fall out of the vagina
- often wash towels or underwear and do not wear reuse underwear
- again prior to washing due to risk of infections
- try to avoid sexual relations when you have vaginal yeast infection, it can be aggravated
- do not use tampons and feminine cleansers
- avoid contact with eye area
- do not eat or swallow
- do not use rubber products such as latex, condoms, and diaphragms together
- do not use other vaginal tablets while using Jil Gyung Yi

Directions
- For adults, use 2 tablets at the same time
- For a first-time, keep use 2 days consecutively and after that, use once a week
- Do not use up to 6 hours prior to any sexual relation
- Do not use up to 6 months after giving birth.
- Symptoms may vary depending on the age of the patient.
Instruction:
- Use only adults and children 12 years age and over
- Wash hands before using this product
- Open the package and vaginal tablets are placed into the vagina in much the same way as a tampon stand, squat, or lie on your back in a comfortable position then insert the tablets into the vagina as far as it will comfortably go.
- If recommended to use the treatment during night time prior to bed.
- Use of a sanitary pad after application, because there may be vaginal secretion (lump) after application, if there are lots of secretion, change a new sanitary pad.
- Wash the area around the vagina (the vulva) gently next day.
- bacteria in your vaginal secretion can cause itching.

Overdose warning
- Avoid direct sunlight
- If you accidentally take the tablet orally, stop using this medication and see doctor immediately.
Stop use and ask a doctor if you have
- fever (over 38°C), pain in abdomen, odor, lumbar pain, abnormal secretion, nausea and abnormal shoulder pain.

Store at cool temperature and dry place with a closed container.
Store at cool temperature and dry place with a closed container.

JilGyungYi
Feminine Cleanser

Inactive ingredients
Butylene glycol, Water, Allantoin, Opuntia ficus-indica extract, Centella asiatica extract, Aloe barbadensis leaf extract, Ginkgo biloba leaf extract, Mentha piperita (peppermint) leaf extract, Sophora angustifolia root extract, Angelica gigas root extract, 1,2-hexanediol, Citric acid, Magnesium stearate, Soluble collagen, Sodium citrate

Manufactured by JILGYUNGYI CO., LTD.
3rd Floor, 310, 40, Imho, Uiwang-si, Gyeonggi-do, Rep. of Korea
11-8, Banpo-dae-ro 8-gil, Seocho-gu, Seoul, Republic of Korea

JILGYUNGYI

glucose, sodium chloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73004-010
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ANHYDROUS DEXTROSE (UNII: 5SLOG7R0OK) (ANHYDROUS DEXTROSE - UNII:5SLOG7R0OK)	ANHYDROUS DEXTROSE	305.82 mg in 600 mg
Sodium Chloride (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)	Sodium Chloride	213 mg in 600 mg

Inactive Ingredients

Ingredient Name	Strength
Butylene glycol (UNII: 3XUS85K0RA)	
Water (UNII: 059QF0K00R)	
Allantoin (UNII: 344S277G0Z)	

Product Characteristics

Color	white	Score	no score
Shape	capsule	Size	15mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73004-010-02	6 in 1 CARTON	03/01/2019	
1	NDC:73004-010-01	600 mg in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:73004-010-03	10 in 1 CARTON	03/01/2019	
2	NDC:73004-010-01	600 mg in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:73004-010-04	20 in 1 CARTON	03/01/2019	
3	NDC:73004-010-01	600 mg in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/01/2019	

Labeler - JILGYUNGYI CO.,LTD. (689204202)**Registrant** - JILGYUNGYI CO.,LTD. (689204202)

Establishment

Name	Address	ID/FEI	Business Operations
JILGYUNGYI CO.,LTD.		689204202	manufacture(73004-010)

Revised: 11/2021

JILGYUNGYI CO.,LTD.