

LOPERAMIDE HCL- loperamide hcl tablet
L.N.K. International, Inc.

Quality Plus 44-375 Loperamide HCl

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

**QUALITY
+PLUS**

NDC 50844-735-08

*Compare to active ingredient
in Imodium® A-D

LOPERAMIDE HCl
TABLETS, 2 mg

ANTI-DIARRHEAL

Controls the symptoms of diarrhea

24 Caplets

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

1. To open, tear
at perforations and
remove a section.

2. Use ✂ to cut
through plastic
for caplet.

*This product is not manufactured or distributed by Johnson
& Johnson Corporation, owner of the registered trademark
Imodium® A-D. 50844 ORG061937508

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

QUALITY PLUS
LOPERAMIDE HCl
TABLETS, 2 mg


QUALITY PLUS

*Compare to active ingredient in Imodium® A-D

LOPERAMIDE HCl
TABLETS, 2 mg
ANTI-DIARRHEAL

Controls the symptoms of diarrhea

NDC 50844-735-08



3 5084473508 9

**DO NOT USE IF PACKAGE IS
 ER UNIT IS TORN, BROKEN
 SIGNS OF TAMPERING**

o varnish
 p date

Drug Facts

Active ingredient (in each caplet) Loperamide HCl 2 mg. **Purpose** Anti-diarrheal

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death.

Drug Facts (continued)

Do not use if you have bloody or black stool.

Ask a doctor before use if you have a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if symptoms get worse or diarrhea lasts for more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over
 loose stool but no more than 4 caplets in 24 hours

children 9-11 years (60-95 lbs)
 1 caplet after the first loose stool, 1/2 caplet after each subsequent loose stool but no more than 3 caplets in 24 hours

children 6-8 years (48-59 lbs)
 1 caplet after the first loose stool, 1/2 caplet after each subsequent loose stool but no more than 2 caplets in 24 hours

children 2-5 years (34-47 lbs)
 ask a doctor

children under 2 years (up to 33 lbs)
 do not use

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ store between 20°-25°C (68°-77°F)
 ■ see end flap for expiration date and lot number

Inactive ingredients corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday



Quality Plus 44-375

LOPERAMIDE HCL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-735
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-735-24	1 in 1 CARTON	05/03/2005	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:50844-735-45	1 in 1 CARTON	05/03/2005	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:50844-735-19	2 in 1 CARTON	05/03/2005	
3		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:50844-735-02	2 in 1 CARTON	05/03/2005	
4		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:50844-735-44	3 in 1 CARTON	05/03/2005	
5		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:50844-735-08	4 in 1 CARTON	05/03/2005	
6		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

--	--	--	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-735) , pack(50844-735)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-735)

Revised: 7/2025

L.N.K. International, Inc.