# ACETAMINOPHEN- acetaminophen tablet Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# **Major Pharmaceuticals Acetaminophen Drug Facts**

# **Active ingredient (in each tablet)**

Acetaminophen 325 mg

# **Purpose**

Pain reliever/fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

# **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

# Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if the user has ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if the user

- has liver disease
- is a child with pain of arthritis

# Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin

# Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

# do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul> <li>take 2 tablets every 4 to 6 hours while symptoms last</li> <li>do not take more than 10 tablets in 24 hours</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul>
children 6-11 years	<ul> <li>take 1 tablet every 4 to 6 hours while symptoms last</li> <li>do not take more than 5 tablets in 24 hours</li> <li>do not use for more than 5 days unless directed by a doctor</li> </ul>
children under 6 years	ask a doctor

## **Inactive ingredients**

croscarmellose sodium\*, povidone, pregelatinized starch, stearic acid

<sup>\*</sup>may contain this ingredient

#### Questions or comments?

1-800-616-2471

#### **Principal Display Panel**

Compare to the active ingredient in Regular Strength Tylenol® Tablets

Acetaminophen

Pain Reliever/Fever Reducer

Aspirin-Free

Regular Strength

**TABLETS** 

50 ACETAMINOPHEN TABLETS – 325 mg. EACH



acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of

- acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours
- taken with other drugs containing acetam inophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening blisters rash
- If a skin reaction occurs, stop use and seek medical help right away.

- (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose
warning: In case of overdose, get medical help or
contact a Poison Control Center right away
(1-800-222-1222). Quick medical attention is
critical for adults as well as for children even if you
do not notice any signs or symptoms.

oo not take more than
5 tablets in 24 hours
■ do not use for more than
5 days unless directed by
a doctor

c hildren under 6 years ask a doctor

Inactive ingredients croscarmellose sodium\*, povidone, pregelatinized starch, stearic acid \*may contain this ingredient

Questions or comments? 1-800-616-2471



40371 5C C2

# **ACETAMINOPHEN**

acetaminophen tablet

# **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0904-6719

Route of Administration ORAL

# **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients		
Ingredient Name	Strength	
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)		
PO VIDO NE (UNII: FZ989GH94E)		
STEARIC ACID (UNII: 4ELV7Z65AP)		

Product Characteristics			
Color	WHITE	Score	no score
Shape	ROUND (beveled edge)	Size	10 mm
Flavor		Imprint Code	325MG;L403
Contains			

P	Packaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0904-6719-50	1 in 1 CARTON	10/18/2018	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0904-6719-60	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/04/2018	
3	NDC:0904-6719-80	1000 in 1 BOTTLE; Type 0: Not a Combination Product	12/31/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	10/18/2018	

# Labeler - Major Pharmaceuticals (191427277)

Revised: 1/2019 Major Pharmaceuticals