# ACID REDUCER PLUS ANTACID- famotidine, calcium carbonate and magnesium hydroxide tablet, chewable Rite Aid Corporation

-----

#### Rite Aid Corporation Acid Reducer Plus Antacid Drug Facts

#### Active ingredients (in each chewable tablet)

Famotidine 10 mg

Calcium carbonate 800 mg

Magnesium hydroxide 165 mg

#### **Purposes**

Acid reducer

Antacid

#### Use

relieves heartburn associated with acid indigestion and sour stomach

#### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

#### Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

#### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

#### Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

#### Stop use and ask a doctor if

• your heartburn continues or worsens

• you need to take this product for more than 14 days

#### If pregnant or breast-feeding,

ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- adults and children 12 years and over:
- do not swallow tablet whole: chew completely
- to relieve symptoms, **chew** 1 tablet before swallowing
- do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

#### Other information

- each tablet contains: calcium 330 mg; magnesium 80 mg
- Phenylketonurics: Contains Phenylalanine 2 mg per tablet
- read the directions and warnings before use
- read the bottle label. It contains important information.
- store at  $20-25^{\circ}$ C ( $68-77^{\circ}$ F)
- protect from moisture

#### **Inactive ingredients**

anhydrous lactose, artificial berry flavor, aspartame, D&C red no. 7 calcium lake, dextrates, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyacrylate dispersion, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, talc

#### Questions or comments?

1-800-719-9260

#### **Principal Display Panel**

Compare to the active ingredients of Pepcid Complete®

ACID REDUCER PLUS ANTACID

famotidine 10 mg / calcium carbonate 800 mg / magnesium hydroxide 165 mg tablets (chewable) acid reducer plus antacid

**DUAL ACTION** 

Just one tablet relieves heartburn due to acid indigestion

ACTUAL SIZE

BERRY FLAVOR



#### DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING Drug Facts Purposes Active ingredients (in each chewable tablet) Famotidine 10 mg.....Acid reducer Calcium carbonate 800 mg......Antacid Magnesium hydroxide 165 mg......Antacid **Use** relieves heartburn associated with acid indigestion and sour stomach Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional

Keep out of reach of children. In case of overdose, get

medical help or

contact a Poison Control Center

right away

before use.

### ADHESINE BAR - NO COPY Drug Facts (continued) Directions ■ adults and children 12 years and over: do not swallow tablet whole: chew completely ■ to relieve symptoms, chew 1 tablet before swallowing do not use more than 2 chewable tablets in 24 hours children under 12 years: ask a doctor Other information each tablet contains: calcium 330 mg; magnesium 80 mg Phenylketonurics: Contains Phenylalanine 2 mg per tablet read the directions and warnings before use ■ read the bottle label. It contains important information. store at 20-25°C (68-77°F) protect from moisture

#### Drug Facts (continued)

Inactive ingredients anhydrous lactose, artificial berry flavor, aspartame, D&C red no. 7 calcium lake, dextrates, FD&C blue no. 1 aluminum lake, FD&Cred no. 40 aluminum lake, glyceryl monostearate, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyacrylate dispersion, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, talc

Questions or comments? 1-800-719-9260

\*This product is not manufactured or distributed by Johnson 8 Johnson Consumer Inc., distributor of Pepcid Complete®. Copy Area





#### **GLUTEN FREE**

DISTRIBUTED BY: RITE AID 30 HUNTER LANE, CAMP HILL, PA 17011 WWW.RITEAID.COM

#### SATISFACTION GUARANTEE:

If you're not satisfied, we'll happily refund your money.

No Copy Area

#### **ACID REDUCER PLUS ANTACID**

famotidine, calcium carbonate and magnesium hydroxide tablet, chewable

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:11822-0500

Route of Administration ORAL

#### **Active Ingredient/Active Moiety** Basis of **Ingredient Name** Strength Strength FAMO TIDINE (UNII: 5QZO 15J2Z8) (FAMO TIDINE - UNII:5QZO 15J2Z8) **FAMOTIDINE** 10 mg MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, MAGNESIUM 165 mg HYDRO XIDE ION - UNII:9159UV381P) HYDRO XIDE CALCIUM CARBONATE (UNII: H0 G9 379 FGK) (CARBONATE ION - UNII: 7UJQ50 PE7D, CALCIUM CALCIUM 800 mg CATION - UNII:2M83C4R6ZB) CARBONATE

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS LACTOSE (UNII: 3S Y5LH9 PMK)		
ASPARTAME (UNII: Z0H242BBR1)		
D&C RED NO. 7 (UNII: ECW0LZ41X8)		
DEXTRATES (UNII: G263MI44RU)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERYL MONOSTEARATE (UNII: 230 OU9 XXE4)		
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)		
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
PO VIDO NE (UNII: FZ989 GH94E)		
TALC (UNII: 7SEV7J4R1U)		

Product Characteristics			
Color	PINK (mottled)	Score	no score
Shape	ROUND (bi-layered)	Size	17mm

Flavor	BERRY	Imprint Code	L321
Contains			

F	Packaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11822-0500-1	25 in 1 BOTTLE; Type 0: Not a Combination Product	11/08/2013	
2	NDC:11822-0500-2	50 in 1 BOTTLE; Type 0: Not a Combination Product	11/08/2013	
3	NDC:11822-0500-3	75 in 1 BOTTLE; Type 0: Not a Combination Product	11/08/2013	04/16/2015

<b>Marketing Info</b>	Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA077355	11/08/2013		

## Labeler - Rite Aid Corporation (014578892)

Revised: 6/2019 Rite Aid Corporation