

RITEAID NIGHTTIME SLEEP AID- diphenhydramine hydrochloride and acetaminophen liquid
RITE AID CORPORATION

RITEAID Nighttime Sleep-aid Pain Reliever

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Diphenhydramine HCl 50 mg

Purpose

Pain reliever

Nighttime sleep-aid

Uses

- for the temporary relief of occasional minor aches and pains with accompanying sleeplessness.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs., which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age

- with any other product containing diphenhydramine, even one used on skin.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic beverages
- drowsiness will occur
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- Sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only one dose (30 mL) per day (24 hours)
- Only use dose cup provided

adults & children 12 yrs & over	30 mL at bedtime
children under 12 yrs	do not use

Other information

- **each 30 mL contains:** sodium 10 mg
- store at room temperature
- very low sodium

Inactive ingredients

citric acid, flavors, FD&C Blue No. 1, FD&C Red No. 40, glycerin, potassium citrate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose.

Questions or comments?

1-866-467-2748

PRINCIPAL DISPLAY PANEL - 354 ml bottle

Compare to the active ingredient in Vicks® ZzzQuil™ Night Pain Sleep-Aid*

NDC# 11822-8791-1

Nighttime Sleep-Aid**Pain Reliever**

Diphenhydramine HCl

Acetaminophen

- Fall Asleep Fast
- Max Strength Pain Reliever
- Non-Habit Forming

Berry Flavor

Naturally and Artificially Flavored

No Added Alcohol

12 FL. OZ. (354 ml)

*This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® ZzzQuil™ Night Pain Sleep-Aid.

Distributed by:

3.5" Height

7.875" Width

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ take only one dose (30 mL) per day (24 hours)
■ only use the dose cup provided
adults & children 12 yrs & over 30 mL at bedtime
children under 12 yrs do not use
Other information
■ each 30 mL contains: sodium 10 mg
■ store at room temperature
■ very low sodium
Inactive ingredients citric acid, flavors, FD&C Blue No. 1, FD&C Red No. 40, glycerin, potassium citrate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose
Questions or comments? 1-866-467-2748
*This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® ZzzQuil™ Night Pain Sleep-Aid.
DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com
60012RITLR
LOT:
EXP:
0 11822 12989 3

Compare to the active ingredients in
Vicks® ZzzQuil™ Night Pain Sleep-Aid®
NDC 11822-8791-1
NIGHTTIME SLEEP-AID PAIN RELIEVER
DIPHENHYDRAMINE HCl
50 mg per 30 mL
ACETAMINOPHEN 1000 mg per 30 mL
max strength pain reliever
fall asleep fast
non-habit forming
no added alcohol
BERRY FLAVOR
NATURALLY AND ARTIFICIALLY FLAVORED
12 FL OZ (354 mL)

Drug Facts **TAMPER EVIDENT:** Do not use if printed shrink band is missing or broken
Active ingredients (in each 30 mL)
Acetaminophen 1000 mg.....Pain reliever
Diphenhydramine HCl 50 mg.....Nighttime sleep-aid
Uses ■ for the temporary relief of occasional minor aches and pains with accompanying sleeplessness.
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
■ taken with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product.
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ for children under 12 years of age
■ with any other product containing diphenhydramine, even one used on skin
Ask a doctor before use if you have ■ liver disease
■ a breathing problem such as emphysema or chronic bronchitis
■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland
Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers
When using this product ■ avoid alcoholic beverages ■ drowsiness will occur ■ do not drive a motor vehicle or operate machinery
Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days ■ new symptoms occur
■ Sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. These could be signs of a serious condition. ▶

RITEAID NIGHTTIME SLEEP AID

diphenhydramine hydrochloride and acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-8791
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	50 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-8791-1	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M010	06/01/2021	

Labeler - RITE AID CORPORATION (014578892)

Revised: 10/2024

RITE AID CORPORATION