

# **CHILDRENS PAIN AND FEVER- acetaminophen suspension**

## **Walgreen Company**

-----

### **Walgreen Co. Children's Pain & Fever Drug Facts**

#### **Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

#### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has**

liver disease

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child’s dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL

60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

### Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

### Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

### Questions or comments?

**1-800-719-9260**

### Package/Label Principal Display Panel

Walgreens

WALGREENS • PHARMACIST RECOMMENDED •

Compare to the active ingredient in Children's Tylenol® Oral Suspension

Dye-Free

Children's Pain & Fever

ACETAMINOPHEN 160 mg PER 5 mL

PAIN RELIEVER / FEVER REDUCER

ORAL SUSPENSION

Aspirin Free

Alcohol Free



Ibuprofen Free

Paraben Free

AGES 2-11 YEARS

4 FL OZ (118 mL)

Cherry flavor



**Drug Facts**

**Active ingredient (in each 5 mL)** **Purpose**  
 Acetaminophen 160 mg.....Pain reliever/fever reducer

**Uses** temporarily:  
 ■ reduces fever  
 ■ relieves minor aches and pains due to:  
 ■ the common cold ■ flu ■ headache  
 ■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.  
 ■ with other drugs containing acetaminophen  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has liver disease**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**  
 ■ pain gets worse or lasts more than 5 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present  
 These could be signs of a serious condition.

**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**  
 ■ this product does not contain directions or complete warnings for adult use  
 ■ do not give more than directed (see overdose warning)  
 ■ shake well before using  
 ■ mL = milliliter  
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.  
 ■ remove the child protective cap and squeeze your child's dose into the dosing cup  
 ■ repeat dose every 4 hours while symptoms last  
 ■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor  
**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**  
 ■ each 5 mL contains: sodium 3 mg  
 ■ store at 20-25°C (68-77°F)  
 ■ do not use if printed neckband is broken or missing

**Inactive ingredients** anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

**Questions or comments?** 1-800-719-9260

\*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
 †This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.

**DOSAGE CUP PROVIDED**  
 Gluten Free

**Walgreens**

**Dye-Free**

**Children's Pain & Fever**

NDC 0363-1230-28

**Walgreens**

Compare to the active ingredient in Children's Tylenol® Oral Suspension\*

**Dye-Free**

**Children's Pain & Fever**

ACETAMINOPHEN 160 mg PER 5 mL  
 PAIN RELIEVER / FEVER REDUCER  
 ORAL SUSPENSION

- Aspirin Free
- Alcohol Free
- Ibuprofen Free
- Paraben Free

**AGES 2-11 YEARS**

**4 FL OZ (118 mL)** **Cherry flavor**

**DIRECTIONS:**

**STEP 1. Shake well before using**




**STEP 2. Remove child protective cap**



**STEP 3. Squeeze your child's dose into the dosing cup**



**IMPORTANT:**  
 Please see dosing chart or ask a doctor for your child's correct dose.




W30RG0322-F  
 REV-1223

DISTRIBUTED BY: WALGREEN CO.  
 DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEED  
 walgreens.com ©2021 Walgreen Co.  
 ITEM 691994 W10106-0123-L

3 11917 04417 0

95926 94 C6



**CHILDRENS PAIN AND FEVER**  
 acetaminophen suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-1239
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

**Product Characteristics**

<b>Color</b>	WHITE (opaque,viscous)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1239-26	1 in 1 CARTON	05/17/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/17/2023	

Revised: 9/2024

Walgreen Company