MUCUS RELIEF DM MAXIMUM STRENGTH- dextromethorphan hbr, guaifenesin liquid Rite Aid Corporation

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg Guaifenesin 400 mg

Purpose

Cough suppressant

Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of cough
 - the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm(mucus)

When using this product,

do not use more than directed

Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent

headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age : do not use

Other information

- each 20 mL contains: sodium 20 mg
- store between 20-25°C (68-77°F). do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C red 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® DM Max* MAXIMUM STRENGTH

MUCUS RELIEF

DM MAX

MULTI-SYMPTOM RELIEF

COUGH SUPPRESSANT EXPECTORANT

DEXTROMETHORPHAN HBr 20 mg

GUAIFENESIN 400 mg

Controls cough

Relieves chest congestion

Thins & loosen mucus

For ages 12 & over

4-hour dosing

For ages 12 and over

FL OZ(mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max.

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

www.riteaid.com

Package Label



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DM Max.



PLD-A409A LB007075



DISTRIBUTED BY: RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011 www.riteaid.com

SATISFACTION GUARANTEE

f you're not satisfied, we'll ppily refund your money.

PEEL CORNER FOR DRUG FACTS

Drug Facts

Active ingredients Purposes (in each 20 mL)

Dextromethorphan HBr 20 mg.....Cough suppressant Guaifenesin 400 mg......Expectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phleam (mucus)

When using this product, do not use more than directed.

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not

Other information

- each 20 mL contains: sodium 20 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Drug Facts (continued)

Inactive ingredients

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate. sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS ▲

MUCUS RELIEF DM MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin liquid

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-7224		
Route of Administration	ORAL				

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL		
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL		

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
EDETATE SODIUM (UNII: MP1J8420LU)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
PROPYL GALLATE (UNII: 8D4SNN7V92)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
GLYCERIN (UNII: PDC6A3C0OX)		
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)		
WATER (UNII: 059QF0KO0R)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:11822- 7224-6	117 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/26/2021	02/01/2027	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	11/26/2021	02/01/2027	

Revised: 11/2025 Rite Aid Corporation