

# **STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

-----  
**Drug Facts**

**Active ingredient (in each 30 mL)**

Bismuth subsalicylate 525 mg

**Purpose**

Upset stomach reliever/Antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - gas
  - indigestion
  - nausea
  - fullness
  - belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

## **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## **Directions**

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
  - 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
  - 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
  - do not exceed 8 doses (240 mL) in 24 hours
  - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

## **Other information**

- **each 30 mL contains:** potassium 8 mg, sodium 9 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

## **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

## **Questions or comments?**

Call **1-888-309-9030**

## **Principal Display Panel**

Compare to the active ingredient of **Pepto-Bismol®\***

Regular Strength

### **Stomach Relief**

Bismuth subsalicylate 525 mg

Upset Stomach Reliever/Antidiarrheal

Relieves:

- heartburn
- indigestion
- nausea
- upset stomach
- diarrhea

Alcohol Free

Sugar Free

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

## **Package Label**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**DANGER: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING. REMOVE DOSE CAP ON UNICEF CAP TO OPEN OR RECLOSE.**



**100%**  
Satisfaction  
Guaranteed!  
(888) 309-9030

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE TN 37072

**DG** health

**VALUE PACK**

Compare to the  
active ingredient  
of Pepto-Bismol®

## Regular Strength Stomach Relief

Bismuth Subsalicylate 525 mg  
Upset Stomach Reliever/  
Antidiarrheal

Relieves:  
• Heartburn  
• Indigestion  
• Nausea  
• Upset stomach  
• Diarrhea

Alcohol free  
Sugar free

## Regular Strength Stomach Relief

Bismuth Subsalicylate 525 mg  
Upset Stomach Reliever/  
Antidiarrheal

Relieves:  
• Heartburn  
• Indigestion  
• Nausea  
• Upset stomach  
• Diarrhea

Alcohol free  
Sugar free



TOTAL 32 fl oz (946 mL) - 2 x 16 fl oz (473 mL)

| Drug Facts   |  |
|--|--|
| <b>Active ingredient (in each 30 mL dose)</b><br>Bismuth subsalicylate 525 mg  | <b>Purpose</b><br>Upset stomach reliever/Antidiarrheal |
| <b>Uses relieves</b> ■ diarrhea ■ traveler's diarrhea<br>■ upset stomach due to overindulgence in food and drink, including<br>■ heartburn ■ gas ■ indigestion ■ nausea ■ fullness ■ belching  |  |
| <b>Warnings</b><br><b>Key's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.<br><b>Do not use if you have</b> ■ an ulcer ■ bleeding problem ■ bloody or black stool<br><b>Ask a doctor before use if you have</b> ■ fever ■ mucus in the stool<br><b>Ask a doctor or pharmacist before use if you are taking any drug for</b> ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)<br>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.<br><b>Stop use and ask a doctor if</b><br>■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days<br><b>If pregnant or breast-feeding,</b> ask a health professional before use.<br><b>Keep out of reach of children.</b><br>In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. |  |
| <b>Directions</b><br>■ mL = milliliter ■ shake well before using<br>■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product<br>■ adults and children 12 years and over<br>■ 30 mL (1 dose) every 30 min or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea<br>■ 30 mL (1 dose) every 15 min as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)<br>■ do not exceed 6 doses (240 mL) in 24 hours<br>■ use until diarrhea stops but not more than 2 days<br>■ children under 12 years of age: ask a doctor<br>■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea   |  |
| <b>Other information</b><br>■ each 30 mL contains: potassium 8 mg, sodium 9 mg<br>■ each 30 mL contains: salicylate 200 mg<br>■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)   |  |
| <b>Inactive ingredients:</b> benzoic acid, D&C red #22, D&C red #28, D&C red #29, hydroxy, glycerin, purified water, sucrose, xanthan gum  |  |
| <b>Questions or comments?</b> Call 1-888-309-9030  |  |

### DOLLAR GENERAL HEALTH Regular Strength Stomach Relief

## STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:55910-624 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength     | Strength        |
|--|-----------------------|-----------------|
| <b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ) | BISMUTH SUBSALICYLATE | 525 mg in 30 mL |

### Inactive Ingredients

| Ingredient Name                              | Strength |
|--|----------|
| <b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)       |          |
| <b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT) |          |
| <b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH) |          |
| <b>WATER</b> (UNII: 059QF0KO0R)              |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)           |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)          |          |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)        |          |

### Packaging

| # | Item Code        | Package Description  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:55910-624-16 | 473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 10/30/2019           |                    |
| 2 | NDC:55910-624-08 | 237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 10/30/2019           |                    |
| 3 | NDC:55910-       | 2 in 1 PACKAGE   | 10/30/2019           |                    |

|   |                  |  |            |  |
|---|------------------|--|------------|--|
| 2 | 624-32           | 2 III 1 PACKAGE  | 10/30/2019 |  |
| 3 |                  | 473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product |            |  |
| 4 | NDC:55910-624-04 | 118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 10/30/2019 |  |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M008                                     | 10/30/2019           |                    |

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 12/2023

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)