

**ALGIDOL FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, doxylamine succinate**  
**Gelpharma S.A. de C.V.**

-----  
**Algidol Flu**

***Drug Facts***

***Active ingredients (in each softgel capsule)***

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

***Uses***

- temporarily relieves
- minor aches and pains associated with the common cold, sore throat, or headache
- cough associated with the common cold
- nasal congestion due to the common cold
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

***Warnings***

This product contains acetaminophen. Severe liver damage may occur if you take: **Liver warning:**

Acetaminophen may cause severe skin reactions. Symptoms may include: **Allergy alert:**

If a skin reaction occurs, stop use and seek medical help right away.

- more than 8 capsules in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- skin reddening

- blisters
- rash

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.
- if you have heart disease, high blood pressure, thyroid disease, diabetes, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor.

**Ask a doctor before use if**

you have liver disease.

**Ask a doctor or pharmacist before use**

if you are taking the blood thinning drug warfarin.

**When using this product**

- do not exceed the recommended dosage
- do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition.
- if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- if symptoms do not improve within 7 days or are accompanied by fever, consult a doctor.

**Stop use and ask a doctor if**

- pain or nasal decongestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- if nervousness, dizziness, or sleeplessness occur

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- do not exceed 8 capsules per 24 hours

---

Adults & children 12 years & over	2 capsules with water every 4 hours
Children 4 to under 12 years	Consult a doctor
Children under 4 years	Do not use

---

### **Other information**

- do not exceed 77°F (25°C)
- avoid excessive heat

### **Inactive ingredients**

gelatin, purified water, glycerin, polyethylene glycol 1000, propylene glycol, polyethylene glycol 600, povidone K30, orange flavor, FD&C Yellow No. 6, BHT, BHA.

### **Drug Facts**

#### **Active ingredients (in each softgel capsule)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

#### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

#### **Uses**

- temporarily relieves
- minor aches and pains associated with the common cold, sore throat, or headache
- cough associated with the common cold

- nasal congestion due to the common cold
- runny nose and alleviates sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever
- temporarily reduces fever

## **Warnings**

This product contains acetaminophen. Severe liver damage may occur if you take: **Liver warning:**

Acetaminophen may cause severe skin reactions. Symptoms may include: **Allergy alert:**

If a skin reaction occurs, stop use and seek medical help right away.

- more than 8 capsules in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
  
- skin reddening
- blisters
- rash

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.
- if you have heart disease, high blood pressure, thyroid disease, diabetes, a breathing problem such as emphysema or chronic bronchitis, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor.

## **Ask a doctor before use if**

you have liver disease.

## **Ask a doctor or pharmacist before use if**

you are taking the blood thinning drug warfarin.

## **When using this product**

- do not exceed the recommended dosage
- do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. because these could be signs of a serious condition.
- if sore throat is severe, persists for more than 2 days, is accompanied or followed by

fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- if symptoms do not improve within 7 days or are accompanied by fever, consult a doctor.
- may cause excitability especially in children
- may cause marked drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages while taking this product. Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if**

- pain or nasal decongestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- if nervousness, dizziness, or sleeplessness occur

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 capsules per 24 hours

---

Adults & children 12 years & over	2 capsules with water every 4 hours
Children 4 to under 12 years	Consult a doctor
Children under 4 years	Do not use

---

**Other information**

- do not exceed 77°F (25°C)
- avoid excessive heat

**Inactive ingredients**

gelatin, purified water, polyethylene glycol 1000, glycerin, polyethylene glycol 600, propylene glycol, povidone K30, citric acid, mint flavor, FD&C Blue No. 1, FD&C Yellow No. 6, BHT, BHA.

# Package Labeling:

MI-1249

Do not take Algidol Flu Daytime and Algidol Flu Nighttime at the same time.  
 Always take at least 4 hours before taking another dose of Algidol Flu.  
 Do not take more than a total of 8 softgels in any 24-hour period.  
 Take only as directed.  
 Keep carton for full information.

**Tamper evident: Use only if blisters are intact and carton is not damaged.**

DAYTIME	NIGHTTIME
<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each softgel) Purpose</b>                      Acetaminophen 325 mg ..... Pain reliever/fever reducer                      Dextromethorphan HBr 10 mg ..... Cough suppressant                      Guaifenesin 200 mg ..... Expectorant                      Phenylephrine HCl 5 mg ..... Nasal decongestant</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves                             <ul style="list-style-type: none"> <li>minor aches and pains associated with the common cold, sore throat, or headache</li> <li>cough associated with the common cold</li> <li>nasal congestion due to the common cold</li> </ul> </li> <li>temporarily reduces fever</li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes</li> </ul> <p><b>Warnings</b></p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> <li>more than 8 softgels in 24 hours, which is the maximum daily amount</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p>Directions</p> <ul style="list-style-type: none"> <li>for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor</li> <li>if you have heart disease, high blood pressure, thyroid disease, diabetes, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor</li> </ul> <p>Ask a doctor before use if you have liver disease.</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product</p> <ul style="list-style-type: none"> <li>do not exceed the recommended dosage</li> <li>do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition.</li> </ul>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each softgel) Purpose</b>                      Acetaminophen 325 mg ..... Pain reliever/fever reducer                      Dextromethorphan HBr 10 mg ..... Cough suppressant                      Doxylamine succinate 6.25 mg ..... Antihistamine                      Phenylephrine HCl 5 mg ..... Nasal decongestant</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves                             <ul style="list-style-type: none"> <li>minor aches and pains associated with the common cold, sore throat, or headache</li> <li>cough associated with the common cold</li> <li>nasal congestion due to the common cold</li> <li>runny nose and alleviates sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever</li> </ul> </li> <li>temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> <li>more than 8 softgels in 24 hours, which is the maximum daily amount</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p>Directions</p> <ul style="list-style-type: none"> <li>for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where cough is accompanied by excessive phlegm (mucus) unless directed by a doctor</li> <li>if you have heart disease, high blood pressure, thyroid disease, diabetes, a breathing problem such as emphysema or chronic bronchitis, or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor</li> </ul> <p>Ask a doctor before use if you have liver disease.</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product</p> <ul style="list-style-type: none"> <li>do not exceed the recommended dosage</li> <li>do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition.</li> </ul>

MI-1249

**Algidol**  
 Max Strength **Flu**  
 NIGHTTIME  
 Severe Cold & Flu

**Drug Facts (Continued)**

symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition.

- if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- if symptoms do not improve within 7 days or are accompanied by fever, consult a doctor.
- may cause marked drowsiness, alcohol, sedatives, and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages while taking this product. Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor. Use caution when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if

- pain or nasal decongestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- if nervousness, dizziness, or sleepiness occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 softgels per 24 hours

Adults & children 12 years & over	2 softgels with water every 4 hours
Children 4 to under 12 years	Consult a doctor
Children under 4 years	Do not use

**Other information**

- do not exceed 77°F (25°C)
- avoid excessive heat

**Inactive ingredients**  
 gelatin, purified water, polyethylene glycol 1000, glycerin, polyethylene glycol 600, propylene glycol, povidone K30, citric acid, mint flavor, FD&C Blue No. 1, FD&C Yellow No. 6, BHT, BHA.

GEL pharma NDC: 69377-003-01

**Algidol**  
 Max Strength **Flu**  
 DAYTIME  
 Severe Cold & Flu

**Relief from: Fever, Minor Aches & Pains, Headache, Cough, Nasal Congestion**

**DAYTIME NIGHTTIME**  
 Severe Cold & Flu

**ACETAMINOPHEN (PAIN RELIEVER- FEVER REDUCER)**  
**DEXTROMETHORPHAN HBr (COUGH SUPPRESSANT)**  
**GUAIFENESIN (EXPECTORANT)**  
**PHENYLEPHRINE HCl (NASAL DECONGESTANT)**

Non-Drowsy Relief from:

- Sore Throat
- Sinus Pressure
- Chest Congestion
- Sneezing
- Runny Nose
- Itchy Throat

Fast Dissolving Liquid Filling

16 DAYTIME 8 NIGHTTIME

**24 Softgels**

**Algidol**  
 Max Strength **Flu**  
 DAYTIME  
 Severe Cold & Flu

**Drug Facts (Continued)**

- if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- if symptoms do not improve within 7 days or are accompanied by fever, consult a doctor.

Stop use and ask a doctor if

- pain or nasal decongestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- if nervousness, dizziness, or sleepiness occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 softgels per 24 hours

Adults & children 12 years & over	2 softgels with water every 4 hours
Children 4 to under 12 years	Consult a doctor
Children under 4 years	Do not use

**Other information**

- do not exceed 77°F (25°C)
- avoid excessive heat

**Inactive ingredients**  
 gelatin, purified water, glycerin, polyethylene glycol 1000, propylene glycol, polyethylene glycol 600, povidone K30, orange flavor, FD&C Yellow No. 6, BHT, BHA.

Manufactured by: Gelpharma S.A. de C.V.  
 Av. Paseo Del Pacifico No. 380  
 GDL Technology Park, Zapopan, JAL. 45010, Mexico  
 MADE IN MEXICO  
 Questions? (346) 326-1728

GEL pharma

8 50025 47601 0





## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	0 CAPSULE	2
Part 2	0 CAPSULE	1

## Part 1 of 2

### ALGIDOL DAYTIME

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule

## Product Information

Item Code (Source)	NDC:69377-004
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 1000 (UNII: U076Q6Q621)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9)	
POVIDONE K30 (UNII: U725QWY32X)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	

## Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	10mm
Flavor		Imprint Code	



**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69377-004-01	2 in 1 BLISTER PACK		
1		8 in 1 CAPSULE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/01/2022	

**Part 2 of 2****ALGIDOL NIGHTTIME**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule

**Product Information**

<b>Item Code (Source)</b>	NDC:69377-005
<b>Route of Administration</b>	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POLYETHYLENE GLYCOL 1000</b> (UNII: U076Q6Q621)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 600</b> (UNII: NL4J9F21N9)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69377-005-01	1 in 1 BLISTER PACK		
1		8 in 1 CAPSULE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/01/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/01/2022	

**Labeler** - Gelpharma S.A. de C.V. (812773665)

Revised: 11/2023

Gelpharma S.A. de C.V.