

**DOLOGESIC DF- acetaminophen, dextbrompheniramine maleate tablet**  
**Llorens Pharmaceutical International Division**

-----  
**Llorens-DologesicDFTab 010**

**Active Ingredients**

(in each caplet)

Acetaminophen 500 mg

Dextbrompheniramine Maleate 1 mg

**Purpose**

Pain reliever/fever reducer

antihistamine

**Uses**

- For the temporary relief of minor aches and pains associated with
- headache
- muscular aches
- backaches
- minor arthritis pain
- common cold
- toothaches
- menstrual cramps
- temporarily reduces fever
- itchy and watery eyes due to hay fever

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 capsules in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blister
- rash.

If a skin reaction occurs, stop and seek medical help right away.

**While using this product**

- drowsiness may occur; alcohol, sedatives, and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages
- do not take this product if you are taking sedatives or tranquilizers without consulting your doctor
- use caution when driving a motor vehicle or operating machinery

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in the product
- for more than 10 days for pain, unless directed by a doctor
- for more than 3 days for fever, unless directed by a doctor

**Ask a doctor before use:**

- if you have liver disease
- if you are taking the blood thinning drug warafin

**Stop using this product and ask a doctor if:**

- pain gets worse or last more than 10 days
- fever gets worse or last more than 3 days
- new symptoms occur
- redness or swelling is present
- these could be signs of a serious condition

**Keep out of reach of children**

**If you are pregnant or breast-feeding,** ask a health professional before use.

**Directions**

**Overdose Warning:** Taking more than recommended dose (overdose), may cause liver damage. In case of accidental overdose, contact a physician or Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

**Do not exceed recommended dosage**

Age	Dose
adults and children 12 years of age and over	Take 2 caplets every 4-6 hours. Do not take more than 8 caplets in 24 hours, or as directed by a doctor
	Do not use in children under



Ingredient Name	Strength
<b>HYDROXYPROPYL METHYLCELLULOSE</b> (UNII: 3NXW29V3WO)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>TRICALCIUM PHOSPHATE</b> (UNII: K4C08XP666)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	LLORENS
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-010-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2016	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/01/2016	

**Labeler** - Llorens Pharmaceutical International Division (037342305)

Revised: 12/2025

Llorens Pharmaceutical International Division