

PHARBETOL- acetaminophen 325 mg tablet
Ulai Health LLC

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 tablets every 4 to 6 hours while symptoms last• do not take more than 10 tablets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children 6 to under 12 years	<ul style="list-style-type: none">• take 1 tablet every 4 to 6 hours while symptoms last• do not take more than 5 tablets in 24 hours• do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Other information

- store at 20-25 °C (68-77 °F)

Inactive ingredients

povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

Questions or comments?

(866) 562-2756 (Mon - Fri 8 AM to 4 PM EST)

PHARBEST

NDC 73057-353-11

Manufactured in the USA

Regular Strength *COMPARE TO the active ingredient in TYLENOL® Regular Strength

Contains no Aspirin

PHARBETOL

Acetaminophen 325 mg each

Pain Reliever/Fever Reducer

1000 TABLETS

PHARBEST
NDC 73057-353-11

Manufactured
in the **USA**

Regular Strength *COMPARE TO the active ingredient
in TYLENOL® Regular Strength

Contains no Aspirin

PHARBETOL

Acetaminophen 325 mg each

Pain Reliever/Fever Reducer

1000 TABLETS

THIS PACKAGE FOR HOUSEHOLD WITHOUT YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	<p>Active ingredient (in each tablet) Acetaminophen 325 mg.....Pain reliever/fever reducer</p> <p>Uses Temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ toothache ■ muscular aches ■ menstrual and menstrual cramps ■ temporarily reduces fever</p> <p>Warnings Liver warnings: This product contains acetaminophen. The maximum daily dose of this product is 4,000 mg (10 tablets) in 24 hours for adults and 3,250 mg (10 tablets) in 24 hours for children. Severe liver damage may occur if: ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ Child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks in 24 hours ■ you have liver disease ■ you are taking other drugs ■ Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription) ■ if you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.</p> <p>Ask a doctor before use if you have liver disease.</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>Stop use and ask a doctor if: ■ pain gets worse or lasts more than 10 days in adults ■ pain gets worse or lasts more than 5 days in children under 12 years ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ these could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>
Directions	<p>do not take more than directed (see overdose warning)</p> <p>adults and children 12 years and over ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a doctor</p> <p>children 6 to under 12 years ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours</p> <p>children under 6 years ■ do not use for more than 5 days unless directed by a doctor ■ ask a doctor</p>
Other information	<p>■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid</p>
Questions or comments?	<p>(866) 562-2756 (Mon- Fri 8 AM to 4 PM EST)</p>

Org. 06/19
3 1730571353116
Distributed by:
ULAI HEALTH LLC,
Alexandria, VA 22312

PHARBETOL

acetaminophen 325 mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73057-353
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
Inactive Ingredients				
Ingredient Name		Strength		
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	PH020	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73057-353-11	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/08/2019	
2	NDC:73057-353-08	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/08/2019	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	07/08/2019		

Labeler - Ulai Health LLC (081181535)

Revised: 1/2025

Ulai Health LLC