


PYRAZINAMIDE- pyrazinamide tablet
Hikma Pharmaceuticals USA Inc

Pyrazinamide tablets, usp

DESCRIPTION

Pyrazinamide, the pyrazine analogue of nicotinamide, is an antituberculous agent. It is a white crystalline powder, stable at room temperature, and sparingly soluble in water.

Pyrazinamide has the following structural formula:



C₅H₅N₃O M.W. 123.11

Each pyrazinamide tablet for oral administration contains 500 mg of pyrazinamide and the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate (dihydrate), microcrystalline cellulose, and stearic acid.

CLINICAL PHARMACOLOGY

Pyrazinamide is well absorbed from the GI tract and attains peak plasma concentrations within 2 hours. Plasma concentrations generally range from 30 to 50 mcg/mL with doses of 20 to 25 mg/kg. It is widely distributed in body tissues and fluids including the liver, lungs and cerebrospinal fluid (CSF). The CSF concentration is approximately equal to concurrent steady-state plasma concentrations in patients with inflamed meninges.¹ Pyrazinamide is approximately 10% bound to plasma proteins.² The half-life (t_{1/2}) of pyrazinamide is 9 to 10 hours in patients with normal renal and hepatic function. The plasma half-life may be prolonged in patients with impaired renal or hepatic function. Pyrazinamide is hydrolyzed in the liver to its major active metabolite, pyrazinoic acid. Pyrazinoic acid is hydroxylated to the main excretory product, 5-hydroxypyrazinoic acid.³

Approximately 70% of an oral dose is excreted in the urine, mainly by glomerular filtration within 24 hours.³

Pyrazinamide may be bacteriostatic or bactericidal against *Mycobacterium tuberculosis* depending on the concentration of the drug attained at the site of infection. The mechanism of action is unknown. *In vitro* and *in vivo* the drug is active only at a slightly acidic pH.

INDICATIONS AND USAGE

Pyrazinamide is indicated for the initial treatment of active tuberculosis in adults and children when combined with other antituberculous agents. (The current recommendation of the CDC for drug-susceptible disease is to use a six-month regimen for initial treatment of active tuberculosis, consisting of isoniazid, rifampin and pyrazinamide given for 2 months, followed by isoniazid and rifampin for 4 months.*⁴)

(Patients with drug-resistant disease should be treated with regimens individualized to their situation. Pyrazinamide frequently will be an important component of such therapy.)

(In patients with concomitant HIV infection, the physician should be aware of current recommendation of CDC. It is possible these patients may require a longer course of treatment).

It is also indicated after treatment failure with other primary drugs in any form of active tuberculosis.

Pyrazinamide should only be used in conjunction with other effective antituberculous agents.

*See recommendations of Center for Disease Control (CDC) and American Thoracic Society for complete regimen and dosage recommendations.⁴

CONTRAINDICATIONS

Pyrazinamide is contraindicated in persons:

- with severe hepatic damage.
- who have shown hypersensitivity to it.
- with acute gout.

WARNINGS

Patients started on pyrazinamide should have baseline serum uric acid and liver function determinations. Those patients with preexisting liver disease or those at increased risk for drug related hepatitis (e.g., alcohol abusers) should be followed closely.

Pyrazinamide should be discontinued and not be resumed if signs of hepatocellular damage or hyperuricemia accompanied by an acute gouty arthritis appear.

PRECAUTIONS

General

Pyrazinamide inhibits renal excretion of urates, frequently resulting in hyperuricemia which is usually asymptomatic. If hyperuricemia is accompanied by acute gouty arthritis, pyrazinamide should be discontinued.

Pyrazinamide should be used with caution in patients with a history of diabetes mellitus, as management may be more difficult.

Primary resistance of *M. tuberculosis* to pyrazinamide is uncommon. In cases with known or suspected drug resistance, *in vitro* susceptibility tests with recent cultures of

M. tuberculosis against pyrazinamide and the usual primary drugs should be performed. There are few reliable *in vitro* tests for pyrazinamide resistance. A reference laboratory capable of performing these studies must be employed.

Information for Patients

Patients should be instructed to notify their physicians promptly if they experience any of the following: fever, loss of appetite, malaise, nausea and vomiting, darkened urine, yellowish discoloration of the skin and eyes, pain or swelling of the joints.

Compliance with the full course of therapy must be emphasized, and the importance of not missing any doses must be stressed.

Laboratory Tests

Baseline liver function studies [especially ALT (SGPT), AST (SGOT) determinations] and uric acid levels should be determined prior to therapy. Appropriate laboratory testing should be performed at periodic intervals and if any clinical signs of symptoms occur during therapy.

Interactions

Drug/Laboratory Test Interactions

Pyrazinamide has been reported to interfere with ACETEST[®] and KETOSTIX[®] urine tests to produce a pink-brown color.⁵

Carcinogenesis, Mutagenesis, Impairment of Fertility^{6,7,8}

In lifetime bioassays in rats and mice, pyrazinamide was administered in the diet at concentrations of up to 10,000 ppm. This resulted in estimated daily doses for the mouse of 2 g/kg, or 40 times the maximum human dose, and for the rat of 0.5 g/kg, or 10 times the maximum human dose. Pyrazinamide was not carcinogenic in rats or male mice and no conclusion was possible for female mice due to insufficient numbers of surviving control mice.

Pyrazinamide was not mutagenic in the Ames bacterial test, but induced chromosomal aberrations in human lymphocyte cell cultures.

Pregnancy

Teratogenic Effects

Animal reproduction studies have not been conducted with pyrazinamide. It is also not known whether pyrazinamide can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Pyrazinamide should be given to a pregnant woman only if clearly needed.

Nursing Mothers

Pyrazinamide has been found in small amounts in breast milk. Therefore, it is advised that pyrazinamide be used with caution in nursing mothers taking into account the risk-benefit of this therapy.⁹

Pediatric Use

Pyrazinamide regimens employed in adults are probably equally effective in pediatric patients.^{4,10,11} Pyrazinamide appears to be well tolerated in pediatric patients.

Geriatric Use¹²

Clinical studies of pyrazinamide did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic or renal function, and of concomitant disease or other drug therapy.

It does not appear that patients with impaired renal function require a reduction in dose. It may be prudent to select doses at the low end of the dosing range, however.¹³

ADVERSE REACTIONS

General

Fever, porphyria and dysuria have rarely been reported. Gout (see **PRECAUTIONS**).

Gastrointestinal

The principal adverse effect is a hepatic reaction (see **WARNINGS**). Hepatotoxicity appears to be dose related, and may appear at any time during therapy. GI disturbances including nausea, vomiting and anorexia have also been reported.

Hematologic and Lymphatic

Thrombocytopenia and sideroblastic anemia with erythroid hyperplasia, vacuolation of erythrocytes and increased serum iron concentration have occurred rarely with this drug. Adverse effects on blood clotting mechanisms have also been rarely reported.

Other

Mild arthralgia and myalgia have been reported frequently. Hypersensitivity reactions including rashes, urticaria, and pruritis have been reported. Fever, acne, photosensitivity, porphyria, dysuria and interstitial nephritis have been reported rarely.

OVERDOSAGE

Overdosage experience is limited. In one case report of overdose, abnormal liver function tests developed. These spontaneously reverted to normal when the drug was stopped. Clinical monitoring and supportive therapy should be employed. Pyrazinamide is dialyzable.¹³

DOSAGE AND ADMINISTRATION

Pyrazinamide should always be administered with other effective antituberculous drugs.

It is administered for the initial 2 months of a 6-month or longer treatment regimen for drug-susceptible patients. Patients who are known or suspected to have drug-resistant disease should be treated with regimens individualized to their situation.

Pyrazinamide frequently will be an important component of such therapy.

Patients with concomitant HIV infection may require longer courses of therapy. Physicians treating such patients should be alert to any revised recommendations from CDC for this group of patients.

Usual dose: Pyrazinamide is administered orally, 15 to 30 mg/kg once daily. Older regimens employed 3 or 4 divided doses daily, but most current recommendations are for once a day. Three grams per day should not be exceeded. The CDC recommendations do not exceed 2 g per day when given as a daily regimen (see table).

Alternatively, a twice weekly dosing regimen (50 to 75 mg/kg twice weekly based on lean body weight) has been developed to promote patient compliance with a regimen on an outpatient basis. In studies evaluating the twice weekly regimen, doses of pyrazinamide in excess of 3 g twice weekly have been administered. This exceeds the recommended maximum 3 g/daily dose. However, an increased incidence of adverse reactions has not been reported.

This table is taken from the CDC-American Thoracic Society joint recommendations.⁴

Recommended Drugs for the Initial Treatment of Tuberculosis in Children and Adults

Drug	Daily Dose*	
	Children	Adults
Isoniazid	10 to 20 mg/kg PO or IM	5 mg/kg PO or IM
Rifampin	10 to 20 mg/kg PO	10 mg/kg PO
Pyrazinamide	15 to 30 mg/kg PO	15 to 30 mg/kg PO
Streptomycin	20 to 40 mg/kg IM	15 mg/kg IM
Ethambutol	15 to 25 mg/kg PO	15 to 25 mg/kg PO
Drug	Maximal Daily Dose in Children and Adults	
Isoniazid	300 mg	
Rifampin	600 mg	
Pyrazinamide	2 g	
Streptomycin	1 g**	
Ethambutol	2.5 g	
Drug	Twice Weekly Dose	
	Children	Adults
Isoniazid	20 to 40 mg/kg Max. 900 mg	15 mg/kg Max. 900 mg
Rifampin	10 to 20 mg/kg	10 mg/kg

Pyrazinamide	Max. 600 mg 50 to 70 mg/kg	Max. 600 mg 50 to 70 mg/kg
Streptomycin	25 to 30 mg/kg IM	25 to 30 mg/kg IM
Ethambutol	50 mg/kg	50 mg/kg

Definition of abbreviations: PO = perorally; IM = intramuscularly.

HOW SUPPLIED

Pyrazinamide Tablets, USP contain pyrazinamide 500 mg. They are supplied as white, round, scored tablets debossed "VP/012" in containers of 60 tablets, NDC# 0054-0812-21, in containers of 90 tablets, NDC# 0054-0812-22, in containers of 100 tablets, NDC# 0054-0812-25, and in containers of 500 tablets, NDC# 0054-0812-29.

Storage: Store in a well-closed container at controlled room temperature 15°C to 30°C (59°F to 86°F).

Dispense in a well-closed container with a child resistant closure.

Rx only

Distributed by:

Hikma Pharmaceuticals USA Inc.
Berkeley Heights, NJ 07922

Manufactured by:

Mikart, LLC
Atlanta, GA 30318

1191B00 H/01

Revised: October 2023

REFERENCES

1. Drug Information, American Hospital Formulary Service. American Society of Hospital Pharmacists. Bethesda, Md. 1991.
2. USPDI, Drug Information for the Health Care Professional. United States Pharmacopeial Convention, Inc. Rockville, Md. 1991:1B:2226-2227.
3. Goodman-Gilman A, Rall TW, Nies AS, Taylor P. The Pharmacological Basis of Therapeutics, ed 8. New York, Pergamon Press. 1990; 1154.
4. Treatment of tuberculosis and tuberculosis infection in adults and children. Am Rev Respir Dis. 1986; 134:363-368.
5. Reynolds JEF, Parfitt K, Parsons AV, Sweetman SC. Martindale The Extra Pharmacopoeia, ed 29. London, The Pharmaceutical Press. 1989; 569-570.
6. Bioassay of pyrazinamide for possible carcinogenicity. National Cancer Institute Carcinogenesis Technical Report Series No. 48, 1978.
7. Zenger E, Anderson B, Haworth S, Lawlor T, Mortelmans K, Speck W. Salmonella mutagenicity tests: III. Results from the testing of 255 chemicals. Environ Mutagen. 1987; 9 (Suppl 9):1109.
8. Roman IC, Georgian L. Cytogenetic effects of some antituberculosis drugs in vitro.

Mutation Research. 1977; 48:215-224.

9. Holdiness M. Antituberculosis drugs and breast-feeding. Arch Intern Med. 1984; 144:1888.
10. Turcios N, Evans H. Preventing and managing tuberculosis in children. J Resp Dis. 1989; 10 (6) (Jun):23.
11. Starke JR. Multidrug therapy for tuberculosis in children. Pediatr Infec Dis J. 1990; 9:785-793.
12. Specific requirements on content and format of labeling for human prescription drugs; proposed addition of "geriatric use" subsection in the labeling. Federal Register. 1990; 55 (212) (Nov 1):46134-46137.
13. Stamathakis G, Montes C, Trouvin JH, et al. Pyrazinamide and pyrazinoic acid pharmacokinetics in patients with chronic renal failure. Clinical Nephrology. 1988; 30:230-234.

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 0054-0812-21 60 Tablets

**Pyrazinamide
Tablets, USP
500 mg
Rx only**



PYRAZINAMIDE

pyrazinamide tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0054-0812
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
PYRAZINAMIDE (UNII: 2KNI5N06TI) (PYRAZINAMIDE - UNII:2KNI5N06TI)		PYRAZINAMIDE	500 mg	
Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)				
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	WHITE	Score	2 pieces	
Shape	ROUND	Size	13mm	
Flavor		Imprint Code	VP;012	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0054-0812-21	60 in 1 CONTAINER; Type 0: Not a Combination Product	08/30/2024	
2	NDC:0054-0812-22	90 in 1 CONTAINER; Type 0: Not a Combination Product	08/30/2024	
3	NDC:0054-0812-25	100 in 1 CONTAINER; Type 0: Not a Combination Product	08/30/2024	
4	NDC:0054-0812-29	500 in 1 CONTAINER; Type 0: Not a Combination Product	08/30/2024	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA081319	08/30/2024		

Labeler - Hikma Pharmaceuticals USA Inc (080189610)

Registrant - Hikma Pharmaceuticals USA Inc (080189610)

Establishment

Name	Address	ID/FEI	Business Operations
Mikart, LLC		030034847	MANUFACTURE(0054-0812) , PACK(0054-0812)