

NYSTATIN- nystatin suspension
Cardinal Health 107, LLC

Nystatin Oral Suspension USP


[100,000 units per mL]

I0868C0622 R06/22

Rx Only

DESCRIPTION

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*.
Structural formula:



Nystatin Oral Suspension USP, for oral administration, contains 100,000 USP Nystatin Units per mL. Inactive ingredients: alcohol ($\leq 1\%$ v/v), artificial peppermint flavor, cherry flavor, citric acid, D&C Yellow No. 10, FD&C Red No. 40, glycerin, magnesium aluminum silicate, methylparaben, potassium phosphate dibasic, propylene glycol, propylparaben, purified water and sucrose.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

Microbiology

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida* species

become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

CONTRAINDICATIONS

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

PRECAUTIONS

General

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

Pregnancy

Teratogenic Effects Category C

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

Pediatric Use

See **DOSAGE AND ADMINISTRATION**.

ADVERSE REACTIONS

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General**).

Gastrointestinal: Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

Dermatologic: Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

Other: Tachycardia, bronchospasm, facial swelling, and non-specific myalgia have also been rarely reported.

OVERDOSAGE

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (see **CLINICAL PHARMACOLOGY, Pharmacokinetics**).

DOSAGE AND ADMINISTRATION

INFANTS

2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

NOTE: Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

CHILDREN AND ADULTS

4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

HOW SUPPLIED

Nystatin Oral Suspension USP, 100,000 USP Nystatin Units per mL, is available in a cherry, peppermint flavored, light creamy yellow, ready-to-use suspension, supplied in the following oral dosage forms:

Overbagged with 5 x 5 mL unit dose cup in each bag, NDC 55154-4314-5

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

Storage

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Avoid freezing.

Distributed by:

MAJOR[®] PHARMACEUTICALS

Indianapolis, IN 46268

Distributed By:

Cardinal Health

Dublin, OH 43017

L58543101024

R06/22

Package/Label Display Panel

NDC 55154-4314-5

NYSTATIN ORAL SUSPENSION, USP

500,000 UNITS / 5 ML

5 UNIT DOSE CUPS



H116

NDC 55154-4314-5

NYSTATIN ORAL SUSPENSION, USP

500,000 UNITS / 5 ML

5 UNIT DOSE CUPS

Delivers 5 mL

Alcohol \leq 1% v/v

Shake Well

Usual dosage: See product insert for prescribing information, precautions and warnings.

STORAGE: Store at 20° to 25° C (68° to 77° F)

[See USP Controlled Room Temperature]. **AVOID FREEZING**

Protect from light

RX ONLY

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only.

Keep this and all drugs out of the reach of children.

MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

www.majorpharmaceuticals.com

Distributed by Cardinal Health

Dublin, OH 43017

L58543101024

NYSTATIN

nystatin suspension

Product Information

HUMAN PRESCRIPTION

Item Code

NDC 55154-4314/NDC 0004

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55154-4314(NDC:0904-7276)
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USP'U] in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
DIBASIC POTASSIUM PHOSPHATE (UNII: CI71S98N1Z)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics			
Color	yellow (light-creamy)	Score	
Shape		Size	
Flavor	CHERRY (with peppermint)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-4314-5	5 in 1 BAG	08/01/2023	
1		5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203621	07/17/2020	

Labeler - Cardinal Health 107, LLC (118546603)

Revised: 10/2025

Cardinal Health 107, LLC