

**LOLLICAINE BUBBLE GUM- benzocaine gel, dentifrice**  
**LOLLICAINE CHERRY- benzocaine gel, dentifrice**  
**LOLLICAINE MINT- benzocaine gel, dentifrice**  
**LOLLICAINE PINA COLADA- benzocaine gel, dentifrice**  
**Centrix, Inc.**

-----  
**Lollicaine Topical Anesthetic Gel/Liquid -  
20% Benzocaine**

**Active ingredients**

Benzocaine 20%

**Purpose**

Oral Anesthetic

**Use**

For the temporary relief of occasional minor irritation and pain, associated with:

- Sore mouth and throat
- Ulcers
- Minor dental procedures
- Needle punctures
- Minor irritation of the mouth and gums caused by dentures or orthodontic appliances

**Warnings**

For external use only

**Methemoglobinemia Warning**

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath
- Dizziness or lightheadedness
- Fatigue or lack of energy

**Allergy alert:**

Do not use if you have a history of allergy to any "caine" anesthetics, such as benzocaine, butacaine, procaine or tetracaine.

## **Sore Throat Warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly.

## **Do Not Use**

- For teething
- In children under 2 years of age

## **When using this product**

Avoid contact with eyes

## **Ask a doctor before use**

- If pregnant or breastfeeding

## **Directions**

**Do not exceed recommended dosage.**

**Adults and children 2 years of age and older:** Apply to affected area up to 4 times daily or as directed by dentist or doctor.

**Children under 12 years of age:** Should be supervised in the use of the product.

**Children under 2 years of age:** Do not use

## **Keep out of the reach of children**

## **Other information**

Store at 59° to 86° F (15° to 30° C)

## **Questions or comments?**

800.235.5862 | [centrixdental.com](http://centrixdental.com)

CENTRIX INCORPORATED

770 River Road

Shelton, CT 06484 USA

Toll Free (USA): 800.235.5862

CENTRIXDENTAL.COM

## **Inactive ingredients**

**Bubble Gum flavor contains:** PEG-8, PEG-75, sodium saccharin

**Cherry flavor contains:** Cherry flavor, PEG-8, PEG-75, sodium saccharin

**Mint flavor contains:** Mint flavor, PEG-8, PEG-75, sodium saccharin

**Pina Colada flavor contains:**PEG-8, PEG-75, PinaColada flavor, sodium saccharin

# LolliCaine®

20% Benzocaine  
Topical Anesthetic Gel

**REF 360094**

**Piña Colada**



**Contains: 120 x LolliPack®**  
**0.014 oz. (0.4 g),**  
**0.01 fl. oz. (0.3 ml) per unit**

**centrix®**  
MAKING DENTISTRY EASIER®

# LolliCaine®

20% Benzocaine  
Topical Anesthetic Gel

**REF 360094**

**Piña Colada**

**Contains: 120 x LolliPack®  
0.014 oz. (0.4 g),  
0.01 fl. oz. (0.3 ml) per unit**



**centrix®**  
MAKING DENTISTRY EASIER®

## Drug Facts

Active ingredient (in each unit-dose pack)	Purpose
Benzocaine 20%.....	Oral Anesthetic

**Uses** For the temporary relief of occasional minor irritation and pain associated with:

- Sore mouth and throat
- Ulcers
- Minor dental procedures
- Needle punctures
- Minor irritation of the mouth and gums caused by dentures or orthodontic appliances

### Warnings

For external use only

**Methemoglobinemia warning**  
Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath
- Dizziness or lightheadedness
- Fatigue or lack of energy

### Allergy alert

## Drug Facts (continued)

### Directions

Do not exceed recommended dosage.

Adults and children 2 years of age and older	Apply to affected area up to 4 times daily or as directed by dentist or doctor.
Children under 12 years of age	Should be supervised in the use of the product.
Children under 2 years of age	Do not use.

### Other information

Store at 59° to 86°F (15° to 30°C)

### Inactive ingredients

PEG-8, PEG-75, Piña Colada flavor, sodium saccharin

### Questions?

800.235.5862 | centrixdental.com

Do not use if you have a history or allergy to any "caine" anesthetics, such as benzocaine, butacaine, procaine or tetracaine.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly.

**Do not use**

- For teething
- In children under 2 years of age

**When using this product**

Avoid contact with eyes.

**Ask a doctor before use**

- If pregnant or breastfeeding

**Keep out of reach of children**



**CENTRIX INCORPORATED**

770 River Road

Shelton, CT 06484 USA

Toll Free (USA): 800.235.5862



**CENTRIXDENTAL.COM**



©2025 Centrix, Inc. | 011104 REV J

NDC: 60640-8800-1

**LOT**



# LolliCaine<sup>®</sup>

20% Benzocaine  
Topical Anesthetic Gel

**REF 360092**

**Mint**



**Contains: 120 x LolliPack®**  
**0.014 oz. (0.4 g),**  
**0.01 fl. oz. (0.3 ml) per unit**

**centrix®**  
MAKING DENTISTRY EASIER®

# LolliCaine®

20% Benzocaine  
Topical Anesthetic Gel

**REF 360092**

**Mint**

**Contains: 120 x LolliPack®**  
**0.014 oz. (0.4 g),**  
**0.01 fl. oz. (0.3 ml) per unit**



**centrix®**  
MAKING DENTISTRY EASIER®

## **Drug Facts**

<b>Active ingredient</b> <b>(in each unit-dose pack)</b>	<b>Purpose</b>
Benzocaine 20%.....	Oral Anesthetic

**Uses** For the temporary relief of occasional minor

## **Drug Facts (continued)**

### **Directions**

**Do not exceed recommended dosage.**

Adults and children  
2 years of age  
and older

Apply to affected area  
up to 4 times daily or as  
directed by dentist or doctor.

**Uses** For the temporary relief of occasional minor irritation and pain associated with:

- Sore mouth and throat
- Ulcers
- Minor dental procedures
- Needle punctures
- Minor irritation of the mouth and gums caused by dentures or orthodontic appliances

### Warnings

**For external use only**

#### Methemoglobinemia warning

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath
- Dizziness or lightheadedness
- Fatigue or lack of energy

#### Allergy alert

Do not use if you have a history of allergy to any "caine" anesthetics, such as benzocaine, butacaine, procaine or tetracaine.

#### Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly.

#### Do not use

- For teething
- In children under 2 years of age

#### When using this product

Avoid contact with eyes.

#### Ask a doctor before use

- If pregnant or breastfeeding

**Keep out of reach of children**

Children under 12 years of age

Should be supervised in the use of the product.

Children under 2 years of age

Do not use.

### Other information

Store at 59° to 86°F (15° to 30°C)

### Inactive ingredients

Mint flavor, PEG-8, PEG-75, sodium saccharin

### Questions?

800.235.5862 | [centrixdental.com](http://centrixdental.com)



**CENTRIX INCORPORATED**

770 River Road

Shelton, CT 06484 USA

Toll Free (USA): 800.235.5862



**CENTRIXDENTAL.COM**



©2025 Centrix, Inc. | 011103 REV J

NDC: 60640-8800-1

**LOT**



# LolliCaine<sup>®</sup>

20% Benzocaine  
Topical Anesthetic Gel

**REF 360090**

**Cherry**



**Contains: 120 x LoliPack®**

**0.014 oz. (0.4 g),**

**0.01 fl. oz. (0.3 ml) per unit**

**centrix®**  
MAKING DENTISTRY EASIER®

# LoliCaine®

20% Benzocaine

Topical Anesthetic Gel

**REF 360090**

**Cherry**

**Contains: 120 x LoliPack®**

**0.014 oz. (0.4 g),**



0.01 fl. oz. (0.3 ml) per unit

**centrix**<sup>®</sup>  
MAKING DENTISTRY EASIER<sup>®</sup>

## Drug Facts

<b>Active ingredient</b> <b>(in each unit-dose pack)</b>	<b>Purpose</b>
---	----------------

Benzocaine 20%.....Oral Anesthetic

**Uses** For the temporary relief of occasional minor irritation and pain associated with:

- Sore mouth and throat
- Ulcers
- Minor dental procedures
- Needle punctures
- Minor irritation of the mouth and gums caused by dentures or orthodontic appliances

### Warnings

**For external use only**

#### Methemoglobinemia warning

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath
- Dizziness or lightheadedness
- Fatigue or lack of energy

#### Allergy alert

Do not use if you have a history of allergy to any "caine" anesthetics, such as benzocaine, butacaine, procaine or tetracaine.

#### Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly.

#### Do not use

- For teething
- In children under 2 years of age

#### When using this product

Avoid contact with eyes.

#### Ask a doctor before use

- If pregnant or breastfeeding

**Keep out of reach of children**

## Drug Facts (continued)

### Directions

**Do not exceed recommended dosage.**

Adults and children 2 years of age and older	Apply to affected area up to 4 times daily or as directed by dentist or doctor.
Children under 12 years of age	Should be supervised in the use of the product.
Children under 2 years of age	Do not use.

### Other information

Store at 59° to 86°F (15° to 30°C)

### Inactive ingredients

Cherry flavor, PEG-8, PEG-75, sodium saccharin

### Questions?

800.235.5862 | centrixdental.com

 **CENTRIX INCORPORATED**  
770 River Road  
Shelton, CT 06484 USA  
Toll Free (USA): 800.235.5862

 **CENTRIXDENTAL.COM**





# LolliCaine<sup>®</sup>

20% Benzocaine  
Topical Anesthetic Gel

**REF 360088**

**Bubble Gum**



**Contains: 120 x LolliPack<sup>®</sup>**  
**0.014 oz. (0.4 g),**  
**0.01 fl. oz. (0.3 ml) per unit**

# LolliCaine®

20% Benzocaine  
Topical Anesthetic Gel

**REF 360088**

**Bubble Gum**

**Contains: 120 x LolliPack®**  
**0.014 oz. (0.4 g),**  
**0.01 fl. oz. (0.3 ml) per unit**



**centrix®**  
MAKING DENTISTRY EASIER®

## Drug Facts

<b>Active ingredient</b> <b>(in each unit-dose pack)</b>	<b>Purpose</b>
---	----------------

Benzocaine 20%.....	Oral Anesthetic
---------------------	-----------------

**Uses** For the temporary relief of occasional minor irritation and pain associated with:

- Sore mouth and throat
- Ulcers
- Minor dental procedures
- Needle punctures
- Minor irritation of the mouth and gums caused by dentures or orthodontic appliances

### Warnings

**For external use only**

#### Methemoglobinemia warning

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath

## Drug Facts (continued)

### Directions

**Do not exceed recommended dosage.**

Adults and children 2 years of age and older	Apply to affected area up to 4 times daily or as directed by dentist or doctor.
Children under 12 years of age	Should be supervised in the use of the product.
Children under 2 years of age	Do not use.

### Other information

Store at 59° to 86°F (15° to 30°C)

### Inactive ingredients

Bubble Gum flavor, PEG-8, PEG-75, sodium saccharin

### Questions?

800.235.5862 | centrixdental.com

- Dizziness or lightheadedness
- Fatigue or lack of energy

**Allergy alert**

Do not use if you have a history of allergy to any "caine" anesthetics, such as benzocaine, butacaine, procaine or tetracaine.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, see your dentist or doctor promptly.

**Do not use**

- For teething
- In children under 2 years of age

**When using this product**

Avoid contact with eyes.

**Ask a doctor before use**

- If pregnant or breastfeeding

**Keep out of reach of children**



**CENTRIX INCORPORATED**  
770 River Road  
Shelton, CT 06484 USA  
Toll Free (USA): 800.235.5862



**CENTRIXDENTAL.COM**



©2025 Centrix, Inc. | 011536 REV I

NDC: 60640-8800-1

**LOT**



## LOLLICAINE BUBBLE GUM

benzocaine gel, dentifrice

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:60640-8800
<b>Route of Administration</b>	DENTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>PEG-8</b> (UNII: B697894SGQ)	
<b>PEG-75</b> (UNII: G2M7P15E5P)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60640-8800-1	120 in 1 CARTON	01/14/2019	
1		0.4 g in 1 DOSE PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/14/2019	

**LOLLICAINE CHERRY**

benzocaine gel, dentifrice

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:60640-9000
<b>Route of Administration</b>	DENTAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>PEG-8</b> (UNII: B697894SGQ)	
<b>PEG-75</b> (UNII: G2M7P15E5P)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

**Product Characteristics**

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:60640-9000-1	120 in 1 CARTON	01/14/2019	
1		0.4 g in 1 DOSE PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/14/2019	

## LOLLICAINE MINT

benzocaine gel, dentifrice

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:60640-9200
Route of Administration	DENTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
PEG-8 (UNII: B697894SGQ)	
PEG-75 (UNII: G2M7P15E5P)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

### Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60640-9200-1	120 in 1 CARTON	01/14/2019	
1		0.4 g in 1 DOSE PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/14/2019	

## LOLLICAINE PINA COLADA

benzocaine gel, dentifrice

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:60640-9400
Route of Administration	DENTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>PEG-8</b> (UNII: B697894SGQ)	
<b>PEG-75</b> (UNII: G2M7P15E5P)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

### Product Characteristics

Color	yellow	Score	
Shape		Size	
Flavor	COCONUT	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60640-9400-1	120 in 1 CARTON	01/14/2019	
1		0.4 g in 1 DOSE PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/14/2019	

---

**Labeler** - Centrix, Inc. (053707303)

**Establishment**

Name	Address	ID/FEI	Business Operations
Centrix Inc		053707303	pack(60640-8800, 60640-9000, 60640-9200, 60640-9400) , label(60640-8800, 60640-9000, 60640-9200, 60640-9400)

Revised: 10/2025

Centrix, Inc.