

**ASPIRIN- aspirin tablet, delayed release**  
**Topco Associates, LLC**

-----  
**TopCare 44-645**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Do not use**

if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments?**

**1-888-423-0139**

**Principal Display Panel**

NDC 36800-545-60

**TopCare  
health®**

COMPARE TO THE ACTIVE INGREDIENT  
IN ST. JOSEPH® LOW DOSE ASPIRIN†

LOW DOSE ♥ ASPIRIN REGIMEN\*\*

**Aspirin 81 mg**  
**PAIN RELIEVER (NSAID)**

Safety Coated

\*\*Talk to your doctor or other healthcare provider  
before using this product for your heart.

actual size

**180 ENTERIC COATED TABLETS**

†This product is not manufactured or distributed by Foundation  
Consumer Healthcare, LLC, owner of the registered trademark  
St. Joseph® Low Dose Aspirin.

50844 REV1024B64560

DISTRIBUTED BY  
TOPCO ASSOCIATES LLC  
ITASCA, IL 60143  
©TOPCO LNKA0125

QUESTIONS?

1-888-423-0139

topcare@topco.com

www.topcarebrand.com

This TopCare® product is laboratory  
tested to guarantee its highest quality.  
Your total satisfaction is guaranteed.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**



NDC 36800-545-60

COMPARE TO THE ACTIVE INGREDIENT IN ST. JOSEPH® LOW DOSE ASPIRIN\*

LOW DOSE ♥ ASPIRIN REGIMEN\*\*

# Aspirin 81 mg

PAIN RELIEVER (NSAID)  
Safety Coated

\*Talk to your doctor or other healthcare provider before using this product for your heart.

180 ENTERIC COATED TABLETS

actual size



NDC 36800-545-60

COMPARE TO THE ACTIVE INGREDIENT IN ST. JOSEPH® LOW DOSE ASPIRIN\*

LOW DOSE ♥ ASPIRIN REGIMEN\*\*

# Aspirin 81 mg

PAIN RELIEVER (NSAID)  
Safety Coated

\*Talk to your doctor or other healthcare provider before using this product for your heart.

180 ENTERIC COATED TABLETS

actual size

This product is not manufactured or distributed by Foundation Consumer Healthcare, LLC, owner of the registered trademark St. Joseph® Low Dose Aspirin.  
50844 REV1024B64560



DISTRIBUTED BY  
TOPCO ASSOCIATES LLC  
ITASCA, IL 60143  
©TOPCO LNK A0125  
QUESTIONS?  
1-888-423-0139  
topcare@topco.com  
www.topcarebrand.com

Scan here for more information or call 1-888-423-0139

**QUALITY GUARANTEED**  
This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

No Varnish/No Print Lot # & Expiry Area



0 36800 110953 7

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

B-1910-645-60-R  
REV1024B64560

### Drug Facts (continued)

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - ringing in the ears or a loss of hearing occurs
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

do not take more than directed

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### Other information

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- see end flap for expiration date and lot number

### Inactive ingredients

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hydroxyethylcellulose, hydroxypropylcellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

### Questions or comments? 1-888-423-0139

### KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

### Drug Facts

#### Active ingredient (in each tablet)

Aspirin 81 mg (NSAID).....Pain reliever  
nonsteroidal anti-inflammatory drug

**Uses** for the temporary relief of minor aches and pains or as recommended by your doctor

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- chest
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- diabetes
- gout
- arthritis

Topcare 44-645

## ASPIRIN

aspirin tablet, delayed release

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:36800-545

Route of Administration

ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

**Product Characteristics**

Color	pink	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-545-60	1 in 1 CARTON	03/19/2021	
1		180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/19/2021	

**Labeler** - Topco Associates, LLC (006935977)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-545)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-545)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(36800-545)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(36800-545)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(36800-545)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(36800-545)

Revised: 5/2025

Topco Associates, LLC