

**DRX CHOICE FIBER THERAPY- methylcellulose powder, for solution
RARITAN PHARMACEUTICALS INC**

DRx Choice Fiber Therapy Methylcellulose 2 g Bulk Forming Laxative

***Active ingredient*(in each rounded tablespoon)**

Methylcellulose (a non-allergenic fiber) 2g

Purpose

Bulk-forming fiber laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 12-72 hours

Warnings

Choking: taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- a sudden change in bowel habits that persists for two weeks
- abdominal pain, nausea or vomiting

Stop use and ask a doctor if

- constipation lasts more than 7 days
- you have rectal bleeding

These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

- **MIX THIS PRODUCT (CHILD OR ADULT DOSE) WITH AT LEAST 8 OUNCES (A FULL GLASS) OF WATER OR OTHER FLUID. TAKING THIS PRODUCT WITHOUT ENOUGH LIQUID MAY CAUSE CHOKING. SEE CHOKING WARNING**
- use product at the first sign of constipation or irregularity
- put one dose in a full glass of cold water

- stir briskly and drink promptly
- drinking another glass of water is helpful

Age	Dose
adults & children 12 years of age and over	start with 1 rounded tablespoon . Increase as needed, 1 rounded tablespoon at a time, up to 3 times per day.
children 6 - 11 years of age	start with 2 level teaspoons . Increase as needed, 2 level teaspoons at a time, up to 3 times per day.
children under 6 years of age	consult a physician

Other information

- **each rounded tablespoon contains:** calcium 85mg and potassium 125mg
- each rounded tablespoon contributes 27 calories from maltodextrin
- store at room temperature
- protect contents from humidity
- keep tightly closed

Inactive ingredients

Anhydrous citric acid, FD&C yellow #6 lake, flavor, maltodextrin, potassium citrate, silica, sucralose, tricalcium phosphate,

Questions or comments?

1-866-467-2748

Principal Display Panel

DRx Choice®

Compare to the Active Ingredient in Citrucel® Powder*

NDC 68163-517-16

Sugar free

Soluble fiber therapy

Methylcellulose Fiber Therapy for Irregularity

- **Gentle and clinically proven effective****
- **Gives you additional fiber to help relieve occasional constipation**

Fiber for irregularity that won't cause excess gas***

Orange Flavor

Naturally & Artificially Flavored

NET WT 16.9 OZ (479g)

Directions for use:

1. Fill glass with at least 8 oz. of cold water.
2. Measure rounded tablespoon of product (one dose) and stir product into water. Stir briskly and drink promptly.

**When Used As Directed.

***Based On Laboratory Testing. Individual Results may Vary

TAMPER EVIDENT: DO NOT USE PRINTED INNER SEAL IS BROKEN OR MISSING

GLUTEN FREE

*This product is not manufactured or distributed by GSK Consumer Healthcare, the distributor of Citrucel® Powder.

Manufactured By:

Raritan Pharmaceuticals

8 Joana Court,

East Brunswick, NJ 08816

Compare to the active ingredient in Citrucel® Powder*
NDC 68163-51716



sugar free

soluble fiber therapy

Methylcellulose
Fiber Therapy for Irregularity

Gentle and clinically proven effective**
Gives you additional fiber to help
relieve occasional constipation
Fiber for irregularity that won't cause excess gas***

Orange

NET WT. 16.9 OZ (479g)

TAMPER EVIDENT: DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each rounded tablespoon)	Purpose
Methylcellulose (a non-allergenic fiber) 2g	Bulk-forming fiber laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 12-72 hours

Warnings

Choking: taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have a difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- a sudden change in bowel habits that persists for two weeks
- abdominal pain, nausea or vomiting

Stop use and ask a doctor if

- constipation lasts more than 7 days
- you have rectal bleeding.

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

- MIX THIS PRODUCT (CHILD OR ADULT DOSE) WITH AT LEAST 8 OUNCES (A FULL GLASS) OF WATER OR OTHER FLUID. TAKING THIS PRODUCT WITHOUT ENOUGH LIQUID MAY CAUSE CHOKING. SEE CHOKING WARNING.
- use product at the first sign of constipation or irregularity
- put one dose in a full glass of cold water
- stir briskly and drink promptly
- drinking another glass of water is helpful

Age	Dose
adults and children 12 years of age and over	start with 1 rounded tablespoon . Increase as needed, 1 rounded tablespoon at a time, up to 3 times per day.
children 6-11 years of age	start with 2 level teaspoons . Increase as needed, 2 level teaspoons at a time, up to 3 times per day.
children under 6 years of age	consult a physician

Other information

- each rounded **tablespoon** contains: calcium 85 mg; potassium 125 mg
- each rounded **tablespoon** contributes 27 calories from maltodextrin
- store at room temperature
- protect contents from humidity
- keep tightly closed

Inactive ingredients anhydrous citric acid, FD&C Yellow No. 6 Lake, flavor, maltodextrin, potassium citrate, silica, sucralose, tricalcium phosphate

Questions or comments? 1-866-467-2748

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Citrucel® Powder.

DIRECTIONS FOR USE (NO SCOOP DOSING):



1. Fill glass with at least 8 oz. of cold water.



2. Measure rounded tablespoon of product (one dose) and stir product into water. Stir briskly and drink promptly.

**When Used As Directed.

***Based On Laboratory Testing. Individual Results May Vary.

LOT: 5176DCLR

THIS IS TO DESIGNATE
NON VARNISH AREA.
DO NOT PRINT.

EXP:

Manufactured by:
Raritan Pharmaceuticals
8 Joana Court,
East Brunswick, NJ 08816



3 68163 51716 3

DRX CHOICE FIBER THERAPY

methylcellulose powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68163-517	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
METHYLCELLULOSE (4000 MPA.S) (UNII: MRJ667KA5E) (METHYLCELLULOSE (4000 MPA.S) - UNII:MRJ667KA5E)		METHYLCELLULOSE (4000 MPA.S)	2 g in 10.2 g	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
POTASSIUM CITRATE (UNII: EE90ONI6FF)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
TRICALCIUM PHOSPHATE (UNII: K4C08XP666)				
Product Characteristics				
Color	orange	Score		
Shape		Size		
Flavor	ORANGE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68163-517-16	479 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/18/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	07/18/2022		

Labeler - RARITAN PHARMACEUTICALS INC (127602287)