

LOW DOSE ASPIRIN ENTERIC SAFETY-COATED- aspirin tablet Fred's Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for safety coated 81 mg aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

Other information

- store between 15-30°C (59-86°F)

Inactive ingredients

D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch (corn), silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triethyl citrate

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to active ingredient in Bayer® Low Dose Aspirin†

Low Dose ASPIRIN

Pain reliever (NSAID)

≠Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.

- Aspirin Regimen≠
- Safety Coated≠≠

TABLETS ENTERIC COATED

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin

≠≠Coating helps protect against stomach upset

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY fred's, Inc.

4300 NEW GETWELL RD., MEMPHIS, TN 38118

www.fredsinc.com

Product Label

fred's Pharmacy
NDC 55315-867-30
Compare to the active ingredient in Bayer® Low Dose Aspirin®

Low Dose ASPIRIN
Pain Reliever (NSAID)

ACTUAL SIZE

81mg each

• Aspirin Regimen®
• Safety Coated™
300 TABLETS ENTERIC COATED

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

†Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.

TAMPER EVIDENT DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active Ingredient (in each tablet)
Aspirin 81 mg (NSAID)[†]
"nonsteroidal anti-inflammatory drug"

Purpose
Pain reliever

Uses
■ for the temporary relief of minor aches and pains or as recommended by your doctor.
■ Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
■ ask your doctor about other uses for safety coated 81 mg aspirin.

Warnings
Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had past stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn

Drug Facts (continued under label)

100% Satisfaction Guarantee or your money back
1-855-331-FRED (37238)

6300 NEW GETTYSBURG RD., MEMPHIS, TN 38118
www.freds.com

Exp. Date: 01845791122349

PLD-06674 LB005546

PEEL HERE

Drug Facts (continued)

■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic ■ you have asthma
Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days ■ redness or swelling is present
■ ringing in the ears or loss of hearing occurs ■ new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ Drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information

■ Store between 15-30°C (59-86°F)

Inactive ingredients D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch (corn), silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, stearic acid, titanium dioxide, triethyl citrate

Questions or comments?
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.
*Coating helps protect against stomach upset

FRED'S PHARMACY Low Dose Aspirin

LOW DOSE ASPIRIN ENTERIC SAFETY-COATED

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55315-867
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	E
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55315-867-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2019	03/31/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/31/2019	03/31/2025

Labeler - Freds Inc (005866116)

Revised: 10/2022

Freds Inc