

**COLD AND FLU RELIEF SEVERE DAYTIME- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated**  
**Rite Aid Corporation**

-----  
**Rite Aid 44-640**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - sinus congestion and pressure
  - sore throat
  - fever
  - headache
  - nasal congestion
  - cough due to minor throat and bronchial irritation
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters

- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

### **When using this product**

**do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

### ***Other information***

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions or comments?***

**1-800-426-9391**

### ***Principal Display Panel***

11822-0640-8

Compare to the active ingredients in **Vicks®**

**DayQuil® VapoCOOL®**

**SEVERE COLD & FLU**

**+ CONGESTION\***

**MAXIMUM STRENGTH**

**SEVERE**

**COLD & FLU RELIEF**

DAY

**ACETAMINOPHEN**

PAIN RELIEVER / FEVER REDUCER

**DEXTROMETHORPHAN HBr**  
COUGH SUPPRESSANT  
**GUAIFENESIN • PHENYLEPHRINE HCl**  
EXPECTORANT • NASAL DECONGESTANT

Temporarily relieves headache, fever,  
sore throat, minor aches & pains, nasal/sinus  
congestion & sinus pressure,  
cough, chest congestion

ACTUAL SIZE

**24**  
CAPLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER  
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by The Procter & Gamble  
Company, owner of the registered trademark Vicks<sup>®</sup> DayQuil<sup>®</sup> VapoCOOL<sup>®</sup>  
SEVERE COLD & FLU + CONGESTION.

50844 REV0722C64008

**DISTRIBUTED BY:** RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319  
**[www.riteaid.com](http://www.riteaid.com)**

**SATISFACTION  
GUARANTEE**

If you're not satisfied, we'll  
happily refund your money.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

Compare to the active ingredients in Vicks® DayQuil® VapoCOOL® SEVERE COLD & FLU + CONGESTION\*

NDC 11822-0640-8

**MAXIMUM STRENGTH**

# SEVERE COLD & FLU RELIEF



**ACETAMINOPHEN**  
PAIN RELIEVER / FEVER REDUCER  
**DEXTROMETHORPHAN HBr**  
COUGH SUPPRESSANT  
**GUAIFENESIN • PHENYLEPHRINE HCl**  
EXPECTORANT • NASAL DECONGESTANT  
Temporarily relieves headache, fever, sore throat, minor aches & pains, nasal/sinus congestion & sinus pressure, cough, chest congestion



**24**  
CAPLETS



**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR PLASTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

no print / no varnish area for stock exp. dated

**PARENTS:**  
Learn about them medicine abuse  
[www.StupidDrugs.com](http://www.StupidDrugs.com)

**Drug Facts**  
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION  
**Active ingredients (in each caplet)**  
Acetaminophen 325 mg, Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg, Cough suppressant  
Guaifenesin 200 mg, Expectorant  
Phenylephrine HCl 5 mg, Nasal decongestant

**Drug Facts (continued)**  
**Uses** ■ temporarily relieves common cold and flu symptoms: ■ minor aches and pains ■ sinus congestion and pressure ■ sore throat ■ fever ■ headache ■ nasal congestion ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ temporarily relieves throat irritation through the nose ■ promotes nasal and/or sinus drainage  
**Drug Facts (continued)**

**Drug Facts (continued)**  
**Inactive ingredients** corn starch, croscollon, FD&C red #40 aluminum lake, F&C yellow #6 aluminum lake, magnesium stearate, methylcellulose, polyethylene glycol, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide  
**Questions or comments?** 1-800-426-9391

**Drug Facts (continued)**  
This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® VapoCOOL® SEVERE COLD & FLU + CONGESTION. 50844 REV0722C64008  
**DISTRIBUTED BY: RITE AID,**  
200 NEUBERRY COMMONS  
EATERS, PA 17319  
[www.riteaid.com](http://www.riteaid.com)

8-1702-640-08-R3  
REV0722C64008

**Drug Facts (continued)**  
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive  
**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin redness  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients  
**Ask a doctor before use if you have**  
■ liver disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ heart disease  
■ difficulty in urination due to enlargement of the prostate gland

**Drug Facts (continued)**  
**Stop use and ask a doctor if:**  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions** ■ do not take more than directed  
do not take more than 8 caplets in 24 hours  
adults and children 12 years and over: take 2 caplets with water every 4 hours  
children under 12 years: ask a doctor  
**Other information** ■ each caplet contains: sodium 3 mg  
**BLISTER IS TORN OR BROKEN**  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ see end flap for expiration date and lot number

no print / no varnish area

no print / no varnish area

Rite Aid 44-640

**COLD AND FLU RELIEF SEVERE DAYTIME**

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0640
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40 ALUMINUM LAKE</b> (UNII: 6T47AS764T)	
<b>FD&amp;C YELLOW NO. 6 ALUMINUM LAKE</b> (UNII: GYP6Z2JR6Q)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;640
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0640-8	2 in 1 CARTON	11/25/2019	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/25/2019	

**Labeler** - Rite Aid Corporation (014578892)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0640) , pack(11822-0640)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-0640)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0640)

Revised: 5/2026

Rite Aid Corporation