SENNA LAXATIVE- sennosides tablet TOP CARE (Topco Associates LLC)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Sennosides 8.6 mg

Purpose

Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 6-12 hours.

Warnings

Do not use

• laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before using if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks.

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• take preferably at bedtime or as directed by a doctor

age	starting dosage	maximum dosage
adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

Other information

-

- each tablet contains: calcium 25 mg
- store at 25°C (77° F); excursions permitted between 15–30°C (59–86°F)

Inactive Ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, hypromellose, liquid paraffin*, magnesium stearate, maltodextrin*, microcrystalline cellulose, mineral oil*, polyethylene glycol*, polyvinyl alcohol*, silicon dioxide*, sodium lauryl sulfate*, stearic acid*, talc*

*contains one or more of these ingredients

Questions or comments?

Call toll free 1-888-423-0139

Principal Display Panel

COMPARE TO SENOKOT® ACTIVE INGREDIENT†

Vegetable Laxative

SENNOSIDES 8.6 mg • LAXATIVE

Gentle, Overnight Relief of Constipation

TABLETS

GLUTEN FREE

†This product is not manufactured or distributed by Purdue Products L.P., distributor of Senokot®.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007

topcare@topco.com www.topcarebrand.com

Product Label

+ TopCare.

SENNOSIDES 8.6 mg • LAXATIVE

Vegetable Laxative

Gentle, Overnight Relief of Constipation

NDC 36800-786-01

TOPCARE HEALTH Vegetable Laxative

		Com		
IOO TABLE	TS	actual size	0	
NOITAMROTNI TOUOOR9 GNA SONING		DESTNARAUS		
EP OUTER CARTON FOR COMPLETE		moo.bnerdeneoqot.www		
e under cap is broken or missing. Evident: do not use if printed safety	VILLAGE, IL 60007	8 ODESTIONS? 1-888-423-0139 06C0 ASSOCIATES LLC, ELK GROVE	DISTRIBUTED BY TO 00901 @	
f Serokof®,		r comments? Call toll free 1- manufactured or distributed by Pu	SAM NOT IGN AND	
		and the second	No. of Concession, name	
ihydrate, hypromellose", liquid paratlin", ylene glycol", polyvinyl alcohol", silicon	n, dibasic calcium phosphale d cellulose, mineral oil*, polyethy	redients croscamellose sodiur ate, maltodextrin*, microcrystalline lauryl sulfate*, stearic acid*, talc* nore of these ingredients	magnesium stear muibos ,*abixoib	
	(7°88-92) J°05-21 ns	rtion Intains: calcium 25 mg 77°F); excursions permitted betwee		
		(panuluoo)) stasi Pung	
				
ssk a doctor	ask a doctor	st 2 years	children unde	
1 tablet twice a day ask a doctor	1/2 tablet once a day ask a doctor	under 6 years ar 2 years		
			of S nenblinta	
t tablet twice a day	1/2 tablet once a day	nuqet 6 years	children 6 to children 2 to	
Z tablet twice a day T tablet twice a day	1 tablet once a day 1/2 tablet once a day	under 12 years under 6 years	children 6 to children 2 to	
4 tablets twice a day 2 tablets twice a day 1 tablet twice a day	starting dosage 2 tablet once a day 1/2 tablet once a day 1/2 tablet once a day	under 12 years under 6 years	age children 6 to children 2 to	
t tablets twice a day 2 tablets twice a day 2 tablets twice a day	e, get medical help or contact a starting dosage 2 tablet once a day 1/2 tablet once a day 1/2 tablet once a day	ich of children. In case of overdos ly at bedtime or as directed by a do ildren 12 years of age and over under 12 years	Keep out of res Directions age children 6 to children 6 to children 2 to	
agesob mumixem 4 tablets twice a day 2 tablet twice a day 7 tablet twice a day	e, get medical help or contact a starting dosage 2 tablet once a day 1/2 tablet once a day 1/2 tablet once a day	brest-feeding, ask a health profe och of children. In case of overdos ily at bedtime or as directed by a do ildren 12 years under 6 years	If pregnant or tree Keep out of res I take preferab age adults and ch children 6 to children 2 to	
i Poison Control Center right away. maximum dosage 4 tablets twice a day 2 tablets twice a day 1 tablet twice a day	e, get medical help or contact a cotor starting dosage 2 tablet once a day 1 tablet once a day 1/2 tablet once a day	us condition. breat-feeding, ask a health profe ly at bedtime or as directed by a do ildren 12 years of age and over conder 12 years under 6 years	Indicate a serior If pregrant or Keep out of rea atke preferab age adults and ch children 6 to children 6 to children 2 to	
wement after use of a laxative. These ma Poison Control Center right away. 4 tablets twice a day 2 tablets twice a day 1 tablet twice a day	ading or fail to have a bowel mo ssional before use. e, get medical help or contact a starting dosage 2 tablet once a day 1 tablet once a day 1/2 tablet once a day	us condition. breat-feeding, ask a health profe ly at bedtime or as directed by a do ildren 12 years of age and over conder 12 years under 6 years	Indicate a sud Stop use and a Stop use areior If pregnant or io Meep out of rea able preferab age adults and ch children 6 to children 6 to children 7 to children 8 to children	
wement after use of a laxative. These ma Poison Control Center right away. 4 tablets twice a day 2 tablets twice a day 7 tablets twice a day 7 tablet twice a day	ontinues over a period of 2 wee ading or fail to have a bowel mo ssional before use. e, get medical help or contact a starting dosage 1 tablet once a day 1 tablet once a day 12 tablet once a day	efore use if you have den change in bowel habits that or den change in bowel habits that or us condition. Inder 12 years of age and over och of children. In case of overdos ch of children. In case of overdos of of children. In case of over overdos of of children. In case of over overdos of of children. In case of over overdos of of children. In case of of children. In case of of children.	Ask a doctor b a stomach pain a tomach aud Stop use and a indicate a serior indicate a serior indicate a serior indicate a serior a stop preterab age age children 6 to children 6 to children 6 to	
wement after use of a laxative. These ma Poison Control Center right away. 4 tablets twice a day 2 tablets twice a day 1 tablet twice a day	ontinues over a period of 2 wee ading or fail to have a bowel mo ssional before use. e, get medical help or contact a starting dosage 1 tablet once a day 1 tablet once a day 12 tablet once a day	Inder change in bowel habits that co den change in bowel habits that co us condition. Is a doctor if you have rectal blee breast-feeding, ask a health profection. Inder 12 years of age and over or of children. In case of overdos of of children. In case of overdos Inder 12 years	Ask a doctor b a stomach pain a tomach aud Stop use and a indicate a serior indicate a serior indicate a serior indicate a serior a stop preterab age age children 6 to children 6 to children 6 to	
wement after use of a laxative. These ma Poison Control Center right away. 4 tablets twice a day 2 tablets twice a day 1 tablet twice a day	directed by a doctor ading or fail to have a bowel mo scional before use. signal before use. 2 tablet once a day 1 tablet once a day 1 tablet once a day 1 tablet once a day	efore use if you have den change in bowel habits that or den change in bowel habits that or us condition. Inder 12 years of age and over och of children. In case of overdos ch of children. In case of overdos of of children. In case of over overdos of of children. In case of over overdos of of children. In case of over overdos of of children. In case of of children. In case of of children.	generally pro generally pro bon not use bon not use bon not use bon servio bongues and a bongues and a indicate a servio forgenant or forgenant or for	
ks wement after use of a laxative. These ma Poison Control Center right away. 4 tablets twice a day 2 tablets twice a day 1 tablets twice a day	directed by a doctor ading or fail to have a bowel mo scional before use. signal before use. 2 tablet once a day 1 tablet once a day 1 tablet once a day 1 tablet once a day	under 6 years under 12 years bedre a bowel movement in 6 -12 ucts for longer than 1 week unless efore use if you have a condition. Inder 12 years of age and over breast-feeding, ask a health prote us condition. Desst-feeding, ask a health prote best-feeding, ask a health protecte best-feeding, ask a health p	Sennosides 8.6 Uses Uses Uses Uses Uses Uses Uses Use	

COMPARE TO SENOKOT® TICLO So. Date: Tot No:

Ρ							
	roduct Info	rmation					
Pı	roduct Type		HUMAN OTC DRUG	Item Code	(Source)	NDC:36	6800-786
Re	oute of Admin	istration	ORAL				
A	ctive Ingred	lient/Active	Moiety				
	J		dient Name		Basis of Str	enath	Strength
SE	NNOSIDES (UN	-	(SENNOSIDES - UNII:3FYP5	MOIJX)	SENNOSIDES		8.6 mg
				-			
In	active Ingr	edients					
			Ingredient Nam	е			Strength
CF	ROSCARMELLOS	SE SODIUM (U	NII: M28OL1HH48)				
			DIHYDRATE (UNII: 07TS)	Z97GEP)			
ΗY	PROMELLOSES	3 (UNII: 3NXW29	V3WO)				
	AGNESIUM STE	•	•				
	ALTODEXTRIN (
			NE (UNII: OP1R32D61U)				
М	NERAL OIL (UNI	I: T5L8T28FGP)					
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)							
STEARIC ACID (UNII: 4ELV7Z65AP)							
TALC (UNII: 7SEV7J4R1U)							
		•					
sc	DIUM LAURYL	SULFATE (UNII					
SC PC	DIUM LAURYL	SULFATE (UNII Glycol, UNSF	ECIFIED (UNII: 3WJQ0SDV	V1A)			
SC PC	DIUM LAURYL	SULFATE (UNII Glycol, UNSF		V1A)			
SC PC	DIUM LAURYL	SULFATE (UNII Glycol, UNSF	ECIFIED (UNII: 3WJQ0SDV	V1A)			
SC PC PC	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC	PECIFIED (UNII: 3WJQ0SDV IFIED (UNII: 532B59J990)	V1A)			
SC PC PC	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC	PECIFIED (UNII: 3WQOSDV IFIED (UNII: 532B59J990)	VIA)	no score		
SC PC PC	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown	ECIFIED (UNII: 3WQ0SDV IFIED (UNII: 532B59J990) Score	V1A)	no score 9mm		
SC PC PC	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char Dlor nape	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC	ECIFIED (UNII: 3WQOSDV IFIED (UNII: 532B59J990) Score Size	VIA)	9mm	9	
SC PC PC PC St Fla	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char Dor nape avor	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown	ECIFIED (UNII: 3WQ0SDV IFIED (UNII: 532B59J990) Score	V1A)		8	
SC PC PC PC St Fla	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char Dlor nape	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown	ECIFIED (UNII: 3WQOSDV IFIED (UNII: 532B59J990) Score Size	V1A)	9mm	8	
SC PC PC PC St Fla	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char Dor nape avor	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown	ECIFIED (UNII: 3WQOSDV IFIED (UNII: 532B59J990) Score Size	VIA)	9mm	8	
SC PC PC Cc Sh Fla Cc	DDIUM LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char Dor nape avor	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown	ECIFIED (UNII: 3WQOSDV IFIED (UNII: 532B59J990) Score Size		9mm 0813;AV;TCL080;S		
SC PC PC ST Fla Cc	odium LAURYL OLYETHYLENE (OLYVINYL ALCO roduct Char olor hape avor ontains ackaging Item Code	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown ROUND	ECIFIED (UNII: 3WQOSDV IFIED (UNII: 532B59J990) Score Size		9mm		keting End Date
SC PC PC ST Fla Cc	odium LAURYL OLYETHYLENE (OLYVINYL ALCO roduct Char olor hape avor ontains ackaging	SULFATE (UNII GLYCOL, UNSF HOL, UNSPEC Cacteristics brown ROUND	PECIFIED (UNII: 3WQOSDW IFIED (UNII: 532B59J990) Score Size Imprint Code		9mm 0813;AV;TCL080;S Marketing Start		keting End Date
SC PC PC SI Fla Cc Pa #	odium LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char blor hape avor ontains ackaging item Code NDC:36800-	SULFATE (UNII GLYCOL, UNSP HOL, UNSPEC Cacteristics brown ROUND	PECIFIED (UNII: 3WQOSDW IFIED (UNII: 532B59J990) Score Size Imprint Code	P 01	9mm 0813;AV;TCL080;S Marketing Start Date		-
SC PC PC Sh Fla Cc Pa #	odium LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char blor hape avor ontains ackaging item Code NDC:36800-	SULFATE (UNII GLYCOL, UNSP HOL, UNSPEC Cacteristics brown ROUND	PECIFIED (UNII: 3WQOSDW IFIED (UNII: 532B59J990) Score Size Imprint Code	P 01	9mm 0813;AV;TCL080;S Marketing Start Date		-
SC PC PC SI Fla Cc Pa # 1	odium LAURYL DLYETHYLENE (DLYVINYL ALCO roduct Char blor hape avor ontains ackaging item Code NDC:36800-	SULFATE (UNII GLYCOL, UNSP HOL, UNSPEC Cacteristics brown ROUND	PECIFIED (UNII: 3WQOSDW IFIED (UNII: 532B59J990) Score Size Imprint Code Package Description	P 01	9mm 0813;AV;TCL080;S Marketing Start Date		-

OTC monograph not final	part334	01/31/2016	

Labeler - TOP CARE (Topco Associates LLC) (006935977)

Revised: 11/2022

TOP CARE (Topco Associates LLC)