

**IBUPROFEN- ibuprofen tablet, film coated**  
**L.N.K. International, Inc.**

-----  
**Quality Plus 44-291**

***Active ingredient (in each brown tablet)***

Ibuprofen USP, 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- skin reddening
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - weakness in one part or side of body
  - leg swelling
  - slurred speech
  - trouble breathing
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear
- redness or swelling is present in the painful area

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

***Principal Display Panel***

**Quality**

**+Plus**

NDC 50844-291-16

†Compare to active ingredient  
in Advil® Tablets

**IBUPROFEN TABLETS**

**USP, 200 mg**

PAIN RELIEVER/  
FEVER REDUCER (**NSAID**)

**1000 Coated Tablets**

**ACTUAL**

# SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Tablets. 50844 REV1221C29116

Distributed by  
**LNK INTERNATIONAL, INC.**  
 60 Arkay Drive, Hauppauge, NY 11788  
 USA

**QUALITY PLUS** NDC 50844-291-16  
 †Compare to active ingredient in Advil® Tablets

**IBUPROFEN TABLETS USP, 200 mg**  
 PAIN RELIEVER/  
 FEVER REDUCER (NSAID)

**1000 Coated Tablets** ACTUAL SIZE

**Drug Facts** TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Active ingredient (in each brown tablet)** Purpose  
 Ibuprofen USP, 200 mg (NSAID)\* ..... Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ toothache ■ backache ■ menstrual cramps  
 ■ the common cold ■ muscular aches ■ minor pain of arthritis  
 ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling  
 ■ asthma (wheezing) ■ skin reddening ■ shock ■ rash ■ blisters  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
 ■ under a doctor's care for any serious condition ■ taking any other drug

**Drug Facts (continued)**  
**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ you have symptoms of heart problems or stroke:  
 ■ chest pain ■ weakness in one part or side of body ■ leg swelling  
 ■ slurred speech ■ trouble breathing  
 ■ pain gets worse or lasts more than 10 days ■ any new symptoms appear  
 ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed  
 ■ the smallest effective dose should be used  
 ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist  
 ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used  
 ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor  
 ■ children under 12 years: ask a doctor

**Other information** ■ store between 20°-25°C (68°-77°F)  
 ■ avoid excessive heat 40°C (104°F) ■ use by expiration date on package

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Tablets.  
 50844 REV1221C29116  
 Distributed by  
**LNK INTERNATIONAL, INC.**  
 60 Arkay Drive, Hauppauge, NY 11788  
 USA

**Quality Plus 44-291**

## IBUPROFEN

ibuprofen tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-291
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-291-50	50 in 1 CARTON	05/24/1988	
1	NDC:50844-291-24	2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:50844-291-03	10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
3	NDC:50844-291-02	1 in 1 CARTON	05/24/1988	
3		12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:50844-291-08	1 in 1 CARTON	05/24/1988	
4		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:50844-291-01	1 in 1 CARTON	05/24/1988	
5		30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:50844-291-10	1 in 1 CARTON	05/24/1988	
6		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
7	NDC:50844-291-15	1 in 1 CARTON	05/24/1988	
7		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
8	NDC:50844-291-11	1 in 1 CARTON	05/24/1988	
8		60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
9	NDC:50844-291-12	1 in 1 CARTON	05/24/1988	

9		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
10	NDC:50844-291-29	1 in 1 CARTON	05/24/1988	
10		150 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
11	NDC:50844-291-13	1 in 1 CARTON	05/24/1988	
11		250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
12	NDC:50844-291-99	250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
13	NDC:50844-291-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
14	NDC:50844-291-16	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
15	NDC:50844-291-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

**Labeler** - L.N.K. International, Inc. (038154464)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-291)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-291)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-291)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-291) , pack(50844-291)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-291)

## Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		117025878	manufacture(50844-291)
-------------------------	--	-----------	------------------------

Revised: 6/2024

L.N.K. International, Inc.