

# CLINDAMYCIN PHOSPHATE- clindamycin phosphate solution

## Viona Pharmaceuticals Inc


### Clindamycin Phosphate Topical Solution USP, 1%

#### DESCRIPTION

Clindamycin phosphate topical solution USP, 1% contains clindamycin phosphate, USP, at a concentration equivalent to 10 mg clindamycin per milliliter.

Clindamycin phosphate is a water soluble ester of the semi-synthetic antibacterial produced by a 7(S)-chloro-substitution of the 7(R)-hydroxyl group of the parent compound lincomycin.

The structural formula is represented below:



The chemical name for clindamycin phosphate is Methyl 7-chloro-6,7,8-trideoxy-6-(1-methyl-*trans*-4-propyl-L-2-pyrrolidinecarboxamido)-1-thio-L-*threo*- $\alpha$ -D-galactooctopyranoside 2-(dihydrogen phosphate).

Clindamycin phosphate topical solution USP, 1% contains isopropyl alcohol, 50% v/v; propylene glycol and purified water. Sodium hydroxide or hydrochloric acid may be added to adjust pH between 4.0 to 7.0.

#### CLINICAL PHARMACOLOGY

##### Mechanism of Action

The mechanism of action of clindamycin in treating acne vulgaris is unknown.

##### Pharmacokinetics

Following multiple topical applications of clindamycin phosphate at a concentration

equivalent to 10 mg clindamycin per mL in an isopropyl alcohol and water solution, very low levels of clindamycin are present in the serum (0 to 3 ng/mL) and less than 0.2% of the dose is recovered in urine as clindamycin.

Although clindamycin phosphate is inactive *in vitro*, rapid *in vivo* hydrolysis converts this compound to the antibacterially active clindamycin.

## **Microbiology**

Clindamycin inhibits bacterial protein synthesis by binding to the 23S RNA of the 50S subunit of the ribosome. Clindamycin is bacteriostatic.

## **Antimicrobial Activity**

Clindamycin is active *in vitro* against most isolates of *Propionibacterium acnes*; however, the clinical significance is unknown.

## **Resistance**

Resistance to clindamycin is most often caused by modification of specific bases of the 23S ribosomal RNA. Cross-resistance between clindamycin and lincomycin is complete. Because the binding sites for these antibacterial drugs overlap, cross resistance is sometimes observed among lincosamides, macrolides and streptogramin B. Macrolide-inducible resistance to clindamycin occurs in some isolates of macrolide-resistant bacteria.

## **INDICATIONS AND USAGE**

Clindamycin phosphate topical solution USP, 1% is indicated in the treatment of acne vulgaris. In view of the potential for diarrhea, bloody diarrhea and pseudomembranous colitis, the physician should consider whether other agents are more appropriate (see CONTRAINDICATIONS, WARNINGS and ADVERSE REACTIONS).

## **CONTRAINDICATIONS**

Clindamycin phosphate topical solution is contraindicated in individuals with a history of hypersensitivity to preparations containing clindamycin or lincomycin, a history of regional enteritis or ulcerative colitis, or a history of antibiotic-associated colitis.

## **WARNINGS**

**Orally and parenterally administered clindamycin has been associated with severe colitis which may result in patient death. Use of the topical formulation of clindamycin results in systemic absorption from the skin surface. Diarrhea, bloody diarrhea, and colitis (including pseudomembranous colitis) have been reported with the use of topical and systemic clindamycin.**

**Studies indicate a toxin(s) produced by clostridia is one primary cause of antibiotic-associated colitis. The colitis is usually characterized by severe persistent diarrhea and severe abdominal cramps and may be associated with the passage of blood and mucus. Endoscopic examination may reveal pseudomembranous colitis. Stool culture for *Clostridioides difficile* and stool assay for *C. difficile* toxin may be helpful diagnostically.**

**When significant diarrhea occurs, the drug should be discontinued. Large bowel endoscopy should be considered to establish a definitive diagnosis in cases of severe diarrhea.**

**Antiperistaltic agents such as opiates and diphenoxylate with atropine may prolong and/or worsen the condition. Vancomycin has been found to be effective in the treatment of antibiotic-associated pseudomembranous colitis produced by *Clostridioides difficile*. The usual adult dosage is 500 milligrams to 2 grams of vancomycin orally per day in three to four divided doses administered for 7 to 10 days. Cholestyramine or colestipol resins bind vancomycin *in vitro*. If both a resin and vancomycin are to be administered concurrently, it may be advisable to separate the time of administration of each drug.**

**Diarrhea, colitis, and pseudomembranous colitis have been observed to begin up to several weeks following cessation of oral and parenteral therapy with clindamycin.**

## **PRECAUTIONS**

### **General**

Clindamycin phosphate topical solution contains an alcohol base which will cause burning and irritation of the eye. In the event of accidental contact with sensitive surfaces (eye, abraded skin, mucous membranes), bathe with copious amounts of cool tap water. The solution has an unpleasant taste and caution should be exercised when applying medication around the mouth.

Clindamycin phosphate topical solution should be prescribed with caution in atopic individuals.

### **Drug Interactions**

Clindamycin has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. Therefore it should be used with caution in patients receiving such agents.

### **Pregnancy**

#### **Teratogenic effects**

In clinical trials with pregnant women, the systemic administration of clindamycin during the second and third trimesters has not been associated with an increased frequency of congenital abnormalities. There are no adequate studies in pregnant women during the first trimester of pregnancy. Clindamycin should be used during the first trimester of pregnancy only if clearly needed.

### **Nursing Mothers**

It is not known whether clindamycin is excreted in breast milk following use of clindamycin phosphate topical solution. However, orally and parenterally administered clindamycin has been reported to appear in breast milk. Clindamycin has the potential to cause adverse effects on the breast-fed infant's gastrointestinal flora. Monitor the breast-fed infant for possible adverse effects on the gastrointestinal flora, such as

diarrhea, candidiasis (thrush, diaper rash) or rarely, blood in the stool indicating possible antibiotic-associated colitis.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for clindamycin and any potential adverse effects on the breast-fed child from clindamycin or from the underlying maternal condition.

### **Clinical Considerations**

If used during lactation and clindamycin phosphate topical solution is applied to the chest, care should be taken to avoid accidental ingestion by the infant.

### **Pediatric Use**

Safety and effectiveness in pediatric patients under the age of 12 have not been established.

### **Geriatric Use**

Clinical studies for Clindamycin Phosphate did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

## **ADVERSE REACTIONS**

In 18 clinical studies of various formulations of Clindamycin phosphate using placebo vehicle and/or active comparator drugs as controls, patients experienced a number of treatment emergent adverse dermatologic events [see table below].

<b>Treatment Emergent Adverse Event</b>	<b>Number of Patients Reporting Events</b>		
	<b>Solution n=553(%)</b>	<b>Gel n=148(%)</b>	<b>Lotion n=160(%)</b>
Burning	62 (11)	15 (10)	17 (11)
Itching	36 ( 7)	15 (10)	17 (11)
Burning/Itching	60 (11)	# ( - )	# ( - )
Dryness	105 (19)	34 (23)	29 (18)
Erythema	86 (16)	10 ( 7)	22 (14)
Oiliness/Oily Skin	8 ( 1)	26 (18)	12* (10)
Peeling	61 (11)	# ( - )	11 ( 7)

# not recorded

\* of 126 subjects

Orally and parenterally administered clindamycin has been associated with severe colitis which may end fatally.

Cases of diarrhea, bloody diarrhea and colitis (including pseudomembranous colitis) have been reported as adverse reactions in patients treated with oral and parenteral formulations of clindamycin and rarely with topical clindamycin (see WARNINGS).

Abdominal pain, gastrointestinal disturbances, gram-negative folliculitis, eye pain and contact dermatitis have also been reported in association with the use of topical

formulations of clindamycin.

## **OVERDOSAGE**

Topically applied clindamycin phosphate can be absorbed in sufficient amounts to produce systemic effects (see WARNINGS).

## **DOSAGE AND ADMINISTRATION**

Apply a thin film of Clindamycin phosphate topical solution, twice daily to affected area.

Clindamycin phosphate topical solution USP, 1% is flammable. Avoid fire, flame, and smoking during and immediately following application.

Keep all liquid dosage forms in containers tightly closed.

## **HOW SUPPLIED**

Clindamycin phosphate topical solution USP, 1% containing clindamycin phosphate equivalent to 10 mg clindamycin per milliliter are clear solution with characteristics odor free from particulate matter. Filled in white HDPE bottle fitted with white PP cap and is available in the following sizes:

Clindamycin phosphate topical solution USP, 1% is

NDC 72578-084-02 in applicator bottle of 30 mL

NDC 72578-084-03 in applicator bottle of 60 mL

## **Storage Conditions**

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

Protect from freezing.

Clindamycin phosphate topical solution USP, 1% is flammable. Keep away from heat, sparks or open flames.

**Call your doctor for medical advice about side effects. You may report side effects to Viona Pharmaceuticals Inc. at 1-888-304-5011 or FDA at 1-800-FDA-1088.**

**Manufactured by:**

**Zydus Lifesciences Ltd.**

Changodar, Ahmedabad, India.

**Distributed by:**

**Viona Pharmaceuticals Inc.**

Cranford, NJ 07016

Rev.: 12/25

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 72578-084-02

Clindamycin phosphate topical solution USP, 1%

Rx only

<p>*Each mL contains clindamycin phosphate, USP equivalent to 10mg/mL (1%) of clindamycin. Also contains isopropyl alcohol 50% v/v, propylene glycol and purified water. Sodium hydroxide or hydrochloric acid may be added to adjust pH between 4.0 to 7.0.</p> <p><b>DOSAGE AND USE:</b> See accompanying prescribing information.</p> <p>Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature]. Protect from freezing. Store in an upright fashion.</p> <p>Keep container tightly closed.</p> <p>For external use only. Avoid contact with eyes.</p> <p>Keep this and all drugs out of the reach of children.</p> <p><b>Patient Information:</b></p> <ol style="list-style-type: none"><li>1. Clean and dry the skin area to be treated.</li><li>2. Apply a thin film of medication to the affected area. Use sparingly, avoiding eyes and mouth. If medication accidentally enters eyes, rinse thoroughly with tap water.</li><li>3. If using the applicator top, use dabbing motion of the tip rather than a rolling action. If tip becomes dry, invert the bottle and depress tip several times until it becomes moist.</li></ol>	<p style="text-align: center;">NDC 72578-084-02</p> <h1 style="text-align: center;">Clindamycin Phosphate Topical Solution, USP</h1> <p style="text-align: center;"><b>1 %*</b></p> <p style="text-align: center;">Solution for topical use only</p> <p style="text-align: center;"> <b>30 mL</b> Rx only</p>	<p><b>Manufactured by:</b> <b>Zydus Lifesciences Ltd.</b> Changodar, Ahmedabad, India</p> <p><b>Distributed by:</b> <b>Viona Pharmaceuticals Inc.</b> Cranford, NJ 07016</p> <p>GUJIDRUGS/G/28/1367 XXXXXX Rev.: 08/22</p>	 <p>N 3 7 2 5 7 8 0 8 4 0 2 8</p>
			



# CLINDAMYCIN PHOSPHATE

clindamycin phosphate solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72578-084
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
CLINDAMYCIN PHOSPHATE (UNII: EH6D7113I8) (CLINDAMYCIN - UNII:3U02EL437C)	CLINDAMYCIN PHOSPHATE	10 mg in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72578-084-02	1 in 1 CARTON	06/15/2020	
1		30 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		
2	NDC:72578-084-03	1 in 1 CARTON	06/15/2020	
2		60 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208767	06/15/2020	

**Labeler** - Viona Pharmaceuticals Inc (081468959)**Registrant** - Zydus Lifesciences Limited (650199482)**Establishment**

Name	Address	ID/FEI	Business Operations
Zydus Lifesciences Limited		650650802	ANALYSIS(72578-084) , MANUFACTURE(72578-084)

Revised: 1/2026

Viona Pharmaceuticals Inc