

ASPIRIN- aspirin tablet
Geri-Care Pharmaceuticals, Corp

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Geri Care 44-249

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- minor pain of arthritis
- toothache
- menstrual pain

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
 - vomit blood
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments?

1-800-540-3765

Principal Display Panel

NDC 57896-901-10

GERICARE

ASPIRIN

(NSAID) • 325 mg each

PAIN RELIEVER / FEVER REDUCER

COMPARE TO THE ACTIVE
INGREDIENT OF BAYER® ASPIRIN**

1000 TABLETS

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

**This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Aspirin.

DISTRIBUTED BY:

GERI-CARE PHARMACEUTICAL CORP.

1650 63 Street, Brooklyn, NY 11204

50844 REV0318B24916

NDC 57896-901-10

GERICARE

ASPIRIN
(NSAID) • 325 mg each

PAIN RELIEVER / FEVER REDUCER

COMPARE TO THE ACTIVE
INGREDIENT OF BAYER® ASPIRIN**

1000 TABLETS Actual Size

Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves: ■ pain and fever of colds ■ headache
■ toothache ■ muscle pain ■ minor pain of arthritis ■ menstrual pain

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older
■ take a blood thinning (anticoagulant) or steroid drug
■ have 3 or more alcoholic drinks every day while using this product
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed
■ have had stomach ulcers or bleeding problems

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you have asthma ■ you are taking a diuretic
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ have bloody or black stools ■ feel faint
■ have stomach pain that does not get better ■ vomit blood
■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

Other information ■ use by expiration date on package
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient corn starch

Questions or comments? 1-800-540-3765

50844 REV0318B24916
*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Aspirin.
DISTRIBUTED BY:
GERI-CARE PHARMACEUTICAL CORP.
1650 63 Street, Brooklyn, NY 11204

3 57896 90110 3

GeriCare 44-249

ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-901	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg	
Inactive Ingredients				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	WHITE	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;249;ASPIRIN	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-901-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
2	NDC:57896-901-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
3	NDC:57896-901-27	250 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/1993	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH NOT FINAL	part343	08/04/1993		

Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(57896-901)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(57896-901)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(57896-901)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	MANUFACTURE(57896-901)

Revised: 8/2019

Geri-Care Pharmaceuticals, Corp