COLD FLU SEVERE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated Harris Teeter

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HTE - 1163 - 2019-1011

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 caplets every 4 hours swallow whole – do not crush, chew, or dissolve do not take more than 10 caplets in 24 hours
children under 12 years	■ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Harris Teeter

Cold & Flu Severe

Compare to the Active Ingredients in Tylenol(R) Cold + Flu Severe

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin

PAIN RELIEVER/FEVER REDUCER, COUGH SUPPRESSANT, NASAL DECONGESTANT, EXPECTORANT

Relieves

Head & Body Aches

Fever & Sore Throat

Cough

Nasal Congestion

Mucus & Chest Congestion

For Adults

24 COOL TASTE CAPLETS

Actual Size



- trouble uninating due to an enlarged prostate gland
 - heart disease
 high blood pressure
- thyroid disease ■ liver disease Ask a doctor before use if you have
- if you have ever had an allergic reaction to this product or any of its ingredients an MAOI, ask a doctor or pharmacist before taking this product. the MAOI drug. If you do not know if your prescription drug contains conditions, or Parkinson's disease), or for 2 weeks after stopping (MAOI) (certain drugs for depression, psychiatric, or emotional if you are now taking a prescription monoamine oxidase inhibitor
 - acetaminophen, ask a doctor or pharmacist. nonprescription). If you are not sure whether a drug contains ■ with any other drug containing acetaminophen (prescription or

vomiting, consult a doctor promptly.

days, is accompanied or followed by fever, headache, rash, nausea, or Sore throat warning: If sore throat is severe, persists for more than 2 If a skin reaction occurs, stop use and seek medical help right away. Symptoms may include: a skin reddening a blisters a rash Allergy alert: acetaminophen may cause severe skin reactions.

- 3 or more alcoholic drinks every day while using this product ■ with other drugs containing acetaminophen
 - hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
- daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 Liver warning: This product contains acetaminophen. The maximum Warnings

Drug Facts (continued)

12 years ask a doctor children under ■ do not take more than 10 caplets in 24 hours years and over swallow whole – do not crush, chew, or dissolve Children 12 ■ take 2 caplets every 4 hours adults and

> do not take more than directed (see overdose warning) Directions

attention is critical for adults as well as for children even if you do not Poison Control Center right away (1-800-222-1222). Quick medical Overdose warning: In case of overdose, get medical help or contact a

Keep out of reach of children. If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.

- cough comes back or occurs with rash or headache that lasts ■ redness or swelling is present uew symptoms occur ■ fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if

When using this product do not exceed recommended dosage thinning drug warfarin

Ask a doctor or pharmacist before use if you are taking the blood condy that occurs with too much phiegm (mucus)

chronic bronchitis, or emphysema persistent or chronic cough such as occurs with smoking, asthma,

Drug Facts (continued)

condus more productive ■ condµ ■ usasi congestion

Nasal decongestant увелујерћипе НСІ 5 тд Expectorant Guaifenesin 200 mg. Cough suppressant Dextromethorphan HBr 10 mg. Active ingredients (in each caplet) Purpose

Drug Facts

■ helps loosen phlegm (mucus) and thin bronchial secretions to make ■ www. scues gug bgius ■ SOUG UNIOSI иезазсие USES • for the temporary relief of the following cold/flu symptoms:

Drug Facts (continued)

†Compare to the Active Ingredients in Tylenol® Cold +

Cold & Flu Severe

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCI, Guaifenesin

PAIN RELIEVER / FEVER REDUCER, COUGH SUPPRESSANT NASAL DECONGESTANT, EXPECTORANT

Relieves

- Head & Body Aches
- Fever & Sore Throat
- Cough
- Nasal Congestion
- Mucus & Chest Congestion
- For Adults

24 COOL TASTE CAPLETS

Actual Size

silicon dioxide, croscarmellose sodium, crospovidone, D&C yellow #10 ▶ acesulfame potassium, colloidal Inactive ingredients

> retain carton for complete product information Other information a store between 20-25°C (68-77°F) in a dry place

> > **Drug Facts** (continued)

1-844-705-4384

povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol,

Drug Facts (continued)

Questions or comments?

aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate,

COLD FLU SEVERE

Syourplusguarantee.

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol[®] Cold + Flu Severe.

PROUDLY DISTRIBUTED BY: HARRIS TEETER, LLC MATTHEWS, NC 28105

USE IF BLISTER UNITS TORN OR BROKEN

NOT L 8





acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72036-155
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6 MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)	
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	

Product Characteristics			
Color	yello w	Score	no score
Shape	OVAL	Size	19 mm
Flavor	MINT	Imprint Code	AAA;1136
Contains			

Pack	aging			
#]	Item Code	Package Description	Marketing Start Date	Marketing End Date

1 NDC:72036-155-05	2 in 1 CARTON	08/16/2016			
1	12 in 1 BLISTER PACK; Type 0: Not a Combination Produc	t			
Maybating Information					
Marketing Information					
Marketing Categor	y Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	08/16/2016			

Labeler - Harris Teeter (047279351)

Revised: 10/2019 Harris Teeter