

GAINPRO 10- bambermycins powder
Huvepharma, Inc.

GAINPRO® 10
(bambermycins Type A Medicated Article)

WITHDRAWAL PERIODS

No withdrawal period is required when used according to labeling.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

INDICATIONS FOR USE:

For increased rate of weight gain and improved feed efficiency in growing beef steers and heifers fed in confinement for slaughter. For increased rate of weight gain in growing beef steers and heifers on pasture (stocker, feeder, and slaughter), and replacement beef and dairy heifers on pasture.

ACTIVE INGREDIENT:

Bambermycins 10 grams per pound (22 grams per kilogram)

INGREDIENTS:

Roughage products and/or calcium carbonate, dried bambermycins fermentation product, ethoxyquin (a preservative), mineral oil.

MIXING DIRECTIONS AND INDICATIONS FOR USE ON REVERSE SIDE.

FOR USE IN MANUFACTURED FEEDS ONLY.

USER SAFETY WARNING:

Not for human use. Keep out of reach of children.

Restricted Drug (California) - use only as directed

STORAGE:

Store at or below 25°C (77°F), excursions permitted to 40°C (104°F).

Approved by FDA under NADA # 141-034

Net Weight - 50 pounds (22.68 kg)

HUVEPHARMA®

Distributed by:

HUVEPHARMA, INC.

Peachtree City, GA 30269

INDICATIONS FOR USE

Cattle Classes	Bambermycins g/ton in feed	Bambermycins mg/hd/d	Indications
Growing Beef Steers and Heifers Fed in Confinement for Slaughter	1 - 4	10 to 20 mg/hd/day**	Increased rate of weight gain, improved feed efficiency
Growing Beef Steers and Heifers on Pasture (Stocker, Feeder, and Slaughter), and Replacement Beef and Dairy Heifers on Pasture	2 - 80	10 to 40 mg/hd/day***	Increased rate of weight gain

*Only a single level of bambermycins (e.g. 2 grams bambermycins/ton) can be indicated on a feed tag. A different feed tag must be submitted for each level of bambermycins needed for your complete feeding program.

GROWING BEEF STEERS AND HEIFERS FED IN CONFINEMENT FOR SLAUGHTER:

****THE APPROVED DOSE IN GROWING BEEF STEERS AND HEIFERS FED IN CONFINEMENT**

FOR SLAUGHTER IS 10 TO 20 MG OF BAMBERMYCINS/HD/DAY. The amount of bambermycins

will range between 1 and 4 grams per ton of feed depending upon the daily feed consumption

of the cattle. The amount of Type A Medicated Article added to the complete ration (Type C

Medicated Feed) must be based on daily feed consumption, the number of animals in the pen

and the level (10-20 mg/hd/day) of bambermycins being fed. From this information the grams

of bambermycins per ton of feed can be calculated in order to fulfill feed assay requirements.

See the following table for conversions from mg/hd/day to g/t.

EXAMPLE OF CONVERTING THE BAMBERMYCINS CATTLE DOSAGE FROM MG/HEAD/DAY TO GRAMS/TON*

Cattle Weight		Average Feed Consumption/ Head/Day	Grams Bambermycins/ Ton of Feed		Mg Bambermycins Fed Daily to each Animal	
Lbs.	Kg	Lbs. ¹	Low	High	Low	High
400	181.4	10.0	2.00	4.00	10.0	20.0
500	226.8	14.0	1.50	2.80	10.5	19.6
600	272.2	17.0	1.25	2.30	10.6	19.6
700	317.5	20.0	1.00	2.00	10.0	20.0
800	362.9	22.0	1.00	1.80	11.0	19.8
900	408.2	23.0	1.00	1.70	11.5	19.6
1000	453.6	24.0	1.00	1.60	12.0	19.2
1100	499.0	25.0	1.00	1.60	12.5	20.0
1200	544.3	26.0	1.00	1.50	13.0	19.5
1300	589.7	27.0	1.00	1.40	13.5	18.9

*Dosage range is 10 to 20 mg/head/day

¹Assumes as fed ad libitum consumption and average daily gain of 2.0 to 3.0 lbs./day

Example Calculation—Amount of GAINPRO®-10 added to feed for a pen of cattle. Given: Average Weight—800 lbs.; Number in pen—100 hd.; Dosage—20 mg/hd/day; Daily

Feed Consumption—22 lbs./day.

- 100 hd x 20 mg = 2,000 mg or 2 grams bambermycins.
- GAINPRO®-10 contains 10 grams bambermycins/lb., therefore add 0.2 pound of GAINPRO®-10 to the daily feed for this pen of 100 animals.
- 22 lbs./day x 100 hd = 2,200 lbs. total feed consumption.
- 2 grams bambermycins ÷ 2,200 lbs. = .000909 g/lb.
- .000909 x 2,000 lbs. = 1.82 g bambermycins/ton of feed.

MIXING DIRECTIONS:

It is recommended that GAINPRO®-10 be diluted with a feed ingredient before addition to the final feed. A dilution of 1 part of GAINPRO®-10 and 9 parts feed ingredient is the suggested working intermediate pre-blend. The table below shows examples of GAINPRO®-10 and working intermediate pre-blend addition levels for various commonly used feeds.

Feed for Growing Beef Steers and Heifers Fed in Confinement for Slaughter			
Bambermycins in Feed g/ton	Making 1 to 9 Intermediate Pre-blend		1 to 9 Intermediate Pre-blend lb to be added per ton
	GAINPRO®-10 lb	Feed Ingredient lb	
1	0.1	0.9	1
2	0.2	1.8	2
3	0.3	2.7	3
4	0.4	3.6	4

Thoroughly mix both working intermediate pre-blend and finished feed to ensure complete and uniform distribution of the GAINPRO®-10.

FEEDING DIRECTIONS:

Feed continuously in a complete feed at the rate of 10-20 mg bambermycins/head/day from the Type C Medicated Feed containing 1-4 g bambermycins/ton.

TYPE B LIQUID MEDICATED FEED INTENDED FOR ADDITION TO DRY FEEDS FOR

GROWING BEEF STEERS AND HEIFERS FED IN CONFINEMENT FOR SLAUGHTER:

• Type B Liquid Medicated Feed should be in a pH range of 3.8 to 7.5 and should have a moisture content between 30 and 45%.

• Type B Liquid Medicated Feed must be mixed into the dry feed within 8 weeks of manufacture.

• Type B Liquid Medicated Feed can be manufactured containing 40.0 to 800.0 g of bambermycins/

ton. Mixing directions into dry feed are as follows:

Before feeding, the Type B Liquid Medicated Feed must be thoroughly mixed with other feed

materials to make a complete feed (Type C Medicated Feed). Examples of addition rates to

complete feeds are shown in the following table.

Bambermycins (g/ton) in Type B Liquid	Addition Rate of Type B in Liquid Medicated Feed (lb/ton) to Complete	Bambermycins (g/ton) Final Complete Feed (Type C Medicated
---------------------------------------	---	--

Medicated Feed	Feed	Feed)
40.0	50.0	1.0
40.0	200.0	4.0
100.0	20.0	1.0
100.0	80.0	4.0
400.0	5.0	1.0
400.0	20.0	4.0
800.0	2.5	1.0
800.0	10.0	4.0

The complete dry feed (Type C Medicated Feed) may only be fed to growing beef steers and heifers fed in confinement for slaughter and must be fed within one week after mixing.

GROWING BEEF STEERS AND HEIFERS ON PASTURE (STOCKER, FEEDER, AND SLAUGHTER),

AND REPLACEMENT BEEF AND DAIRY HEIFERS ON PASTURE:

***THE APPROVED DOSE IN GROWING BEEF STEERS AND HEIFERS ON PASTURE (STOCKER, FEEDER, AND SLAUGHTER), AND REPLACEMENT BEEF AND DAIRY HEIFERS ON PASTURE IS

10 TO 40 MG OF BAMBERMYCINS/HD/DAY. Daily intakes of bambermycins in excess of 20 mg/head/day have not been shown to be more effective than 20 mg/head/day.

MIXING DIRECTIONS:

It is recommended that GAINPRO®-10 be diluted with a feed ingredient before addition to the supplemental feed. A dilution of 1 part of GAINPRO®-10 and 9 parts feed ingredient is the suggested working intermediate pre-blend. The table below shows examples of GAINPRO®-10 and working intermediate pre-blend addition levels and feeding rates.

Feed for Growing Beef Steers and Heifers on Pasture (Stocker, Feeder, and Slaughter) and Replacement Beef and Dairy Heifers on Pasture				
Bambermycins in Feed g/ton	Making 1 to 9 Intermediate Pre-blend		1 to 9 Intermediate Pre-blend lb to be added per ton	Feeding Rate lb/hd/day
	GAINPRO®-10 lb	Feed Ingredient lb		
2	0.2	1.8	2	10
4	0.4	3.6	4	5 - 10
8	0.8	7.2	8	2.5 - 10
20	2.0	18	20	1 - 4
40	4.0	36	40	1 - 2
80	8.0	72	80	1

Thoroughly mix both working intermediate pre-blend and finished feed to ensure complete and uniform distribution of the GAINPRO®-10.

To manufacture Type B medicated feeds containing 100.0 to 800.0 g/ton bambermycins mix 10 to 80 lbs. of GAINPRO®-10 with 1990 to 1920 lbs. of non-medicated feed.

To manufacture Type C medicated feeds containing 2.0 to 80.0 g/ton bambermycins from

Type B medicated feeds mix:

40 to 1600 lbs. of Type B feed containing 100 g/ton bambermycins with 1960 to 400 lbs.

of non-medicated feed, or

5 to 200 lbs. of Type B feed containing 800 g/ton bambermycins with 1995 to 1800 lbs. of non-medicated feed.

Thoroughly mix both Type B medicated feed and finished feed to ensure complete and uniform distribution of the GAINPRO®-10.

FEEDING DIRECTIONS FOR HAND-FED TYPE C MEDICATED FEED:

Feed continuously on a hand-fed basis at a rate of 10 to 40 mg bambermycins per head per day. The drug must be contained in 1 to 10 lbs. of supplemental Type C medicated feed containing 2 to 80 g bambermycins per ton. Daily intakes of bambermycins in excess of 20 mg/head/day have not been shown to be more effective than 20 mg/head/day.

TYPE C FREE-CHOICE MEDICATED FEED:

An FDA approved formula must be used to manufacture free-choice medicated feeds containing bambermycins. Feed mills that use a proprietary formula are required to be licensed.

HUVEPHARMA®

Rev. 02-2022

1114113001

GAINPRO[®] 10

(bambermycins Type A medicated article)

GAINPRO[®] 10

(bambermycins Type A medicated article)

WITHDRAWAL PERIODS

No withdrawal period is required when used according to labeling.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

INDICATIONS FOR USE:

For increased rate of weight gain and improved feed efficiency in growing beef steers and heifers fed in confinement for slaughter. For increased rate of weight gain in growing beef steers and heifers on pasture (stocker, feeder, and slaughter), and replacement beef and dairy heifers on pasture.

ACTIVE INGREDIENT:

Bambermycins 10 grams per pound (22 grams per kilogram)

INGREDIENTS:

Roughage products and/or calcium carbonate, dried bambermycins fermentation product, ethoxyquin (a preservative), mineral oil.

MIXING DIRECTIONS AND INDICATIONS FOR USE ON REVERSE SIDE.

FOR USE IN MANUFACTURED FEEDS ONLY.

USER SAFETY WARNING:

Not for human use. Keep out of reach of children.

Restricted Drug (California) – use only as directed

STORAGE:

Store at or below 25°C (77°F), excursions permitted to 40°C (104°F).

Approved by FDA under NADA # 141-034

Net Weight – 50 pounds (22.68 kg)



Distributed by:
HUVEPHARMA, INC.
Peachtree City, GA 30269

GAINPRO[®] 10

(bambermycins Type A medicated article)

INDICATIONS FOR USE

Cattle Classes	Bambermycins g/ton in feed	Bambermycins mg/hd/d	Indications
Growing Beef Steers and Heifers Fed in Confinement for Slaughter	1 - 4	10 to 20 mg/hd/day**	Increased rate of weight gain, improved feed efficiency
Growing Beef Steers and Heifers on Pasture (Stocker, Feeder, and Slaughter), and Replacement Beef and Dairy Heifers on Pasture	2 - 80	10 to 40 mg/hd/day***	Increased rate of weight gain

*Only a single level of bambermycins (e.g. 2 grams bambermycins/ton) can be indicated on a feed tag. A different feed tag must be submitted for each level of bambermycins needed for your complete feeding program.

GAINPRO[®] 10

(bambermycins Type A medicated article)

GROWING BEEF STEERS AND HEIFERS FED IN CONFINEMENT FOR SLAUGHTER:

**THE APPROVED DOSE IN GROWING BEEF STEERS AND HEIFERS FED IN CONFINEMENT FOR SLAUGHTER IS 10 TO 20 MG OF BAMBERMYCINS/HD/DAY. The amount of bambermycins will range between 1 and 4 grams per ton of feed depending upon the daily feed consumption of the cattle. The amount of Type B Medicated Article is added to the complete ration (Type C Medicated Feed) must be based on daily feed consumption, the number of animals in the pen and the level (10-20 mg/hd/day) of bambermycins being fed. From this information the grams of bambermycins per ton of feed can be calculated in order to fulfill feed assay requirements. See the following table for conversions from mg/hd/day to g/t.

EXAMPLE OF CONVERTING THE BAMBERMYCINS CATTLE DOSAGE FROM MG/HEAD/DAY TO GRAMS/TON*

Cattle Weight	Average Feed Consumption/Head/Day	Grams Bambermycins/Ton of Feed		Mg Bambermycins Fed Daily to each Animal		
		Low	High	Low	High	
400	181.4	10.0	2.00	4.00	10.0	20.0
500	226.8	14.0	1.50	2.93	10.5	19.6
600	272.2	17.0	1.25	2.30	10.6	19.6
700	317.5	20.0	1.00	2.00	10.0	20.0
800	362.9	22.0	1.00	1.80	11.0	19.8
900	408.2	23.0	1.00	1.70	11.5	19.6
1000	453.6	24.0	1.00	1.60	12.0	19.2
1100	499.0	25.0	1.00	1.60	12.5	20.0
1200	544.3	26.0	1.00	1.50	13.0	19.5
1300	589.7	27.0	1.00	1.40	13.5	18.9

*Dosage range is 10 to 20 mg/head/day.
 **Assumes an feed intake consumption and average daily gain of 2.0 to 3.0 lbs/day.
 Example Calculation—Amount of GAINPRO-10 added to feed for a pen of cattle:
 Given: Average Weight—800 lbs., Number in pen—100 hd., Dosage—20 mg/head/day, Daily Feed Consumption—22 lbs./day.
 • 100 hd x 20 mg = 2,000 mg or 2 grams bambermycins.
 • GAINPRO-10 contains 10 grams bambermycins/lb., therefore add 0.2 pound of GAINPRO-10 to the daily feed for this pen of 100 animals.
 • 22 lbs./day x 100 hd = 2,200 lbs. total feed consumption.
 • 2 grams bambermycins ÷ 2,200 lbs. = .000909 g/lb.
 • .000909 x 2,000 lbs. = 1.82 g bambermycins/ton of feed.

MIXING DIRECTIONS:

It is recommended that GAINPRO-10 be diluted with a feed ingredient before addition to the final feed. A dilution of 1 part of GAINPRO-10 and 9 parts feed ingredient is the suggested working intermediate pre-blend. The table below shows examples of GAINPRO-10 and working intermediate pre-blend addition levels for various commonly used feeds:

Bambermycins in Feed g/ton	Making 1 to 9 Intermediate Pre-blend GAINPRO-10	Feed Ingredient	1 to 9 Intermediate Pre-blend lb to be added per ton
1	0.1	0.9	2
2	0.2	1.8	3
3	0.3	2.7	4
4	0.4	3.6	4

Thoroughly mix both working intermediate pre-blend and finished feed to ensure complete and uniform distribution of the GAINPRO-10.

FEEDING DIRECTIONS:

Feed continuously in a complete feed at the rate of 10-20 mg bambermycins/head/day from the Type C Medicated Feed containing 1-4 g bambermycins/ton.

TYPE B LIQUID MEDICATED FEED INTENDED FOR ADDITION TO DRY FEEDS FOR GROWING BEEF STEERS AND HEIFERS FED IN CONFINEMENT FOR SLAUGHTER:

*Type B Liquid Medicated Feed should be in a pH range of 3.8 to 7.5 and should have a moisture content between 30 and 45%.
 • Type B Liquid Medicated Feed must be mixed into the dry feed within 8 weeks of manufacture.
 • Type B Liquid Medicated Feed can be manufactured containing 40.0 to 800.0 g of bambermycins/ton. Mixing directions into dry feed are as follows:

Before feeding, the Type B Liquid Medicated Feed must be thoroughly mixed with other feed materials to make a complete feed (Type C Medicated Feed). Examples of addition rates to complete feeds are shown in the following table.

Bambermycins (g/ton) Type B Liquid Medicated Feed	Addition Rate of Type B Liquid Medicated Feed (lb/ton) to Complete Feed	Bambermycins (g/ton) in Final Complete Feed (Type C Medicated Feed)
40.0	50.0	1.0
40.0	200.0	4.0
100.0	20.0	1.0
100.0	80.0	4.0
400.0	5.0	1.0
400.0	20.0	4.0
800.0	2.5	1.0
800.0	10.0	4.0

The complete dry feed (Type C Medicated Feed) may only be fed to growing beef steers and heifers fed in confinement for slaughter and must be fed within one week after mixing.

GROWING BEEF STEERS AND HEIFERS ON PASTURE (STOCKER, FEEDER, AND SLAUGHTER), AND REPLACEMENT BEEF AND DAIRY HEIFERS ON PASTURE:

***THE APPROVED DOSE IN GROWING BEEF STEERS AND HEIFERS ON PASTURE (STOCKER, FEEDER, AND SLAUGHTER), AND REPLACEMENT BEEF AND DAIRY HEIFERS ON PASTURE IS 10 TO 40 MG OF BAMBERMYCINS/HD/DAY. Daily intakes of bambermycins in excess of 20 mg/head/day have not been shown to be more effective than 20 mg/head/day.

MIXING DIRECTIONS:

It is recommended that GAINPRO-10 be diluted with a feed ingredient before addition to the supplemental feed. A dilution of 1 part of GAINPRO-10 and 9 parts feed ingredient is the suggested working intermediate pre-blend. The table below shows examples of GAINPRO-10 and working intermediate pre-blend addition levels and feeding rates.

Bambermycins in Feed g/ton	Making 1 to 9 Intermediate Pre-blend GAINPRO-10	Feed Ingredient	1 to 9 Intermediate Pre-blend lb to be added per ton	Feeding Rate lb/head/day
2	0.2	1.8	2	10
4	0.4	3.6	4	5 - 10
20	2.0	18.0	20	2.5 - 10
40	4.0	36.0	40	1 - 4
80	8.0	72.0	80	1 - 2

Thoroughly mix both working intermediate pre-blend and finished feed to ensure complete and uniform distribution of the GAINPRO-10.

To manufacture Type B medicated feeds containing 100.0 to 800.0 g/ton bambermycins mix 10 to 80 lbs. of GAINPRO-10 with 1990 to 1920 lbs. of non-medicated feed.

To manufacture Type C medicated feeds containing 2.0 to 80.0 g/ton bambermycins from Type B medicated feeds mix:

40 to 1600 lbs. of Type B feed containing 100 g/ton bambermycins with 1960 to 400 lbs. of non-medicated feed, or

5 to 200 lbs. of Type B feed containing 800 g/ton bambermycins with 1995 to 1900 lbs. of non-medicated feed.

Thoroughly mix both Type B medicated feed and finished feed to ensure complete and uniform distribution of the GAINPRO-10.

FEEDING DIRECTIONS FOR HAND-FED TYPE C MEDICATED FEED:

Feed continuously on a hand-fed basis at a rate of 10 to 40 mg bambermycins per head per day. The drug must be contained in 1 to 10 lbs. of supplemental Type C medicated feed containing 2 to 80 g bambermycins per ton. Daily intakes of bambermycins in excess of 20 mg/head/day have not been shown to be more effective than 20 mg/head/day.

TYPE C FREE-CHOICE MEDICATED FEED:

An FDA approved formula must be used to manufacture free-choice medicated feeds containing bambermycins. Feed mills that use a proprietary formula are required to be licensed.



1114112901
Rev. 02-2022

bambermycins powder

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:23243-2070
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BAMBERMYCINS (UNII: PP922A42V2) (BAMBERMYCINS - UNII:PP922A42V2)	BAMBERMYCINS	22 g in 1 kg

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
ETHOXYQUIN (UNII: 9T1410R4OR)	

Product Characteristics

Color	brown ((Off White to Tan))	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-2070-5	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141034	08/22/2016	

Labeler - Huvepharma, Inc. (619153559)

Registrant - Huvepharma EOOD (552691651)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc.		883128204	manufacture, analysis, label, pack, medicated animal feed manufacture

