

**ACETAMINOPHEN 500 MG- acetaminophen tablet**  
**NUVICARE LLC**

-----  
**Extra Strength Pain Relief ACETAMINOPHEN 500 mg**

***Drug Facts***

***Active ingredient (in each caplet)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- If you are allergic to acetaminophen or any of the inactive ingredient in this product.

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor/pharmacist before use if you are**

- taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**KEEP OUT OF REACH OF CHILDREN.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

**Adults and children 12 years and over:**

- take 2 Caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

**Children under 12 Years**

- ask a doctor

**Other information**

- store at 20° - 25°C (68° -77°F)

**Inactive ingredients**

Pregelatinized starch, Polyvinyl pyrrolidone, stearic acid powder, microcrystalline cellulose, magnesium stearate, Hypromellose-5, Hypromellose-15, polyethylene glycol 400, titanium dioxide, talc

**Questions or comments?**

Call 1 (718) 337-8733 or visit: [support@nuvicare.com](mailto:support@nuvicare.com)

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc.,



**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84324-021-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2025	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2025	

**Labeler** - NUVICARE LLC (119257565)

Revised: 12/2025

NUVICARE LLC