# MUCUS RELIEF COUGH AND CONGESTION DM- guaifenes in and dextromethorphan hbr tablet

## Richmond Pharmaceuticals, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# Guaifenes in 400 mg - DextromethorphanCaplets 20 mg Caplets

#### Active ingredients (in each caplet)

Dextromethorphan HBr 20 mg Guaifenesin 400 mg

# Purpose

Cough suppressant Expectorant

#### Uses

- temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

#### **Warnings**

#### Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- cough accompanied by too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

#### When using this product

do not use more than directed.

#### Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

#### If pregnant or breast-feeding,

ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- take with a full glass of water
- **adults and children 12 years of age and over:** 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.
- children under 12 years: do not use

# Other information

■ store at 15°-30°C (59°-86°F)

# **Inactive ingredients**

Colloidal silicon dioxide, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, stearic acid

# **Questions or comments?**

call 804-270-4498, 8.30 am-4.30 pm ET, Monday - Friday TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

# **Principal Display Panel**

 $Mucus\ Relief\ Cough\ and\ Congestion\ DM$ 

NDC- 54738-985-30....30 caplets

LR0116 NDC 54738-985-30 Drug Facts (continued on back of label depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before Purposes Cough Suppressant
..... Expectorant TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING monoamine oxidase inhibitor (MAOI) (certain drugs for temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or helps loosen philegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more GUAIFENESIN Distributed By: Richmond Pharmaceuticals, Inc. Do not use if you are now taking a prescription 400 mg Richmond, VA 23233 USA & Ask a doctor before use if you have **DEXTROMETHORPHAN HBr** Dextromethorphan HBr 20 mg NO UV AREA 20 mg Active ingredients **Expectorant &** in each caplet) Guaifenesin 400 mg **Drug Facts** using this product **Cough Suppressant** emphysema Warnings productive **30 CAPLETS** Jses Richmond Pharmaceuticals, Inc.

TOP LAYER

■ cough accompanied by too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right avay.

Directions

■ do not take more than 6 doses in any 24-hour period

■ this product is not intended for use in children

period

this product is not intended for use in children under 12 years of age

adults & children 12 years and over: 1 caplet every 4 hours

children under 12 years; do not use

■ store at 15°-30°C (59°-86°F)

Inactive ingredients colloida silicon dioxide, magnesium stearate, matodextrin, microcrystalline cellulose, povidone, stearic acit

Questions or comments? call 804-270-4498, 8:30 am – 4:30 pm ET, Mondav – Fridav


NDC- 54738-985-60....60 caplets

GUAIFENESIN
400 mg
&
DEXTROMETHORPHAN HBr
20 mg
Expectorant &
Cough Suppressant

Richmond Pharmaceuticals, Inc.

LR0116 MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the Cough Suppressant monoamine oxidase inhibitor (MAOI) (certain drugs for temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or helps loosen phlegm (mucus) and thin bronchia secretions to rid the bronchial passageways of Do not use if you are now taking a prescription bothersome mucus and make coughs more Distributed By: Richmond Pharmaceuticals, Richmond, VA 23233 USA Ask a doctor before use if you have Sextramethorphan HBr 20 mg NO UV AREA Active ingredients (in each caplet) Drug Facts using this product. nses

## TOP LAYER



#### MUCUS RELIEF COUGH AND CONGESTION DM

guaifenesin and dextromethorphan hbr tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54738-985	
Route of Administration	ORAL			

l	Active Ingredient/Active Moiety		
ı	Ingredient Name	Basis of Strength	Strength

,	DEXTROMETHORPHAN HYDROBROMIDE	20 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg

Inactive Ingredients				
Ingredient Name	Strength			
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MALTO DEXTRIN (UNII: 7CVR7L4A2D)				
PO VIDO NE (UNII: FZ989 GH9 4E)				
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)				

Product Characteristics				
Color	white	Score	2 pieces	
Shape	OVAL	Size	17mm	
Flavor		Imprint Code	AP;150	
Contains				

P	Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
1	NDC:54738-985-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2016		
2	NDC:54738-985-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2016		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	04/01/2016		

# Labeler - Richmond Pharmaceuticals, Inc (043569607)

# Registrant - Advance Pharmaceutical Inc. (078301063)

Establishment				
Name	Address	ID/FEI	<b>Business Operations</b>	
Advance Pharmaceutical Inc.		078301063	manufacture(54738-985)	

Revised: 10/2017 Richmond Pharmaceuticals, Inc