

REGULAR STRENGTH ANTACID MINT - aluminum hydroxide, magnesium hydroxide, simethicone liquid
Topco Associates, LLC

Topco Regular Strength Antacid Mint

ACTIVE INGREDIENTS (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 400 mg
Magnesium hydroxide 400 mg
Simethicone 40 mg

PURPOSE

Antacid
Antacid
Antigas

USE(S)

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

WARNINGS

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- kidney disease
- a magnesium-restricted diet

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR

if symptoms last more than two weeks

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS

- **shake well before use**
- adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor
- children under 12 years: consult a doctor
- mL = milliliter

OTHER INFORMATION

- **each 10 mL contains:** magnesium 170 mg, sodium 5 mg
- store at 20°C-25°C (68°-77°F)
- do not freeze

INACTIVE INGREDIENTS

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

PRINCIPAL DISPLAY PANEL

Topcare Health
NDC 36800-582-40

REGULAR STRENGTH

Antacid

Advanced

ANTACID & ANTIGAS

FAST RELIEF

- Heartburn
- Acid Indigestion
- Pressure
- Bloating (Gas)

ALCOHOL 0.4% v/v

12 FL OZ (355 mL)

MINT FLAVOR

TopCare
health®

NDC 36800-582-40

REGULAR STRENGTH

Antacid

Advanced

ANTACID & ANTIGAS



FAST RELIEF

- Heartburn
- Acid Indigestion
- Pressure
- Bloating (Gas)



Scan here for more information or call 1-888-423-0139

ALCOHOL 0.4% v/v

12 FL OZ
(355 mL)



MINT
FLAVOR

Drug Facts

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each 10 mL) Purpose
 Aluminum hydroxide (equiv. to dried gel, USP) 400 mg....Antacid
 Magnesium hydroxide 400 mg.....Antacid
 Simethicone 40 mg.....Antigas

Uses relieves: ■ acid indigestion ■ heartburn ■ sour stomach ■ upset stomach and gas associated with these symptoms

Warnings

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than two weeks

Keep out of reach of children.

Directions ■ shake well before use ■ adults and children 12 years and older: take 10 - 20 mL four times a day, or as directed by a doctor ■ children under 12 years: consult a doctor ■ mL = milliliter

Other information ■ each 10 mL contains: magnesium 170 mg, sodium 5 mg ■ store at 20°C-25°C (68°F-77°F) ■ do not freeze

Inactive ingredients ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol.

DISTRIBUTED BY TOPCO ASSOCIATES LLC
 ELK GROVE VILLAGE, IL 60007
 ©TOPCO GUAA1020
 QUESTIONS? 1-888-423-0139
 topcare@topco.com www.topcarebrand.com



MFR# 53041 REV 1020



0 36800 92213 6

REGULAR STRENGTH ANTACID MINT

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-582
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-582-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	10/01/2020	

Labeler - Topco Associates, LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(36800-582)