# XERAC AC- xerac ac liquid Person and Covey

-----

Xerac AC

# **OTC - PURPOSE SECTION**

For topical application as an antiperspirant (anhidrotic).

# **DOSAGE & ADMINISTRATION SECTION**

Xerac AC when applied to the underarms, should be applied to absolutely dry skin. Only apply Xerac AC before bedtime, since medication should be kept on the skin for 6-8 hours. To reduce irritation, let the alcohol evaporate, leaving a thin film of antiperspirant on the skin. Wear a T-shirt, while sleeping to prevent medication from being rubbed off on the bed linens. Do not apply Xerac AC to broken, irritated or recently shaved skin. Wash the treated area next morning with soap and water. Do not apply other deodorants or antiperspirants while using Xerac AC.

# WARNINGS SECTION

For external use only. Avoid contact with eyes. If contact occurs, wash eyes thoroughly with water. If irritation or sensitization occurs, discontinue use or contact a physician. Xerac AC may be harmful to cotton fibers and certain metals.

# OTC - KEEP OUT OF REACH OF CHILDREN SECTION

Yes. If swallowed, get medical help or contact a Poison Control Center right away.

# **OTC - ACTIVE INGREDIENT SECTION**

Aluminum Chloride Hexahydrate

#### **INACTIVE INGREDIENT SECTION**

Anhydrous ethanol

# **Indicatins and Usage Section**

For patients suffering from hyperhidrosis, or patients whose jobs or sports activities cause excessive sweating that exacerbates other skin problems.

# **Principal Display Panel**

**DIRECTIONS:** Apply to affected areas at bedtime or as directed by physician.

warning: Keep this and all medication out of reach of children. For external use only. Avoid contact with eyes. This product may be harmful to certain metals and fabrics. Keep container tightly closed. See package insert for full prescribing information.



# xerac™ AC

ALUMINUM CHLORIDE HEXAHYDRATE
IN ANHYDROUS ETHANOL

35 cc

A SOLUTION OF:

Aluminum Chloride (Hexahydrate) 6.25% (w/v) in Anhydrous Ethyl Alcohol (S.D. Alcohol 40) 96% (v/v)

**CAUTION:** Federal Law Prohibits Dispensing Without Prescription.



**DIRECTIONS:** Apply to affected areas at bedtime or as directed by physician.

WARNING: Keep this and all medication out of reach of children. For external use only. Avoid contact with eyes. This product may be harmful to certain metals and fabrics. Keep container tightly closed. See package insert for full prescribing information.



xerac® AC

NDC 0096-0709-60

ALUMINUM CHLORIDE HEXAHYDRATE IN ANHYDROUS ETHANOL

60 cc

A SOLUTION OF:

Aluminum Chloride (Hexahydrate) 6.25% (w/v) in Anhydrous Ethyl Alcohol (S.D. Alcohol 40) 96% (v/v)

CAUTION: Federal Law

Prohibits Dispensing Without Prescription.



# **XERAC AC**

xerac ac liquid

D		Informa	<b>_:</b>
Proc	IIICT I	Intorma	TION
1 1 0 0	исс		LIVII

Product Type HUMAN OTC DRUG Item Code (Source)

NDC:0096-0709

Route of Administration

**TOPICAL** 

# **Active Ingredient/Active Moiety**

Ingredient Name

Basis of Strength

Strength

ALUMINUM CHLORIDE (UNII: 3CYT62D3GA) (ALUMINUM CATION -

UNII:3XHB1D032B)

ALUMINUM CHLORIDE

0.0625 g in 1 g

# **Inactive Ingredients**

**Ingredient Name** 

Strength

ALCOHOL (UNII: 3K9958V90M)

# **Packaging**

#	Item Code	Package Description	Marketing Start  Date	Marketing End Date
	NDC:0096- 0709-60	49.86 g in 1 PACKAGE; Type 0: Not a Combination Product	06/01/1996	
	NDC:0096- 0709-35	29.09 g in 1 PACKAGE; Type 0: Not a Combination Product	06/01/1996	

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M019	06/01/1996			

# Labeler - Person and Covey (008482473)

Establishment					
Name	Address	ID/FEI	Business Operations		
Person and Covey		008482473	manufacture(0096-0709)		

Revised: 1/2024 Person and Covey