TANDEM PLUS BARRIER- iodine solution IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

TANDEM+ BARRIER NPE FREE SANITIZING TEAT DIP A HIGHLY EFFECTIVE AND ADVANCED FORMULA CONTAINS TWO GERMICIDES AND 10% GLYCERIN.

ACTIVE INGREDIENT

This product contains highly effective levels of free odine for fast kill of mastitis causing organisms.

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

- 1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
- 2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
- 3. Do not use this product for cleaning or sanitizing equipment
- 4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION A. PRE-MILKING (PRE-DIPPING)

- 1. REMOVE VISIBLE SOILS FROM THE TEATS.
- 2. OBSERVE FOREMILK BY STRIPPING TWO OR THREE STREAMS OF MILK INTO A STRIP CUP.
- 3. DIP OR SPRAY THE COW'S TEATS FULL LENGTH WITH TANDEM+ BARRIER TEAT DIP. 4 WAIT APPROXIMATELY 30 SECONDS.
- 5. REMOVE ALL TEAT DIP BY THOROUGHLY DRYING THE TEATS AND TEAT ORIFICES WITH A CLEAN, SINGLE SERVICE PAPER TOWEL(S).
- 6. ATTACH MILKING UNIT

B. POST-MILKING (POST DIPPING)

Immediately after milking dip all teats full length with TANDEM+ BARRIER Teat Dip Ensure good coverage of all sides of the teats and teat orifices.

WARNING

DO NOT DILUTE THIS PRODUCT

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS. PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AIDE PROCEDURES


EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

TANDEM+ BARRIER



TANDEM PLUS BARRIER

iodine solution

Product Information					
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-0074		
Route of Administration	TOPICAL				

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	2.50 g in 1 L		
DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)	DODECYLBENZ ENES ULFONIC ACID	1.90 g in 1 L		

Inactive Ingredients		
Ingredient Name	Strength	
XANTHAN GUM (UNII: TTV12P4NEE)		
POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9)		

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:29983-0074-4	18.9 L in 1 DRUM					
2	NDC:29983-0074-6	56.7 L in 1 PACKAGE					
3	NDC:29983-0074-8	207.9 L in 1 DRUM					
4	NDC:29983-0074-9	1039.5 L in 1 PACKAGE					

Marketing Information					
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
	04/21/2010				
	Application Number or Monograph	Application Number or Monograph Marketing Start Citation Date			

Labeler - IBA (019494160)

Revised: 12/2024 IBA