

LOW DOSE MINIPRIN ENTERIC SAFETY COATED- aspirin tablet, delayed release
Time-Cap Labs, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

330R ASA 81 MG MINIPRIN

active ingredient (in each tablet) Purpose

Aspirin 81 mg (NSAID*).....Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains
 - for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effects may occur
-

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, d&c yellow#10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, Polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
-

Ask a doctor before use if

- stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 - you are taking a diuretic
 - you have asthma
 - you have not been drinking fluids
 - you have lost a lot of fluid due to vomiting or diarrhea
-

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
 - taking any other drug
 - under a doctor's care for any serious condition
-

-
-

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - ◦ feel faint
 - ◦ have bloody or black stools
 - ◦ vomit blood
 - ◦ have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - ringing in the ears or a loss of hearing occurs
-

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

PAIN RELIEVER

-
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
-

Time-Cap Labs, Inc. NDC 49483-330-10
 *Compare to the active ingredient in Bayer® Low Dose

LOW DOSE ASPIRIN REGIMEN
MINIPRIN®
ENTERIC
 Safety Coated

Pain Reliever (NSAID)
 SEE NEW WARNINGS INFORMATION

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

1000 ENTERIC COATED TABLETS New Improved Formula

Drug Facts
 Active ingredient (in each tablet) Purpose
 Aspirin 81 mg (NSAID)* Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses • temporarily relieves minor aches and pains
 • for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effects may occur

Warnings
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
 Allergy alert: Aspirin may cause a severe allergic reaction, which may include: • hives • facial swelling • shock • asthma (wheezing)
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs

Drug Facts (continued)
 (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed
 Do not use • if you are allergic to aspirin or any other pain reliever/fever reducer
 Ask a doctor before use if: • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have not been drinking fluids • you have lost a lot of fluid due to vomiting or diarrhea
 Ask a doctor or pharmacist before use if you are: • taking a prescription drug, for diabetes, gout, or arthritis • under a doctor's care for any serious condition • taking any other drug
 Stop use and ask a doctor if: • you experience any of the following signs of stomach bleeding: • belching • have bloody or black stools • vomit blood • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear

Drug Facts (continued)
 • ringing in the ears or a loss of hearing occurs
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions • drink a full glass of water with each dose • adults and children 12 years and over: take 4 to 6 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years: consult a doctor

Other information
 • store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) • use by expiration date on package

Inactive ingredients anhydrous lactose, croscarmellose sodium, colloidal silicon dioxide, croscarmellose sodium, D5C yellow #10 aluminum lake, iron oxide color, methacrylic acid copolymer, microcrystalline cellulose, poly sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer® Low Dose. 3308 0810

Manufactured by: Time-Cap Labs, Inc.
 7 Michael Avenue, Palm Springs, NY 11765

WARNING: DO NOT USE THIS PRODUCT IF THE IMPRINT PAIN RELIEVER INFORMATION ON THE BOTTOM OF THE TABLET IS BLIND, UNREADABLE OR MISSING

LOT #: EXP. DATE:

LOW DOSE MINIPRIN ENTERIC SAFETY COATED

aspirin tablet, delayed release

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:49483-330 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 81 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------|----------|
| ANHYDROUS LACTOSE (UNII: 3S5YLH9PMK) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| TALC (UNII: 7SEV7J4R1U) | |

| | |
|---|--|
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FERRIC OXIDE YELLOW (UNII: EX438O2MRT) | |
| METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| SODIUM HYDROXIDE (UNII: 55X04QC32I) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|----------|--------|--------------|------------------------------|
| Color | yellow | Score | no score |
| Shape | ROUND | Size | 7mm |
| Flavor | | Imprint Code | embossed;upper;8;lower;plain |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:49483-330-12 | 120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 12/19/2018 | |
| 2 | NDC:49483-330-10 | 1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 12/19/2018 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part343 | 08/11/2010 | |

Labeler - Time-Cap Labs, Inc (037052099)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------------|---------|-----------|------------------------|
| Time-Cap Labs, Inc | | 037052099 | manufacture(49483-330) |