

FAMOTIDINE 20 MG - famotidine 20 mg tablet, coated
MARKSANS PHARMA LIMITED

Famotidine Tablets USP, 20 mg - Cool Mint Flavor

ACTIVE INGREDIENT(S)

Active ingredient (in each tablet)

Famotidine USP 20 mg

PURPOSE

Purpose

Acid reducer

USE(S)

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

WARNINGS

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

DO NOT USE

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

ASK A DOCTOR BEFORE USE IF

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting

- stomach pain
- kidney disease

ASK A DOCTOR OR PHARMACIST BEFORE USE IF

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR IF

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

PREGNANCY/BREASTFEEDING

If pregnant or breast-feeding, ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- Meets USP dissolution test 2

INACTIVE INGREDIENTS

Inactive ingredients carnauba wax, colloidal silicon dioxide, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized

starch, sodium saccharin, talc, titanium dioxide

QUESTIONS OR COMMENTS

Questions or comments? Call **1-877-376-4271** (weekdays 9 AM to 5 PM)

• **JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Tips for Managing Heartburn


- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking


Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue,
Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83
Verna Indl. Estate
Verna, Goa-403722, India


PRINCIPAL DISPLAY PANEL

NDC: 25000-124-94
Famotidine Tablets USP 20 mg- Cool mint flavor
25's (5 x 5's blister) count Carton Label





NDC: 25000-124-84
Famotidine Tablets USP 20 mg- Cool mint flavor



Marksans

NDC 25000-124-84


MAXIMUM STRENGTH

Famotidine Tablets USP, 20 mg


Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn
Due to Acid Indigestion



Cool Mint
Flavor



ACTUAL SIZE


25 Tablets

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts
<p>Active ingredient Purpose (in each tablet)</p> <p>Famotidine USP 20 mg.....Acid reducer</p>
<p>Uses</p> <ul style="list-style-type: none"> ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages ▼
<p>Drug Facts (continued under label)</p>
<p>Manufactured for:</p> <p>Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735, USA</p> <p>Manufactured by:</p> <p>Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa-403722, India</p> <p>Mfg. Lic. No.:GO/DRUGS/515</p> <p>PO90924</p>

LOT: _____

EXP: _____

PEEL HERE 

Varnish Omit Area



LOT: Varnish Omit
EXP:

Manufactured for: **Time-Cap Labs, Inc.**
7 Michael Avenue, Farmingdale, NY 11735, USA
Manufactured by: **Marksans Pharma Ltd.**
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, Goa-403722, India
Mfg. Lic. No. GOORUGS515 R00924

Drug Facts

Active ingredient
(in each tablet) **Purpose**
Famotidine USP 20 mg..... Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms,

Drug Facts (continued)

- neck or shoulders, or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn

Tips for Managing Heartburn

- Quit smoking
- If overweight, lose weight
- Eat slowly and avoid big meals
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine,
- Raise the head of your bed
- Do not eat before bedtime
- Do not wear tight-fitting clothing around the stomach
- Do not lie flat or bend over after eating

Drug Facts (continued)

- do not use more than 2 tablets in 24 hours
- children under 12 years, ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- Meets USP dissolution test 2

Inactive ingredients

carminative wax, colloidal silicon dioxide, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium saccharin, talc, titanium dioxide

Questions or comments?

Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

- **JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc. Marksans Pharma Ltd. is the owner of the registered trademark Maximum Strength Peppid® AC.



NDC 25000-124-84
Compare to the active ingredient in Maximum Strength Peppid® AC††

MAXIMUM STRENGTH

Famotidine Tablets USP, 20 mg

Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn
Due to Acid Indigestion



Varnish/Print Omit Area

Varnish Omit

Varnish Omit

NDC: 25000-124-68
Famotidine Tablets USP 20 mg- Cool mint flavor
300's count Bottle Label

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)			FAMOTIDINE	20 mg
Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
CARNAUBA WAX (UNII: R12CBM0EIZ)				
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I3O)				
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)				
STARCH, CORN (UNII: O8232NY3SJ)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	WHITE	Score	no score	
Shape	ROUND	Size	8mm	
Flavor	MINT	Imprint Code	124	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-124-94	5 in 1 CARTON	05/23/2025	
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:25000-124-84	1 in 1 CARTON	05/23/2025	
2		25 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:25000-124-68	300 in 1 BOTTLE; Type 0: Not a Combination Product	05/23/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA217543		05/23/2025	

Labeler - MARKSANS PHARMA LIMITED (925822975)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-124)

Revised: 5/2025

MARKSANS PHARMA LIMITED