

FRINDOVYX- cyclophosphamide injection

Avyxa Pharma, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use FRINDOVYX safely and effectively. See full prescribing information for FRINDOVYX.

FRINDOVYX™ (cyclophosphamide) injection, for intravenous use
Initial U.S. Approval: 1959

-----RECENT MAJOR CHANGES-----

Dosage and Administration (2.2, 2.3) 9/2024
Warnings and Precautions, Embryo Fetal Toxicity (5.7)
9/2024

-----INDICATIONS AND USAGE-----

FRINDOVYX is an alkylating drug indicated for treatment of adults and pediatric patients with:

- Malignant Diseases: malignant lymphomas: Hodgkin's disease, lymphocytic lymphoma, mixed-cell type lymphoma, histiocytic lymphoma, Burkitt's lymphoma; multiple myeloma, leukemias, mycosis fungoides, neuroblastoma, adenocarcinoma of ovary, retinoblastoma, breast carcinoma (1)

-----DOSAGE AND ADMINISTRATION-----

During or immediately after FRINDOVYX administration, administer adequate amounts of fluid to reduce the risk of urinary tract toxicity (2.1).

Malignant Diseases: Adult and Pediatric Patients (2.2)

- Intravenous: Initial course for patients with no hematologic deficiency: 40 mg/kg to 50 mg/kg in divided doses over 2 to 5 days. Other regimens include 10 mg/kg to 15 mg/kg given every 7 to 10 days or 3 mg/kg to 5 mg/kg twice weekly. (2.2)
- See full prescribing information for instructions on preparation, handling, and administration. (2.3)

-----DOSAGE FORMS AND STRENGTHS-----

- Injection: 500 mg/mL, 1 g/2 mL (500 mg/mL), and 2 g/4mL (500 mg/mL) in multiple-dose vial (3)

-----CONTRAINDICATIONS-----

- Hypersensitivity to cyclophosphamide (4)
- Urinary outflow obstruction (4)

-----WARNINGS AND PRECAUTIONS-----

- Myelosuppression, Immunosuppression, Bone Marrow Failure and Infections: Severe immunosuppression may lead to serious and sometimes fatal infections. Close hematological monitoring is required. (5.1)
- Urinary Tract and Renal Toxicity: Hemorrhagic cystitis, pyelitis, ureteritis, and hematuria can occur. Urotoxicity can be fatal. Exclude or correct any urinary tract obstructions prior to treatment. (5.2)
- Cardiotoxicity: Myocarditis, myopericarditis, pericardial effusion, arrhythmias and congestive heart failure, which may be fatal, have been reported. Monitor patients, especially those with risk factors for cardiotoxicity or pre-existing cardiac disease. (5.3)
- Pulmonary Toxicity: Pneumonitis, pulmonary fibrosis and pulmonary veno-occlusive disease leading to respiratory failure may occur. Monitor patients for signs and symptoms of pulmonary toxicity. (5.4)
- Secondary malignancies(5.5)
- Veno-occlusive Liver Disease: Fatal outcome can occur. (5.6)
- Alcohol content: The alcohol content in a dose of FRINDOVYX may affect the central nervous system. This may include impairment of a patient's ability to drive or use machines immediately after infusion. (5.7)
- Embryo-Fetal Toxicity: Can cause fetal harm. Advise patients of reproductive potential of the potential risk to a fetus and to use effective contraception. (5.8, 8.1, 8.3)

-----ADVERSE REACTIONS-----

Adverse reactions reported most often include neutropenia, febrile neutropenia, fever, alopecia, nausea, vomiting, and diarrhea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Avyxa Pharma, LLC at 1-888-520-0954

or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **USE IN SPECIFIC POPULATIONS** -----

- Lactation: Advise not to breastfeed. (8.2)
- Renal Impairment: Monitor for toxicity in patients with moderate and severe renal impairment. (8.6, 12.3)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Administration Instructions
- 2.2 Recommended Dosage for Malignant Diseases
- 2.3 Preparation, Handling and Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Myelosuppression, Immunosuppression, Bone Marrow Failure and Infections
- 5.2 Urinary Tract and Renal Toxicity
- 5.3 Cardiotoxicity
- 5.4 Pulmonary Toxicity
- 5.5 Secondary Malignancies
- 5.6 Veno-occlusive Liver Disease
- 5.7 Alcohol Content
- 5.8 Embryo-Fetal Toxicity
- 5.9 Infertility
- 5.10 Impairment of Wound Healing
- 5.11 Hyponatremia

6 ADVERSE REACTIONS

- 6.1 Clinical Trials and Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Effect of Other Drugs on Cyclophosphamide Exposure
- 7.2 Drugs that Potentiate Cyclophosphamide Toxicities
- 7.3 Effect of Cyclophosphamide on Other Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Malignant Diseases

FRINDOVYX is indicated for the treatment of adult and pediatric patients with:

- malignant lymphomas (Stages III and IV of the Ann Arbor staging system), Hodgkin's disease, lymphocytic lymphoma (nodular or diffuse), mixed-cell type lymphoma, histiocytic lymphoma Burkitt's lymphoma
- multiple myeloma
- leukemias: chronic lymphocytic leukemia, chronic granulocytic leukemia (it is usually ineffective in acute blastic crisis), acute myelogenous and monocytic leukemia, acute lymphoblastic (stem-cell) leukemia (cyclophosphamide given during remission is effective in prolonging its duration)
- mycosis fungoides (advanced disease)
- neuroblastoma (disseminated disease)
- adenocarcinoma of the ovary
- retinoblastoma
- carcinoma of the breast

Cyclophosphamide, although effective alone in susceptible malignancies, is more frequently used concurrently or sequentially with other antineoplastic drugs.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

During or immediately after the administration of FRINDOVYX, adequate amounts of fluid should be ingested or infused to force diuresis in order to reduce the risk of urinary tract toxicity. Therefore, FRINDOVYX should be administered in the morning.

2.2 Recommended Dosage for Malignant Diseases

Adults and Pediatric Patients

Intravenous

When used as the only oncolytic drug therapy, the recommended dosage for the initial

course of FRINDOVYX for patients with no hematologic deficiency is 40 mg/kg to 50 mg/kg given intravenously in divided doses over a period of 2 to 5 days. Other intravenous regimens include 10 mg/kg to 15 mg/kg given every 7 to 10 days or 3 mg/kg to 5 mg/kg twice weekly.

Adjust the dosage of FRINDOVYX based on the specific regimen administered, response to treatment, myelosuppression or other adverse reactions, and patient risk factors [see *Warnings and Precautions (5)*].

2.3 Preparation, Handling and Administration

FRINDOVYX is a hazardous drug. Follow applicable special handling and disposal procedures¹. Caution should be exercised when handling and preparing FRINDOVYX. To minimize the risk of dermal exposure, always wear gloves when handling vials containing FRINDOVYX.

FRINDOVYX

Intravenous Administration

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use cyclophosphamide vials if there are signs of particulate matter.

Cyclophosphamide does not contain any antimicrobial preservative and thus care must be taken to assure the sterility of prepared solutions. Use aseptic technique.

For Direct Intravenous Injection

Withdraw the prescribed dose of FRINDOVYX from the vial with a syringe and dilute with 0.9% Sodium Chloride Injection, USP to a concentration of 20 mg/mL of cyclophosphamide.

For Intravenous Infusion

Withdraw the prescribed dose of FRINDOVYX from the vial with a syringe and dilute FRINDOVYX to a concentration of 2 mg/mL with any of the following diluents:

- 0.45% Sodium Chloride Injection, USP
- 5% Dextrose Injection, USP
- 5% Dextrose and 0.9% Sodium Chloride Injection, USP

To reduce the likelihood of adverse reactions that appear to be administration rate-dependent (e.g., facial swelling, headache, nasal congestion, scalp burning), cyclophosphamide should be injected or infused very slowly. Duration of the infusion also should be appropriate for the volume and type of carrier fluid to be infused.

Storage of Diluted Cyclophosphamide Solution:

If not used immediately, for microbiological integrity, cyclophosphamide solutions should be stored as described in Table 1.

Table 1: Storage of Cyclophosphamide Solutions

Diluent	Storage	
	Room	

Diluent	Room Temperature	Refrigerated
Diluted Solution (20 mg/mL) for Direct Intravenous Injection		
0.9% Sodium Chloride Injection, USP	up to 24 hours	up to 6 days
Diluted Solutions (2 mg/mL) for Intravenous Infusion		
0.45% Sodium Chloride Injection, USP	up to 24 hours	up to 6 days
5% Dextrose Injection, USP	up to 24 hours	up to 36 hours
5% Dextrose and 0.9% Sodium Chloride Injection, USP	up to 24 hours	up to 36 hours

Storage of Undiluted Cyclophosphamide Solution (Multiple-Dose Vial):

After first use, store partially used multiple-dose vial in the original carton refrigerated at 2°C to 8°C (36°F to 46°F) for up to 28 days or at room temperature at 20°C to 25°C (68°F to 77°F) for up to 7 days. Discard unused portion.

3 DOSAGE FORMS AND STRENGTHS

FRINDOVYX: Clear, colorless or slight yellow, sterile solution of cyclophosphamide in multiple-dose vial in the following three strengths.

- 500 mg/mL
- 1 g/2 mL (500 mg/mL)
- 2 g/4 mL (500 mg/mL)

4 CONTRAINDICATIONS

Hypersensitivity

FRINDOVYX is contraindicated in patients who have a history of severe hypersensitivity reactions to cyclophosphamide, any of its metabolites, or to other components of the product. Anaphylactic reactions including death have been reported with cyclophosphamide. Cross-sensitivity with other alkylating agents can occur.

Urinary Outflow Obstruction

FRINDOVYX is contraindicated in patients with urinary outflow obstruction [see *Warnings and Precautions (5.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 Myelosuppression, Immunosuppression, Bone Marrow Failure and Infections

Cyclophosphamide can cause myelosuppression (leukopenia, neutropenia, thrombocytopenia and anemia), bone marrow failure, and severe immunosuppression which may lead to serious and sometimes fatal infections, including sepsis and septic shock. Latent infections can be reactivated [see *Adverse Reactions (6)*].

Antimicrobial prophylaxis may be indicated in certain cases of neutropenia at the discretion of the managing physician. In case of neutropenic fever, antibiotic therapy is indicated. Antimycotics and/or antivirals may also be indicated.

Monitoring of complete blood counts is essential during cyclophosphamide treatment so that the dose can be adjusted, if needed. Cyclophosphamide should not be administered to patients with neutrophils $\leq 1,500/\text{mm}^3$ and platelets $< 50,000/\text{mm}^3$.

Cyclophosphamide treatment may not be indicated, or should be interrupted, or the dose reduced, in patients who have or who develop a serious infection. G-CSF may be administered to reduce the risks of neutropenia complications associated with cyclophosphamide use. Primary and secondary prophylaxis with G-CSF should be considered in all patients considered to be at increased risk for neutropenia complications. The nadirs of the reduction in leukocyte count and thrombocyte count are usually reached in weeks 1 and 2 of treatment. Peripheral blood cell counts are expected to normalize after approximately 20 days. Bone marrow failure has been reported. Severe myelosuppression may be expected particularly in patients pretreated with and/or receiving concomitant chemotherapy and/or radiation therapy.

5.2 Urinary Tract and Renal Toxicity

Hemorrhagic cystitis, pyelitis, ureteritis, and hematuria have been reported with cyclophosphamide. Medical and/or surgical supportive treatment may be required to treat protracted cases of severe hemorrhagic cystitis. Discontinue cyclophosphamide therapy in case of severe hemorrhagic cystitis.

Urotoxicity (bladder ulceration, necrosis, fibrosis, contracture and secondary cancer) may require interruption of cyclophosphamide treatment or cystectomy. Urotoxicity can be fatal. Urotoxicity can occur with short-term or long-term use of cyclophosphamide.

Before starting treatment, exclude or correct any urinary tract obstructions [see *Contraindications (4)*]. Urinary sediment should be checked regularly for the presence of erythrocytes and other signs of urotoxicity and/or nephrotoxicity. Cyclophosphamide should be used with caution, if at all, in patients with active urinary tract infections. Aggressive hydration with forced diuresis and frequent bladder emptying can reduce the frequency and severity of bladder toxicity. Mesna has been used to prevent severe bladder toxicity.

5.3 Cardiotoxicity

Myocarditis, myopericarditis, pericardial effusion including cardiac tamponade, and congestive heart failure, which may be fatal, have been reported with cyclophosphamide therapy.

Supraventricular arrhythmias (including atrial fibrillation and flutter) and ventricular arrhythmias (including severe QT prolongation associated with ventricular tachyarrhythmia) have been reported after treatment with regimens that included cyclophosphamide.

The risk of cardiotoxicity may be increased with high doses of cyclophosphamide, in patients with advanced age, and in patients with previous radiation treatment to the cardiac region and/or previous or concomitant treatment with other cardiotoxic agents.

Particular caution is necessary in patients with risk factors for cardiotoxicity and in patients with pre-existing cardiac disease.

Monitor patients with risk factors for cardiotoxicity and with pre-existing cardiac disease.

5.4 Pulmonary Toxicity

Pneumonitis, pulmonary fibrosis, pulmonary veno-occlusive disease and other forms of pulmonary toxicity leading to respiratory failure have been reported during and following treatment with cyclophosphamide. Late onset pneumonitis (greater than 6 months after start of cyclophosphamide) appears to be associated with increased mortality. Pneumonitis may develop years after treatment with cyclophosphamide.

Monitor patients for signs and symptoms of pulmonary toxicity.

5.5 Secondary Malignancies

Cyclophosphamide is genotoxic [see *Nonclinical Toxicology (13.1)*]. Secondary malignancies (urinary tract cancer, myelodysplasia, acute leukemias, lymphomas, thyroid cancer, and sarcomas) have been reported in patients treated with cyclophosphamide-containing regimens. The risk of bladder cancer may be reduced by prevention of hemorrhagic cystitis.

5.6 Veno-occlusive Liver Disease

Veno-occlusive liver disease (VOD) including fatal outcome has been reported in patients receiving cyclophosphamide-containing regimens. A cytoreductive regimen in preparation for bone marrow transplantation that consists of cyclophosphamide in combination with whole-body irradiation, busulfan, or other agents has been identified as a major risk factor. VOD has also been reported to develop gradually in patients receiving long-term low-dose immunosuppressive doses of cyclophosphamide. Other risk factors predisposing to the development of VOD include preexisting disturbances of hepatic function, previous radiation therapy of the abdomen, and a low performance status.

5.7 Alcohol Content

The alcohol content in a dose of FRINDOVYX may affect the central nervous system and should be taken into account for patients in whom alcohol intake should be avoided or minimized. Consideration should be given to the alcohol content in FRINDOVYX on the ability to drive or use machines immediately after the infusion.

Each administration of FRINDOVYX at 50 mg per kg delivers 0.0448 g/kg of ethanol. For a 75 kg patient this would deliver 3.36 grams of ethanol [see *Description (11)*]. Other cyclophosphamide products may have a different amount of alcohol or no alcohol.

5.8 Embryo-Fetal Toxicity

Based on its mechanism of action and published reports of effects in pregnant patients or animals, FRINDOVYX can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*, *Clinical Pharmacology (12.1)*, and *Nonclinical Toxicology (13.1)*]. Exposure to cyclophosphamide during pregnancy may cause birth defects, miscarriage, fetal growth retardation, and fetotoxic effects in the newborn. Cyclophosphamide is teratogenic and embryo-fetal toxic in mice, rats, rabbits and monkeys.

Advise pregnant women and females of reproductive potential of the potential risk to the fetus [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with FRINDOVYX and for up to 1 year after completion of therapy. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with FRINDOVYX and for 4 months after completion of therapy [see *Use in Specific Populations (8.1, 8.3)*].

5.9 Infertility

Male and female reproductive function and fertility may be impaired in patients being treated with FRINDOVYX. Cyclophosphamide interferes with oogenesis and spermatogenesis. It may cause sterility in both sexes. Development of sterility appears to depend on the dose of cyclophosphamide, duration of therapy, and the state of gonadal function at the time of treatment. Cyclophosphamide-induced sterility may be irreversible in some patients. Advise patients on the potential risks for infertility [see *Use in Specific Populations (8.3 and 8.4)*].

5.10 Impairment of Wound Healing

Cyclophosphamide may interfere with normal wound healing.

5.11 Hyponatremia

Hyponatremia associated with increased total body water, acute water intoxication, and a syndrome resembling SIADH (syndrome of inappropriate secretion of antidiuretic hormone), which may be fatal, has been reported.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in more detail in other sections of the labeling.

- Hypersensitivity [see *Contraindications (4)*]
- Myelosuppression, Immunosuppression, Bone Marrow Failure, and Infections [see *Warnings and Precautions (5.1)*]
- Urinary Tract and Renal Toxicity [see *Warnings and Precautions (5.2)*]
- Cardiotoxicity [see *Warnings and Precautions (5.3)*]
- Pulmonary Toxicity [see *Warnings and Precautions (5.4)*]
- Secondary Malignancies [see *Warnings and Precautions (5.5)*]

- Veno-occlusive Liver Disease [see Warnings and Precautions (5.6)]
- Alcohol Content [see Warnings and Precautions (5.7)]
- Infertility [see Warnings and Precautions (5.9) and Use in Specific Populations (8.3 and 8.4)]
- Impaired Wound Healing [see Warnings and Precautions (5.10)]
- Hyponatremia [see Warnings and Precautions (5.11)]

6.1 Clinical Trials and Postmarketing Experience

The following adverse reactions associated with the use of cyclophosphamide were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The most common adverse reactions were neutropenia, febrile neutropenia, fever, alopecia, nausea, vomiting, and diarrhea.

Cardiac: cardiac arrest, ventricular fibrillation, ventricular tachycardia, cardiogenic shock, pericardial effusion (progressing to cardiac tamponade), myocardial hemorrhage, myocardial infarction, cardiac failure (including fatal outcomes), cardiomyopathy, myocarditis, pericarditis, carditis, atrial fibrillation, supraventricular arrhythmia, ventricular arrhythmia, bradycardia, tachycardia, palpitations, QT prolongation.

Congenital, Familial and Genetic: intra-uterine death, fetal malformation, fetal growth retardation, fetal toxicity (including myelosuppression, gastroenteritis).

Ear and Labyrinth: deafness, hearing impaired, tinnitus.

Endocrine: water intoxication.

Eye: visual impairment, conjunctivitis, lacrimation.

Gastrointestinal: gastrointestinal hemorrhage, acute pancreatitis, colitis, enteritis, cecitis, stomatitis, constipation, parotid gland inflammation, nausea, vomiting, diarrhea.

General Disorders and Administrative Site Conditions: multiorgan failure, general physical deterioration, influenza-like illness, injection/infusion site reactions (thrombosis, necrosis, phlebitis, inflammation, pain, swelling, erythema), pyrexia, edema, chest pain, mucosal inflammation, asthenia, pain, chills, fatigue, malaise, headache, febrile neutropenia.

Hematologic: myelosuppression, bone marrow failure, disseminated intravascular coagulation and hemolytic uremic syndrome (with thrombotic microangiopathy).

Hepatic: veno-occlusive liver disease, cholestatic hepatitis, cytolytic hepatitis, hepatitis, cholestasis; hepatotoxicity with hepatic failure, hepatic encephalopathy, ascites, hepatomegaly, blood bilirubin increased, hepatic function abnormal, hepatic enzymes increased.

Immune: immunosuppression, anaphylactic shock and hypersensitivity reaction.

Infections: The following manifestations have been associated with myelosuppression and immunosuppression caused by cyclophosphamide: increased risk for and severity of pneumonias (including fatal outcomes), other bacterial, fungal, viral, protozoal and, parasitic infections; reactivation of latent infections, (including viral hepatitis, tuberculosis), *Pneumocystis jiroveci*, herpes zoster, *Strongyloides*, sepsis and septic shock.

Investigations: blood lactate dehydrogenase increased, C-reactive protein increased.

Metabolism and Nutrition: hyponatremia, fluid retention, blood glucose increased, blood glucose decreased.

Musculoskeletal and Connective Tissue: rhabdomyolysis, scleroderma, muscle spasms, myalgia, arthralgia.

Neoplasms: acute leukemia, myelodysplastic syndrome, lymphoma, sarcomas, renal cell carcinoma, renal pelvis cancer, bladder cancer, ureteric cancer, thyroid cancer.

Nervous System: encephalopathy, convulsion, dizziness, neurotoxicity has been reported and manifested as reversible posterior leukoencephalopathy syndrome, myelopathy, peripheral neuropathy, polyneuropathy, neuralgia, dysesthesia, hypoesthesia, paresthesia, tremor, dysgeusia, hypogeusia, parosmia.

Pregnancy: premature labor.

Psychiatric: confusional state.

Renal and Urinary: renal failure, renal tubular disorder, renal impairment, nephropathy toxic, hemorrhagic cystitis, bladder necrosis, cystitis ulcerative, bladder contracture, hematuria, nephrogenic diabetes insipidus, atypical urinary bladder epithelial cells.

Reproductive System: infertility, ovarian failure, ovarian disorder, amenorrhea, oligomenorrhea, testicular atrophy, azoospermia, oligospermia.

Respiratory: pulmonary veno-occlusive disease, acute respiratory distress syndrome, interstitial lung disease as manifested by respiratory failure (including fatal outcomes), obliterative bronchiolitis, organizing pneumonia, alveolitis allergic, pneumonitis, pulmonary hemorrhage; respiratory distress, pulmonary hypertension, pulmonary edema, pleural effusion, bronchospasm, dyspnea, hypoxia, cough, nasal congestion, nasal discomfort, oropharyngeal pain, rhinorrhea.

Skin and Subcutaneous Tissue: toxic epidermal necrolysis, Stevens-Johnson syndrome, erythema multiforme, palmar-plantar erythrodysesthesia syndrome, radiation recall dermatitis, toxic skin eruption, urticaria, dermatitis, blister, pruritus, erythema, nail disorder, facial swelling, hyperhidrosis, alopecia.

Tumor lysis syndrome: like other cytotoxic drugs, cyclophosphamide may induce tumor-lysis syndrome and hyperuricemia in patients with rapidly growing tumors.

Vascular: pulmonary embolism, venous thrombosis, vasculitis, peripheral ischemia, hypertension, hypotension, flushing, hot flush.

7 DRUG INTERACTIONS

7.1 Effect of Other Drugs on Cyclophosphamide Exposure

Protease Inhibitors

Concomitant use of protease inhibitors may increase the concentration of cytotoxic metabolites and may enhance the toxicities of cyclophosphamide, including higher incidence of infections, neutropenia, and mucositis. Monitor for increased toxicities in patients receiving protease inhibitors.

7.2 Drugs that Potentiate Cyclophosphamide Toxicities

Drugs or agents with similar toxicities to FRINDOVYX and can potentiate these effects are listed in Table 2.

Table 2: Drugs that Potentiate Cyclophosphamide Toxicities

Toxicity	Drugs
Increased hematotoxicity and/or immunosuppression	<ul style="list-style-type: none"> ● ACE inhibitors: ACE inhibitors can cause leukopenia. ● Natalizumab ● Paclitaxel: Increased hematotoxicity has been reported when cyclophosphamide was administered after paclitaxel infusion. ● Thiazide diuretics ● Zidovudine
Increased cardiotoxicity	<ul style="list-style-type: none"> ● Anthracyclines ● Cytarabine ● Pentostatin ● Radiation therapy of the cardiac region ● Trastuzumab
Increased pulmonary toxicity	<ul style="list-style-type: none"> ● Amiodarone ● G-CSF, GM-CSF (granulocyte colony-stimulating factor, granulocyte macrophage colony-stimulating factor)
Increased nephrotoxicity	<ul style="list-style-type: none"> ● Amphotericin B ● Indomethacin: Acute water intoxication has been reported with concomitant use of indomethacin
Increase in other toxicities:	<ul style="list-style-type: none"> ● Azathioprine: Increased risk of hepatotoxicity (liver necrosis) ● Busulfan: Increased incidence of hepatic veno-occlusive disease and mucositis has been reported. ● Protease inhibitors: Increased incidence of mucositis
Increased risk of hemorrhagic cystitis	<ul style="list-style-type: none"> ● Radiation treatment: Increased risk of hemorrhagic cystitis may result from a combined effect of cyclophosphamide and past or concomitant radiation treatment.

7.3 Effect of Cyclophosphamide on Other Drugs

Metronidazole

Acute encephalopathy has been reported in a patient receiving cyclophosphamide and metronidazole. Monitor for neurologic toxicities in patients receiving metronidazole.

Tamoxifen

Concomitant use of tamoxifen and a cyclophosphamide-containing chemotherapy regimen may increase the risk of thromboembolic complications. Monitor for signs and symptoms of thromboembolic events in patients receiving tamoxifen.

Coumarins

Both increased and decreased warfarin effect have been reported in patients receiving warfarin and cyclophosphamide. Monitor anticoagulant activity closely in patients receiving warfarin or other coumarins.

Cyclosporine

Concomitant administration of cyclophosphamide may decrease serum concentrations of cyclosporine. This interaction may result in an increased incidence of graft-versus-host disease. Monitor for signs and symptoms of graft-versus-host disease in patients receiving cyclosporine.

Depolarizing Muscle Relaxants

If a patient has been treated with cyclophosphamide within 10 days of general anesthesia, alert the anesthesiologist.

Cyclophosphamide causes a marked and persistent inhibition of cholinesterase activity. Prolonged apnea may occur with concurrent depolarizing muscle relaxants (e.g., succinylcholine).

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on its mechanism of action and published reports of effects in pregnant patients or animals, FRINDOVYX can cause fetal harm when administered to a pregnant woman [see *Clinical Pharmacology (12.1)* and *Nonclinical Toxicology (13.1)*]. Exposure to cyclophosphamide during pregnancy may cause fetal malformations, miscarriage, fetal growth retardation, and toxic effects in the newborn [see *Data*]. Cyclophosphamide is teratogenic and embryo-fetal toxic in mice, rats, rabbits and monkeys [see *Data*]. Advise pregnant women and females of reproductive potential of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects is 2% - 4% and of miscarriage is 15% - 20% of clinically recognized pregnancies.

Data

Human Data

Malformations of the skeleton, palate, limbs and eyes as well as miscarriage have been reported after exposure to cyclophosphamide in the first trimester. Fetal growth retardation and toxic effects manifesting in the newborn, including leukopenia, anemia, pancytopenia, severe bone marrow hypoplasia, and gastroenteritis have been reported after exposure to cyclophosphamide.

Animal Data

Administration of cyclophosphamide to pregnant mice, rats, rabbits and monkeys during the period of organogenesis at doses at or below the dose in patients based on body surface area resulted in various malformations, which included neural tube defects, limb and digit defects and other skeletal anomalies, cleft lip and palate, and reduced skeletal ossification.

8.2 Lactation

Risk Summary

Cyclophosphamide is present in breast milk. Neutropenia, thrombocytopenia, low hemoglobin, and diarrhea have been reported in infants breast fed by women treated with cyclophosphamide. Because of the potential for serious adverse reactions in a breastfed child, advise lactating women not to breastfeed during treatment with FRINDOVYX and for 1 week after the last dose.

8.3 Females and Males of Reproductive Potential

FRINDOVYX can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

Pregnancy Testing

Verify the pregnancy status of females of reproductive potential prior to the initiation of FRINDOVYX [see *Use in Specific Populations (8.1)*].

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment with FRINDOVYX and for up to 1 year after completion of therapy.

Males

Based on findings in genetic toxicity and animal reproduction studies, advise male patients with female partners of reproductive potential to use effective contraception during treatment with FRINDOVYX and for 4 months after completion of therapy [see *Nonclinical Toxicology (13.1)*].

Infertility

Females

Amenorrhea, transient or permanent, associated with decreased estrogen and increased gonadotropin secretion develops in a proportion of women treated with cyclophosphamide. Affected patients generally resume regular menses within a few months after cessation of therapy. The risk of premature menopause with cyclophosphamide increases with age. Oligomenorrhea has also been reported in association with cyclophosphamide treatment.

Animal data suggest an increased risk of failed pregnancy and malformations may persist after discontinuation of cyclophosphamide as long as oocytes/follicles exist that were exposed to cyclophosphamide during any of their maturation phases. The exact duration of follicular development in humans is not known, but may be longer than 12 months [see *Nonclinical Toxicology (13.1)*].

Males

Men treated with cyclophosphamide may develop oligospermia or azoospermia which are normally associated with increased gonadotropin but normal testosterone secretion.

8.4 Pediatric Use

The safety and effectiveness of FRINDOVYX have been established in pediatric patients and information on this use is discussed throughout the labeling.

The alcohol content of FRINDOVYX should be taken into account when given to pediatric patients [see *Warnings and Precautions (5.7)*].

Pre-pubescent females who receive cyclophosphamide generally develop secondary sexual characteristics normally and have regular menses. Ovarian fibrosis with apparently complete loss of germ cells after prolonged administration of cyclophosphamide in late pre-pubescence has been reported. Females who received cyclophosphamide who have retained ovarian function after completing treatment are at increased risk of developing premature menopause.

Pre-pubescent males who receive cyclophosphamide develop secondary sexual characteristics normally, but may have oligospermia or azospermia and increased gonadotropin secretion. Some degree of testicular atrophy may occur. Cyclophosphamide-induced azospermia is reversible in some patients, though the reversibility may not occur for several years after cessation of therapy.

8.5 Geriatric Use

There is insufficient data from clinical studies of cyclophosphamide available for patients 65 years of age and older to determine whether they respond differently than younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac functioning, and of concomitant disease or other drug therapy.

8.6 Renal Impairment

In patients with severe renal impairment, decreased renal excretion may result in increased plasma levels of cyclophosphamide and its metabolites which may increase toxicity [see *Clinical Pharmacology (12.3)*]. Monitor patients with severe renal impairment (CL_{cr} = 10 mL/min to 24 mL/min) for signs and symptoms of toxicity.

Cyclophosphamide and its metabolites are dialyzable although there are probably quantitative differences depending upon the dialysis system. In patients requiring dialysis, consider using a consistent interval between cyclophosphamide administration and dialysis.

8.7 Hepatic Impairment

Patients with severe hepatic impairment have reduced conversion of cyclophosphamide to the active 4-hydroxyl metabolite, potentially reducing efficacy [see *Clinical Pharmacology (12.3)*]. Monitor patients with severe hepatic impairment (total bilirubin > 3 x ULN and any aspartate aminotransferase (AST)) for reduced effectiveness of cyclophosphamide.

The alcohol content of FRINDOVYX should be taken into account when given to patients with hepatic impairment [see *Warnings and Precautions (5.7)*].

10 OVERDOSAGE

No specific antidote for cyclophosphamide is known.

Overdosage should be managed with supportive measures, including appropriate

treatment for any concurrent infection, myelosuppression, or cardiac toxicity should it occur.

Serious consequences of overdose include manifestations of dose dependent toxicities such as myelosuppression, urotoxicity, cardiotoxicity (including cardiac failure), veno-occlusive hepatic disease, and stomatitis [see *Warnings and Precautions (5.1, 5.2, 5.3, and 5.6)*].


Patients who received an overdose should be closely monitored for the development of toxicities, and hematologic toxicity in particular.

Cyclophosphamide and its metabolites are dialyzable. Therefore, rapid hemodialysis is indicated when treating any suicidal or accidental overdose or intoxication [see *Clinical Pharmacology (12.3)*].

Cystitis prophylaxis with mesna may be helpful in preventing or limiting urotoxic effects with cyclophosphamide overdose.

11 DESCRIPTION

Cyclophosphamide is an alkylating drug. It is an antineoplastic drug chemically related to the nitrogen mustards. The chemical name for cyclophosphamide is (\pm)-2-[Bis(2-chloroethyl)amino]tetrahydro-2H-1,3,2-oxazaphosphorine 2-oxide monohydrate, and has the following structural formula:



Cyclophosphamide is a white crystalline powder with the molecular formula $C_7H_{15}Cl_2N_2O_2 P \cdot H_2O$ and a molecular weight of 279.1. Cyclophosphamide is soluble in water, saline, or ethanol. The pK_a of cyclophosphamide is 5.7.

FRINDOVYX is a clear, colorless to slight yellow sterile solution available as 500 mg/mL, 1 g/2 mL, and 2 g/4 mL in multiple dose vial for dilution prior to intravenous administration.

- 500 mg/mL vial contains 534.5 mg cyclophosphamide monohydrate, equivalent to 500 mg cyclophosphamide and 500 mg dehydrated alcohol (equivalent to 63% v/v).
- 1 g/2 mL vial contains 1069.0 mg cyclophosphamide monohydrate equivalent to 1 g cyclophosphamide and 1000 mg dehydrated alcohol (equivalent to 63% v/v).
- 2 g/4 mL vial contains 2138.0 mg cyclophosphamide monohydrate, equivalent to 2 g cyclophosphamide and 2000 mg dehydrated alcohol (equivalent to 63% v/v).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action has not been fully characterized. However, cross-linking of

tumor cell DNA may be involved.

The active alkylating metabolites of cyclophosphamide interfere with the growth of susceptible rapidly proliferating malignant cells.

12.2 Pharmacodynamics

Cyclophosphamide exposure-response relationships and the time course of pharmacodynamic response have not been fully characterized.

12.3 Pharmacokinetics

Cyclophosphamide is a prodrug. Cyclophosphamide pharmacokinetics are linear over the approved recommended dose range.

Distribution

The volume of distribution of cyclophosphamide is 30 to 50 L. Cyclophosphamide is approximately 20% protein bound, with no dose dependent changes. Some metabolites are greater than 60% protein bound.

Elimination

The elimination half-life ($t_{1/2}$) of cyclophosphamide ranges from 3 to 12 hours, and the clearance (CL) ranges from 4 to 5.6 L/h.

When cyclophosphamide was administered at 4 g/m² (approximately 2 times the approved recommended dosage) over a 90-minute infusion, concentration-time data demonstrate saturable elimination in parallel with first-order renal elimination.

Metabolism

Cyclophosphamide is metabolized by cytochrome P450s including CYP2A6, 2B6, 3A, 2C9, and 2C19. Cyclophosphamide is activated to form 4-hydroxycyclophosphamide, which is in equilibrium with its ring-open tautomer aldophosphamide. 4-hydroxycyclophosphamide and aldophosphamide can undergo oxidation by aldehyde dehydrogenases to form the inactive metabolites 4-ketocyclophosphamide and carboxyphosphamide, respectively. Aldophosphamide can undergo β -elimination to form active metabolites phosphoramidate mustard and acrolein. This spontaneous conversion can be catalyzed by albumin and other proteins. At high doses, the fraction of parent compound cleared by 4-hydroxylation is reduced resulting in non-linear elimination of cyclophosphamide.

Cyclophosphamide appears to induce its own metabolism. This auto-induction results in an increase in CL, increased formation of active 4-hydroxycyclophosphamide and shortened $t_{1/2}$ following multiple doses administered at 12-to 24-hour interval.

Excretion

Cyclophosphamide and its metabolites are eliminated by hepatic and renal pathways. Cyclophosphamide is primarily excreted as metabolites. Ten to 20% is excreted unchanged in the urine. A small percentage of cyclophosphamide may be eliminated unchanged in bile.

Specific Populations

Renal Impairment

Following one-hour intravenous infusion, cyclophosphamide AUC increased by 38% in patients with CLcr of 25 to 50 mL/min, by 77% in patients with CLcr of 10 to 24 mL/min and by 23% in the hemodialysis group (CLcr of < 10 mL/min) compared to the control group (CLcr ≥ 80 mL/min).

Cyclophosphamide is dialyzable. Dialysis clearance averaged 104 mL/min, which is similar to the metabolic clearance of 95 mL/min for cyclophosphamide. A mean of 37% of the administered dose of cyclophosphamide was removed during a 4-hour hemodialysis period. The $t_{1/2}$ was 3.3 hours in patients during hemodialysis, a 49% reduction compared to $t_{1/2}$ of 6.5 hours in uremic patients.

Hepatic Impairment

Cyclophosphamide CL is decreased by 40% (45 ± 8.6 L/kg) and $t_{1/2}$ is prolonged by 64% (12.5 ± 1 hours) in patients with hepatic impairment with a mean bilirubin 3.5 mg/dL and mean AST 90 IU/L compared to patients with normal hepatic function (mean bilirubin 0.5 mg/dL, mean AST 10 IU/L).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Cyclophosphamide administered by different routes, including intravenous, subcutaneous or intraperitoneal injection, or in drinking water, caused tumors in both mice and rats. In addition to leukemia and lymphoma, benign and malignant tumors were found at various tissue sites, including urinary bladder, mammary gland, lung, liver, and injection site [see *Warnings and Precautions (5.5)*].

Cyclophosphamide was mutagenic and clastogenic in multiple *in vitro* and *in vivo* genetic toxicology studies.

Cyclophosphamide is genotoxic in male and female germ cells. Animal data indicate that exposure of oocytes to cyclophosphamide during follicular development may result in a decreased rate of implantations and viable pregnancies, and in an increased risk of malformations. Male mice and rats treated with cyclophosphamide show alterations in male reproductive organs (e.g., decreased weights, atrophy, changes in spermatogenesis), and decreases in reproductive potential (e.g., decreased implantations and increased post-implantation loss) and increases in fetal malformations when mated with untreated females [see *Use in Specific Populations (8.3)*].

15 REFERENCES

1. OSHA Hazardous Drugs. *OSHA*.
<http://www.osha.gov/SLTC/hazardousdrugs/index.html>.

16 HOW SUPPLIED/STORAGE AND HANDLING

FRINDOVYX™ is a ready-to-dilute clear, colorless to slight yellow sterile solution containing cyclophosphamide and is supplied in multiple-dose vial.

FRINDOVYX™ Injection

NDC Number	Strength	Vial Presentation
83831-119-01	500 mg/mL	Multiple-dose vial, carton of 1
83831-120-02	1 g/2 mL	Multiple-dose vial, carton of 1
83831-121-04	2 g/4 mL	Multiple-dose vial, carton of 1

Store vials refrigerated at 2°C to 8°C (36°F to 46°F).

Store diluted solutions of cyclophosphamide according to Table 1 [see *Dosage and Administration (2.3)*].

Cyclophosphamide is a hazardous product. Follow special handling and disposal procedures.¹

17 PATIENT COUNSELING INFORMATION

Advise the patient of the following:

Myelosuppression, Immunosuppression, and Infections

- Inform patients of the possibility of myelosuppression, immunosuppression, and infections (sometimes fatal). Explain the need for routine blood cell counts. Instruct patients to monitor their temperature frequently and immediately report any occurrence of fever [see *Warnings and Precautions (5.1)*].

Urinary Tract and Renal Toxicity

- Advise the patient to report urinary symptoms (patients should report if their urine has turned a pink or red color) and the need for increasing fluid intake and frequent voiding [see *Warnings and Precautions (5.2)*].

Cardiotoxicity

- Inform patients of the possibility of cardiotoxicity (which may be fatal).
- Advise patients to contact a health care professional immediately for any of the following: new onset or worsening shortness of breath, cough, swelling of the ankles/legs, palpitations, weight gain of more than 5 pounds in 24 hours, dizziness or loss of consciousness [see *Warnings and Precautions (5.3)*].

Pulmonary Toxicity

- Warn patients of the possibility of developing non-infectious pneumonitis. Advise patients to report promptly any new or worsening respiratory symptoms [see *Warnings and Precautions (5.4)*].

Secondary Malignancies

- Inform patients that there is an increased risk of secondary malignancies with FRINDOVYX [see *Warnings and Precautions (5.5)*].

Alcohol Content

- Explain to patients the possible effects of the alcohol content in FRINDOVYX, including possible effects on the central nervous system. Patients in whom alcohol should be avoided or minimized should consider the alcohol content of FRINDOVYX. Alcohol could impair their ability to drive or use machines immediately after infusion [see

Warnings and Precautions (5.7)].

Embryo-Fetal Toxicity

- Inform female patients of the risk to a fetus and potential loss of pregnancy. Advise females to inform their healthcare provider of a known or suspected pregnancy [*see Warnings and Precautions (5.8) and Use in Specific Populations (8.1)*].
- Advise female patients of reproductive potential to use effective contraception during treatment and for up to 1 year after completion of therapy [*see Warnings and Precautions (5.8) and Use in Specific Populations (8.1, 8.3)*].
- Advise male patients with female partners of reproductive potential to use effective contraception during treatment and for 4 months after completion of therapy [*see Warnings and Precautions (5.8) and Use in Specific Populations (8.1, 8.3)*].

Lactation

- Advise lactating women not to breastfeed during treatment and for 1 week after the last dose of FRINDOVYX [*see Use in Specific Populations (8.2)*].

Infertility

- Advise males and females of reproductive potential that FRINDOVYX may impair fertility [*see Warnings and Precautions (5.9) and Use in Specific Populations (8.3, 8.4)*].

Common Adverse Reactions

- Explain to patients that side effects such as nausea, vomiting, stomatitis, impaired wound healing, amenorrhea, premature menopause, sterility and hair loss may be associated with cyclophosphamide administration. Other undesirable effects (including, e.g., dizziness, blurred vision, visual impairment) could affect the ability to drive or use machines [*see Adverse Reactions (6.1)*].

Hydration and Important Administration Instructions


- Advise the patients that during or immediately after the administration, adequate amounts of fluid are required to reduce the risk of urinary tract toxicity [*see Dosage and Administration (2.1)*].

Manufactured for:




Avyxa Pharma, LLC
New Jersey 07054, USA
Made in India




PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

<p>NDC 83831-119-01</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>500 mg/mL</p> <p>Must Dilute Before Intravenous Use</p> <p>Multiple-Dose Vial HAZARDOUS DRUG Rx Only</p>	<p>Each vial contains 500 mg cyclophosphamide, USP.</p> <p>Recommended Dosage: See Prescribing Information.</p> <p>Caution: Use only 0.45% Sodium Chloride Injection, USP; 5% Dextrose Injection, USP; or 5% Dextrose and 0.9% Sodium Chloride Injection, USP for diluting this product prior to intravenous infusion.</p> <p>Store vials refrigerated at 2°C to 8°C (36°F to 46°F).</p>	<p>Manufactured for: Avyxa Pharma, LLC New Jersey 07054, USA Made in India</p> <p>Code No: AP/DRUGS/41/2006 119-01 v2/01-2025 150099977</p>  <p>3 8 3 8 3 1 1 9 0 1 2</p>
---	--	--




Frindovyx (cyclophosphamide) Injection, 500mg/mL - Vial Label


<p>Manufactured for: Avyxa Pharma, LLC New Jersey 07054, USA</p> <p>Made in India</p> <p>119-01 v2/01-2025</p> <p>J Code: J9072</p> <p>Code No: AP/DRUGS/41/2006</p>  <p>3 8 3 8 3 1 1 9 0 1 2</p>	<p>NDC 83831-119-01</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>500 mg/mL</p> <p>Must Dilute Before Intravenous Use</p> <p>Multiple-Dose Vial HAZARDOUS DRUG</p> <p>Sterile Rx Only</p> 	<p>Each vial contains 500 mg cyclophosphamide, USP and 500 mg dehydrated alcohol, USP (equivalent to 63% v/v).</p> <p>Recommended Dosage: See Prescribing Information.</p> <p>Caution: Use only 0.45% Sodium Chloride Injection, USP; 5% Dextrose Injection, USP; or 5% Dextrose and 0.9% Sodium Chloride Injection, USP for diluting this product prior to intravenous infusion.</p> <p>Store vials refrigerated at 2°C to 8°C (36°F to 46°F).</p>	<p>NDC 83831-119-01</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>500 mg/mL</p> <p>Must Dilute Before Intravenous Use</p> <p>Multiple-Dose Vial HAZARDOUS DRUG</p> <p>Sterile Rx Only</p> 
---	--	---	--

Frindovyx (cyclophosphamide) Injection, 500 mg/ mL - Carton Label

<p>NDC 83831-120-02</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>1 gram/2 mL (500 mg/mL)</p> <p>Must Dilute Before Intravenous Use</p> <p>Multiple-Dose Vial HAZARDOUS DRUG Rx Only</p>	<p>Each vial contains 1 g cyclophosphamide, USP.</p> <p>Recommended Dosage: See Prescribing Information.</p> <p>Caution: Use only 0.45% Sodium Chloride Injection, USP; 5% Dextrose Injection, USP; or 5% Dextrose and 0.9% Sodium Chloride Injection, USP for diluting this product prior to intravenous infusion.</p> <p>Store vials refrigerated at 2°C to 8°C (36°F to 46°F).</p>	<p>Manufactured for: Avyxa Pharma, LLC New Jersey 07054, USA Made in India</p> <p>Code No: AP/DRUGS/41/2006 120-02 v2/01-2025 150099975</p>  <p>3 8 3 8 3 1 1 2 0 0 2 5</p>
---	---	--

Frindovyx (cyclophosphamide) Injection, 1 g/2 mL - Vial Label

<p>Manufactured for: Avyxa Pharma, LLC New Jersey 07054, USA</p> <p>Made in India</p> <p>120-02 v2/01-2025</p> <p>J Code: J9072</p> <p>Code No: AP/DRUGS/41/2006</p>  <p>3 83831 12002 5</p>	<p>NDC 83831-120-02</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>1 gram/2 mL (500 mg/mL)</p> <p>Must Dilute Before Intravenous Use Multiple-Dose Vial HAZARDOUS DRUG</p> <p>Sterile Rx Only</p> 	<p>Each vial contains 1 g cyclophosphamide, USP and 1000 mg dehydrated alcohol, USP (equivalent to 63% v/v).</p> <p>Recommended Dosage: See Prescribing Information.</p> <p>Caution: Use only 0.45% Sodium Chloride Injection, USP; 5% Dextrose Injection, USP; or 5% Dextrose and 0.9% Sodium Chloride Injection, USP for diluting this product prior to intravenous infusion.</p> <p>Store vials refrigerated at 2°C to 8°C (36°F to 46°F).</p>	<p>NDC 83831-120-02</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>1 gram/2 mL (500 mg/mL)</p> <p>Must Dilute Before Intravenous Use Multiple-Dose Vial HAZARDOUS DRUG</p> <p>Sterile Rx Only</p> 
---	---	--	---

<p>NDC 83831-121-04</p> <p>Frindovyx™ (cyclophosphamide) Injection</p> <p>2 gram/4 mL (500 mg/mL)</p> <p>Must Dilute Before Intravenous Use Multiple-Dose Vial HAZARDOUS DRUG</p> <p>Rx Only</p>	<p>Each vial contains 2 g cyclophosphamide, USP.</p> <p>Recommended Dosage: See Prescribing Information.</p> <p>Caution: Use only 0.45% Sodium Chloride Injection, USP; 5% Dextrose Injection, USP; or 5% Dextrose and 0.9% Sodium Chloride Injection, USP for diluting this product prior to intravenous infusion.</p> <p>Store vials refrigerated at 2°C to 8°C (36°F to 46°F).</p> <p>Manufactured for: Avyxa Pharma, LLC New Jersey 07054, USA</p> <p>Made in India</p> <p>Code No: AP/DRUGS/41/2006</p>	<p>121-04 v2/01-2025</p>  <p>3 83831 12104 6</p> <p>150099976</p>
---	---	---

Frindovyx (cyclophosphamide) Injection, 2 g/4 mL - Vial Label



Frindovyx (cyclophosphamide) Injection, 2 g/ 4 mL - Carton Label

FRINDOVYX

cyclophosphamide injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83831-119
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CYCLOPHOSPHAMIDE (UNII: 8N3DW7272P) (4-HYDROXYCYCLOPHOSPHAMIDE - UNII:1XBF4E50HS)	CYCLOPHOSPHAMIDE ANHYDROUS	500 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
DEHYDRATED ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83831-119-01	1 in 1 CARTON	01/24/2025	
1		1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA210852	01/24/2025	

FRINDOVYX

cyclophosphamide injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83831-120
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CYCLOPHOSPHAMIDE (UNII: 8N3DW7272P) (4-HYDROXYCYCLOPHOSPHAMIDE - UNII:1XBF4E50HS)	CYCLOPHOSPHAMIDE ANHYDROUS	1 g in 2 mL

Inactive Ingredients

Ingredient Name	Strength
DEHYDRATED ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83831-120-02	1 in 1 CARTON	01/24/2025	
1		2 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA210852	01/24/2025	

FRINDOVYX

cyclophosphamide injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83831-121
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
CYCLOPHOSPHAMIDE (UNII: 8N3DW7272P) (4-HYDROXYCYCLOPHOSPHAMIDE - UNII:1XBF4E50HS)		CYCLOPHOSPHAMIDE ANHYDROUS	2 g in 4 mL	
Inactive Ingredients				
Ingredient Name			Strength	
DEHYDRATED ALCOHOL (UNII: 3K9958V90M)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83831-121-04	1 in 1 CARTON	01/24/2025	
1		4 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA210852	01/24/2025		

Labeler - Avyxa Pharma, LLC (128918748)

Registrant - Avyxa Holdings, LLC. (119187191)

Establishment

Name	Address	ID/FEI	Business Operations
Dr.Reddys Laboratories Limited		650446243	ANALYSIS(83831-119, 83831-120, 83831-121) , MANUFACTURE(83831-119, 83831-120, 83831-121) , PACK(83831-119, 83831-120, 83831-121)

Revised: 1/2025

Avyxa Pharma, LLC