

# **TYLENOL PM- acetaminophen and diphenhydramine hydrochloride tablet, film coated**

**Select Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## **Tylenol® PM**

### **Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 500 mg	Pain reliever
Diphenhydramine HCl 25 mg	Nighttime sleep aid

### **Uses**

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

### **Warnings**

#### **Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

**Stop use and ask a doctor if**

- sleeplessness persists continuously for more than 2 weeks.  
Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets at bedtime</li><li>▪ do not take more than 2 caplets of this product in 24 hours</li></ul>
children under 12 years	do not use

**Other information**

- store between 20-25°C (68-77°F)

- **do not use if pouch is torn or damaged**

### **Inactive ingredients**

carnauba wax, crospovidone, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

### **Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:  
McNeil Consumer Healthcare  
Division of McNEIL-PPC, Inc.  
Fort Washington, PA 19034 USA

### **PRINCIPAL DISPLAY PANEL - 2 Caplet Pouch Carton**

Extra Strength

TYLENOL®  
PM

Acetaminophen, Diphenhydramine HCl  
Pain Reliever, Nighttime Sleep Aid  
Non-habit forming

Contains No Aspirin

50 Pouches of  
2 Caplets each

How can we help?  
(1-877-895-3665)



**TYLENOL<sup>®</sup>** Extra Strength  
**PM**

**Acetaminophen, Diphenhydramine HCl**  
Pain Reliever, Nighttime Sleep Aid Non-habit forming

Contains No Aspirin

50 Pouches of  
2 Caplets each

**TYLENOL<sup>®</sup>** Extra Strength  
**PM**

**Acetaminophen, Diphenhydramine HCl**  
Pain Reliever, Nighttime Sleep Aid Non-habit forming

Contains No Aspirin

50 Pouches of  
2 Caplets each

↑  
TO OPEN  
PUSH IN TAB AND PULL OUT

DO NOT USE IF PACKAGE OR POUCH IS OPENED

50 Pouches of 2 Caplets each

### Drug Facts

#### Active ingredients (in each caplet)

Acetaminophen 500 mg.....	Pain reliever
Diphenhydramine HCl 25 mg.....	Nighttime sleep aid

**Uses** temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks.
  - Insomnia may be a symptom of serious underlying medical illness.
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Drug Facts (continued)

#### Directions

• do not take more than directed (see overdose warning)

adults and children 12 years and over	• take 2 Caplets at bedtime
	• do not take more than 2 caplets of this product in 24 hours
children under 12 years	do not use

#### Other information

- store between 20-25°C (68-77°F)
- do not use if pouch is torn or damaged

#### Inactive ingredients

carnauba wax, croscopolidone, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

#### Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



Care To Recycle™

Distributed by:

McNeil Consumer Healthcare  
Division of McNEIL-PPC, Inc.  
Fort Washington, PA 19034 USA  
©McNEIL-PPC, INC. 2020  
Visit us at [www.tylenol.com](http://www.tylenol.com)  
or call toll-free 1-877-TYLENOL  
(1-877-895-3665)

Product Repackaged and Distributed  
with permission of manufacturer by:  
Select Corporation, Carrollton, TX 75007


Select  
Corporation



6 55708 02951 0



2002951142



## TYLENOL PM

acetaminophen and diphenhydramine hydrochloride tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52904-945(NDC:50580-608)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>acetaminophen</b> (UNII: 362O9ITL9D) (acetaminophen - UNII:362O9ITL9D)	acetaminophen	500 mg
<b>diphenhydramine hydrochloride</b> (UNII: TC2D6JAD40) (diphenhydramine - UNII:8GTS82S83M)	diphenhydramine hydrochloride	25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>carnauba wax</b> (UNII: R12CBM0EIZ)	
<b>crospovidone</b> (UNII: 2S7830E561)	
<b>FD&amp;C blue no. 1 aluminum lake</b> (UNII: J9EQA3S2JM)	
<b>hypromellose, unspecified</b> (UNII: 3NXW29V3WO)	
<b>magnesium stearate</b> (UNII: 70097M6I30)	
<b>microcrystalline cellulose</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>polysorbate 80</b> (UNII: 6OZP39ZG8H)	
<b>povidone, unspecified</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>stearic acid</b> (UNII: 4ELV7Z65AP)	
<b>titanium dioxide</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	TY;PM
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52904-945-04	1 in 1 BLISTER PACK	03/01/1997	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:52904-945-05	2 in 1 BLISTER PACK	03/01/1997	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:52904-945-20	20 in 1 CARTON	03/01/1997	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:52904-945-25	25 in 1 CARTON	03/01/1997	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:52904-945-50	50 in 1 CARTON	03/01/1997	
5		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	03/01/1997	

**Labeler** - Select Corporation (053805599)

Revised: 3/2022

Select Corporation