

**LOW DOSE ASPIRIN- aspirin tablet, chewable**  
**WinCo Foods, LLC**

-----  
**1105B-WNC-2021-0830**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for this product

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

**Questions or comments?**

Call 800-824-1706 Monday - Friday 9am - 4pm MST

**PRINCIPAL DISPLAY PANEL**

WinCo® FOODS

\*\*Compare to the active ingredient in BAYER® Chewable Aspirin

Low Dose Aspirin

Orange Flavored

Chewable

ASPIRIN (NSAID) TABLETS / PAIN RELIEVER

36 Tablets 81 mg EACH

INK AND COATING FREE  
FOR LOT AND  
EXPIRATION STAMPING



02032255070  
DO NOT USE IF IMPRINTED SEAL  
UNDER CAP IS BROKEN OR MISSING



WF207 F105801WNC\_RO

**Drug Facts**

**Active ingredient (in each tablet)** **Purpose**  
Aspirin 81 mg (NSAID).....Pain reliever  
\* nonsteroidal anti-inflammatory drug

**Uses**  
■ for the temporary relief of minor aches and pains or as recommended by your doctor  
■ ask your doctor about other uses for this product

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling  
■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug

**Drug Facts (continued)**

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:**  
■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days

**Drug Facts (continued)**

■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur  
■ ringing in the ears or a loss of hearing occurs  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: consult a doctor

**Other information**  
■ store between 20°-25°C (68°-77°F) in a dry place  
■ retain carton for complete product information and warnings

**Inactive ingredients** corn starch, FD&C yellow #6 aluminum lake, titanium dioxide, silicon dioxide, sucrose

**100% SATISFACTION GUARANTEE**  
Chewable Aspirin.  
BAYER HEALTHCARE LLC, distributor of BAYER®  
WINCO FOODS, LLC, BOISE, ID 83704  
DISTRIBUTED BY:  
Monday - Friday 9am - 4pm MST  
Questions? Call 800-824-1706  
**Drug Facts (continued)**



\*\*Compare to the active ingredient in BAYER® Chewable Aspirin

NC

NC

NC

**LOW DOSE ASPIRIN**

aspirin tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-389
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	AP;009
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-389-36	1 in 1 PACKAGE	08/30/2021	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/30/2021	

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 12/2024

WinCo Foods, LLC