DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled Lil' Drug Store Products, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Daytime Cold & Flu

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours

- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

Distributed by: Lil' Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404

PRINCIPAL DISPLAY PANEL - 16 Softgel Blister Pack Carton

QUALITY

* GUARANTEED *

Compare to the Active Ingredients in Vicks® DayQuil® Cold & Flu LiquiCaps®**

Non-drowsy

Cold & Flu Daytime

Acetaminophen, Dextromethorphan HBr, and Phenylephrine HCl Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant

Relieves

- Aches
 Fever
- Sore Throat Cough
- Nasal Congestion

ACTUAL SIZE

16

Softgels

Lil

Drug Store ®

Non-drowsy

Cold & Flu Daytime



Compare to the Active Ingredients in Vicks® DayQuil® Cold & Flu LiquiCaps®**

Non-drowsy

Cold & Flu Daytime

Acetaminophen, Dextromethorphan HBr, and Phenylephrine HCl Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant

Relieves

- Aches Fever
- Sore Throat Cough
- Nasal Congestion



ACTUAL SIZE

16 Softgels



Distributed by: Softg
Lil' Drug Store Products, Inc. 9300 Earhart Lane SW

Softgels are liquid-filled capsules.





Non-drowsy

Daytime

Setaminophen, Dextromethorphan HBr, and Phenylephrine HCI in Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant Cedar Rapids, IA 52404

PLD-A568B FC006103

Lot No: Exp Date:



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts

Active ingredients (In each softgel)

Acetaminophen 325 mg...... Pain relievenfever reducer

Purposes

Dextromethorphan HBr 10 mg......Cough suppressant Phenylephrine HCI 5 mg....... Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:

- minor aches and pains
 headache
 sore throat
- nasal congestion
- fever
- cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- · with other drugs containing acetaminophen
- · 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening
 blisters
 rash

If a skin reaction occurs, stop use and seek medical help

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's

Drug Facts (continued)

disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- · thyroid disease
- · high blood pressure
- trouble urinating due to an enlarged prostate gland · persistent or chronic cough such as occurs with
- smoking, asthma, or emphysema
- cough that occurs with too much phleam (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended

Stop use and ask a doctor if

- · pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- · redness or swelling is present
- new symptoms occur
- · cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- · adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other Information

- store between 15"-30"C (59"-86"F)
- · avoid excessive heat

Inactive ingredients


butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments? call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

**This product is not manufactured or distributed by The Procter & Gamble Company, Vicks[®], DayQuil[®], and LiquiCaps® are registered trademarks of The Procter & Gamble Company.



68158C-US-11-19-PA



DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

Product	Inform	ation
PIOUUCL		auvii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:66715-6715

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients		
Ingredient Name	Strength	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)		
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		
MANNITOL (UNII: 3OWL53L36A)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SORBITAN (UNII: 6092ICV9RU)		
SORBITOL (UNII: 506T60A25R)		

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	P19
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715- 6715-8	8 in 1 CARTON	02/15/2020	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/05/2014	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 10/2022 Lil' Drug Store Products, Inc.