

**DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled  
Lil' Drug Store Products, Inc.**

-----  
**Daytime Cold & Flu**

***Drug Facts***

<b><i>Active ingredients (in each softgel)</i></b>	<b><i>Purposes</i></b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

### **Other information**

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

### **Inactive ingredients**

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

call toll-free **1-877-507-6516 (M-F 8AM-4:30PM CST)**

Distributed by:

Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW  
Cedar Rapids, IA 52404

### **PRINCIPAL DISPLAY PANEL - 16 Softgel Blister Pack Carton**

QUALITY

★ GUARANTEED ★

Compare to the Active Ingredients in  
Vicks® DayQuil® Cold & Flu LiquiCaps®\*\*

Non-drowsy

Cold & Flu  
Daytime

Acetaminophen, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant

Relieves

- Aches • Fever
- Sore Throat • Cough
- Nasal Congestion

ACTUAL SIZE

16  
Softgels

Lil'  
Drug Store®

Non-drowsy

# Cold & Flu Daytime



Compare to the Active Ingredients in  
Vicks® DayQuil® Cold & Flu LiquiCaps®\*\*

Non-drowsy

# Cold & Flu Daytime

**Acetaminophen**, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant

Relieves

- Aches • Fever
- Sore Throat • Cough
- Nasal Congestion



ACTUAL SIZE

**16**  
Softgels



Distributed by:  
Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW

Softgels are  
liquid-filled  
capsules.



Non-drowsy



**16 Softgels**

**Cold & Flu  
Daytime**

**Acetaminophen**, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant


Cedar Rapids, IA 52404

PLD-A568B FC006103



Lot No:

Exp Date:



**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**  
**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Drug Facts**

**Active Ingredients**      **Purposes**  
**(In each softgel)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
- minor aches and pains    • headache    • sore throat
- nasal congestion        • fever
- cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening    • blisters    • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's

**Drug Facts (continued)**

disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease            • diabetes
- heart disease            • thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other Information**

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

**Inactive Ingredients**

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**

call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

\*\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

68158C-US-11-19-PA

PRINT SIDE GLUE



## DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-6715
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P19
<b>Contains</b>			

### Packaging

		<b>Marketing Start</b>	<b>Marketing End</b>
--	--	------------------------	----------------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6715-8	8 in 1 CARTON	02/15/2020	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/05/2014	

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2025

Lil' Drug Store Products, Inc.