ISOPROPYL ALCOHOL 70 PERCENT- isopropyl alcohol liquid MEIJER, INC.

Drug Facts

Active ingredient (by volume)

Isopropyl alcohol (70% conc.)

Purpose

First aid antiseptic

Uses

first aid to help prevent the risk of infection in minor cuts, scrapes, and burns.

Warnings

For external use only.

• if taken internally, serious gastic disturbances will result

flammable, keep away from fire or flame.

• use only in a well-ventilated area; fumes may be toxic

Ask a doctor before use if you have

deep or puncture wounds, animal bites, or serious burns.

When using this product

- do not get into eyes
- do not apply over large areas of the body
- do not use longer than one week unless directed by a doctor

Stop use and ask a doctor if

condition persists or gets worse.

Keep out of reach of children.

In case of ingestion, get medical help or contact a Poison Control Center right away (1-800-222-1222.

Directions

- clean affected area
- apply small amount of this product on the area 1-3 times daily
- if bandaged, let dry first

• may be covered with a sterile bandage

Other information

- store at room temperature
- does not contain, nor is intended as a substitute for grain or ethyl alcohol

Inactive ingredient

purified water

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

70% Isopropyl Alcohol

First Aid Antiseptic

For Rubbing and Massaging

WARNING: FLAMMABLE, KEEP AWSY FROM SPARK, HEAT OR FLAME.

FL OZ (mL)


DIST. BY MEIJER

DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

Package Label



Drug Facts

Active ingredient (by volume)

Purpose

Isopropyl alcohol (70% conc.)......First aid antiseptic

Uses first aid to help prevent the risk of infection in minor cuts, scrapes, and burns,

Warnings

For external use only.

- if taken internally, serious gastric disturbances will result Flammable, keep away from fire or flame.
- use only in a well-ventilated area; fumes may be toxic

Ask a doctor before use if you have deep or puncture wounds, animal bites, or serious burns.

When using this product do not get into eyes

- do not apply over large areas of the body
- do not use longer than one week unless directed by a doctor

Stop use and ask a doctor if condition persists or gets worse.

Keep out of reach of children. In case of ingestion, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ clean affected area

- apply small amount of this product on the area 1-3 times daily
- if bandaged, let dry first
- may be covered with a sterile bandage

Other information store at room temperature

does not contain, nor is intended as a substitute for grain or ethyl alcohol

Inactive ingredient purified water

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

DIST. BY MEIJER DISTRIBUTION, INC. **GRAND RAPIDS, MI 49544** www.meijer.com



PID 364689 PLD-A226C LB006910



MEIJER Isopropyl Alcohol 70%

ISOPROPYL ALCOHOL 70 PERCENT

isopropyl alcohol liquid

Product Information

HUMAN OTC DRUG NDC:41250-265 Item Code (Source) **Product Type**

Route of Administration TOPICAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	70 mL in 100 mL		

Inactive Ingredients				
Ingredient Name	Strength			
WATER (UNII: 059QF0KO0R)				

l	P	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:41250- 265-32	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2019	06/30/2025	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M003	06/30/2019	06/30/2025	

Labeler - MEIJER, INC. (006959555)

Revised: 1/2024 MEIJER, INC.