LEADER CHILDRENS PAIN AND FEVER- acetaminophen suspension Cardinal Health 110, LLC. dba Leader

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cardinal Health Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- · the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

36-47	4-5 years	5 years 7.5 mL	
48-59	6-8 years	10 mL	
60-71	9-10 years	12.5 mL	
72-95	11 years	15 mL	

^{*}or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Children's Pain & Fever

Acetaminophen 160 mg / 5 mL

Oral Suspension

Pain Reliever / Fever Reducer

Cherry Flavor

Aspirin Free

Ibuprofen Free

Alcohol Free

Dye Free

Ages 2-11 Years

COMPARE TO CHILDREN'S TYLENOL® active ingredient

100% Money Back Guarantee

4 FL OZ (118mL)



Pain Reliever / Fever Reducer Action of the control Fever & nigq Children's

LEADEŔÄ

LEADER 2

Children's

Acetaminophen 160 mg / 5 mL **Oral Suspension** Pain Reliever / Fever Reducer

Cherry

Aspirin Free Ibuprofen Free Alcohol Free Dye Free

Ages 2-11 Years

DIRECTIONS: STEP 1. Shake well



STEP 2. Remove child protective . сар



STEP 3. Squeeze your child's dose into the dosina cup



IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.

Drug Facts

Active ingredient (in each 5 mL)

Purpose

Acetaminophen 160 mg....Pain reliever/fever reducer

- Uses temporarily:
 reduces fever
 relieves minor aches and pains due to:
 the common cold
 the use headache
 sore throat
 toothache

Warnings

- Liver warning: This product contains acetaminophen.

 Severe liver damage may occur if your child takes

 more than 5 doses in 24 hours, which is the maximum daily amount

daily amount us containing acetaminophen Allergy alert. Acetaminophen may cause severe skin reactions. Symptoms may include:

skin readening ### bitests ### askin reactions cocurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by rever, headache, rash, nausea, or vomitting, consult a doctor promptly.

- Do not use

 with any other drug containing acetaminophen
 (prescription or nonprescription). If you are not sure
 whether a drug contains acetaminophen, ask a doctor
 or pharmacist.

 if your child has ever had an allergic reaction to this
 product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days
- new symptoms occur
 redness or swelling is present
 These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- This product does not contain directions or complete warnings for adult use do not give more than directed (see overdose).
- warning)
 shake well before using
 mL = milliliter

- m L = millitier
 find right dose on chart below. If possible, use weight to dose, otherwise, use age.
 remove the child protective cap and squeeze your child's dose into the dosing cup
 repeat dose every 4 hours while symptoms last
 do not give more than 5 times in 24 hours

weight (ID)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or missing

Inactive ingredients

INACTIVE INGREMENTS
anhydrous citric acid, calcium sulfate, carrageenan,
flavor, glycerin, hydroxyethyl cellulose, microcrystallin
cellulose and carboxymethylcellulose sodium,
procylene glycoi, procylparaben, purified water,
sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

CAPDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. DISTRIBUTED BY CARDINAL HEALTH DUBLIN, OHIO 43017 www.myleader.com 1-800-200-6313

100% Money Back Guarantee Return to place of purchase.



COMPARE TO

CHILDREN'S

TYLENOL®

active ingredien

4 FL OZ (118mL)



95926 E9 C5

LEADER CHILDRENS PAIN AND FEVER

ORAL

acetaminophen suspension

Product Information

Route of Administration

Product Type HUMAN OTC DRUG Item Code (Source) NDC:70000-0481

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)			
CARRAGEENAN (UNII: 5C69YCD2YJ)			
GLYCERIN (UNII: PDC6A3C0OX)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K6790BS311)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
PROPYLPARABEN (UNII: Z8IX2SC10H)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)			

Product Characteristics			
Color	WHITE (opaque, viscous)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

F	Packaging			
#	tem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000- 0481-1	1 in 1 CARTON	02/25/2019	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/25/2019	
rinai			

Labeler - Cardinal Health 110, LLC. dba Leader (063997360)

Revised: 6/2021