

**PAMPRIN MAX MENSTRUAL PAIN RELIEF- acetaminophen and aspirin and caffeine tablet, film coated**  
**Focus Consumer Healthcare, LLC**

-----  
**Pamprin Max Menstrual Pain Relief**

Pain reliever

Diuretic

\*nonsteroidal anti-inflammatory drug

**(in each caplet)**

Acetaminophen 250 mg

Aspirin 250 mg

Caffeine 65 mg

D&C Red No. 27 Lake, FD&C Red No. 40 Lake, fractionated coconut oil, hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, silica, sodium starch glycolate, starch, stearic acid, talc, titanium dioxide (245-171)

For the temporary relief of these symptoms associated with menstrual periods:

- cramps
- headache
- backache
- fatigue
- bloating

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Allergy warning:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- 

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.

### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

### **Ask a doctor or pharmacist before use**

if you are taking a prescription drug for gout, diabetes or arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- redness or swelling is present
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- ringing in the ears or loss of hearing occurs

## **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **When using this product**

- you may get drowsy, avoid alcoholic beverages
- alcohol sedatives and tranquilizers may increase drowsiness
- use caution when driving or operating machinery
- excitability may occur especially in children

## **stop use and ask doctor if**

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

**If pregnant or breastfeeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a poison control center right away.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

## **Adults and children 12 years and over:**

- take 2 caplets with water every 6 hours as needed
- do not exceed 8 caplets in a 24 hour period or as directed by a doctor
- do not use more than directed (see warnings)

**children under 12 years:** ask a doctor

Maximum Strength Pamprin


Menstrual Pain Relief

Max Pain+Energy

contains Acetaminophen, Aspirin (NSAID) & Caffeine

2 Pain Relievers + Caffeine

Cramps Headache Fatigue Bloating Backache



**Drug Facts (continued)**

- ringing in the ears or loss of hearing occurs. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions adults and children 12 years and over:**


- take 2 caplets with water every 6 hours as needed
  - do not exceed 8 caplets in a 24 hour period or as directed by a doctor
  - do not use more than directed (see warnings)
- children under 12 years:** ask a doctor

**Inactive ingredients** fractionated coconut oil, hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, red 27 lake, red 40 lake, silica, starch, stearic acid, talc, titanium dioxide

0044946-01

www.pamprih.com

Distributed by: Focus  
Consumer Healthcare, LLC  
801 Broad Street  
Chattanooga, TN 37402  
©2022 Focus Consumer  
Healthcare, LLC



**DRUG FACTS CONTINUED UNDER FRONT FLAP**

**Drug Facts (continued)**

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day using this product
- Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
  - take a blood thinning (anticoagulant) or steroid drug
  - have had stomach ulcers or bleeding problems
  - are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too

LOT  
EXP

**Drug Facts (continued)**

- more than 8 caplets in 24 hours, which is the maximum daily amount may occur if you take
- Liver warning: This product contains acetaminophen. Severe liver damage if a skin reaction occurs, stop use and seek medical help right away.
- may include: • skin reddening • blisters • rash
- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms include: • hives • facial swelling • asthma (wheezing) • shock
- Allergy alert: Aspirin may cause a severe allergic reaction which may syndrome, a rare but serious illness.
- consult a doctor because these symptoms could be an early sign of Reye's using this product; if changes in behavior with nausea and vomiting occur, from chicken pox or flu-like symptoms should not use this product. When
- Reye's syndrome: Children and teenagers who have or are recovering

GA R50990  
GB R50991

- Uses for the temporary relief of minor aches and pains and temporary:
  - water-weight gain • bloating • swelling • full feeling • fatigue
- associated with the premenstrual and menstrual periods.

**IF PRINTED INNER SEAL ON BOTTLE IS BROKEN OR MISSING, DO NOT USE.**

Easy-open bottle with child-resistant packaging. Purchase only if carton end flaps are sealed. Questions or comments? [www.pamprih.com](http://www.pamprih.com)



## PAMPRIN MAX MENSTRUAL PAIN RELIEF

acetaminophen and aspirin and caffeine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71687-3006
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	

### Product Characteristics

<b>Color</b>	pink (FD & C Red No. 40) , pink (D&C REd No. 27)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	MAX
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71687-3006-6	1 in 1 CARTON	10/18/2017	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/18/2017	

**Labeler** - Focus Consumer Healthcare, LLC (080743737)

Revised: 12/2023

Focus Consumer Healthcare, LLC