# SUNMARK ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated Strategic Sourcing Services LLC

-----

## McKesson Anti-Diarrheal Drug Facts

## Active ingredient (in each caplet)

Loperamide HCl 2 mg

## **Purpose**

Anti-diarrheal

#### Use

controls symptoms of diarrhea, including Travelers' Diarrhea

## Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

#### Do not use

if you have bloody or black stool

## Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

## Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

## When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

## Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

#### Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

## **Inactive ingredients**

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

## Questions or comments?

1-800-719-9260

## **Principal Display Panel**

sunmark®

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT

See New Warnings

Loperamide Hydrochloride Tablets, 2 mg

anti-diarrheal

Controls the symptoms of diarrhea

**ANTI-DIARRHEAL** 

**Actual Size** 

12 CAPLETS\*

(\*Capsule-shaped tablets)

**GLUTEN FREE** 

cel Ini ose, polyethylen e glycol, pregelati nized starch l ake, h yp romel lose, ma gnesium ste arate, micro crystal line D &C yell own o. 1 Oal umin um I ake, FD &C b lue n o. 1 alu minu m *In active ing redien ts* anhydrouslætoæ, carnauba wax,

Orug Facts (continued)

1-800-71 9-9260 Questions or comments?

**Drug Facts** (continued)

noitemrotni redto

es n 10 n ob

sak a doctor

more than 2 caplets in 24 hours

more than 3 caplets in 24 hours

sub sequent loose stool; but no

m ore than 4 caplets in 24 hours

subsequent loose stool; but no

S ca plets after the first loose

stool; 1 caplet after each

o n fu d ;loofs esoo l fine upes du s

stool; 1/2 cap let after each

t ca plet after the first loose

stool; 1/2 caplet after each

t caplet after the first loose

(adl 65 of qu)

un der 2 years

(3d Th of 1bs)

2-5 years

(99169-81P)

6-8 years

(¤q196-09)

9-11 years

children

and over

12 years

nenblida

adults and

Dire ctions

children

children

children

e xbiration date profe ssi onal be fore use. se end panel for lot number and If pre gnant or breast-fee ding, a sk a health ■ store at 20°-25°C (68°-77°F)

- These may be signs of a serious condition. ■you getabdominal swelling or bulging.
  - diarrhea lasts for more than 2 days
    - eymptoms get worse Stop use and ask a doctor if

when driving or operating machinery. drowsiness or dizziness may occur. Be careful When using this product tiredness,

brescription drugs. Loperamide mayinteract with certain it you are taking a prescription drug. Ask a doctoror pharmacist before use

- mativith theart is mortal in eart in yithm
  - ■a. history of liver disease
- ■fever mucusin the stool Ask a doctorbefore use if you have

Do not use if you have blo ody or black stool

can se serion s heart problems or death Heart alert: Taking more than drected can rash o r other all ergic reaction to lo peramide H Cl Allergy a lert: Do not use if you have ever had a sbuju je M

including Travelers' Diambea

USE controls symptoms of distribea,

Loperamide HCl's mg`.....hrti-diarmeal (in each caplet) es od ind Active ingredient

Drug Facts

\*\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

weight to dose; otherwise, use age.

drink plenty of clear fluids to help

i find right do se on chart. If possible, use prevent dehydration caused by diarrhea

Control Centerright away (1-800-222-1222).

Keep out of reach of children. In case of

overdo se, g et medical hel porco ntact a Poison

Drug Facts (continued)

## **M**SKESSON

Another Quality Product Distributed By McKesson 6555 State Highway 161, Irving, TX 75039 Money Back Guarantee Please visit us at www.sunmarkbrand.com



COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT\*\* NDC 49348-529-02

See New Warnings

sunmark<sup>®</sup>

Loperamide Hydrochloride Tablets, 2 mg

anti-diarrhea

OPEN OTHER END

22453 S1 C6



#### SUNMARK ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:49348-529

Route of Administration ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength

LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - LOPERAMIDE - UNII:6X9OC3H4II)

LOPERAMIDE HYDROCHLORIDE 2 mg

## **Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)
MAGNESIUM STEARATE (UNII: 70097M6I30)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

**D&C YELLOW NO. 10** (UNII: 35SW5USQ3G)

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)

#### **Product Characteristics**

1 Todact Characteristics					
Color	GREEN	Score	2 pieces		
Shape	OVAL	Size	10mm		
Flavor		Imprint Code	L2		
Contains					

Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:49348-529- 34	1 in 1 CARTON	11/16/2004	12/31/2021		
1		96 in 1 BOTTLE; Type 0: Not a Combination Product				
2	NDC:49348-529- 02	12 in 1 CARTON	08/21/2003			
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product				
3	NDC:49348-529- 04	24 in 1 CARTON	07/15/2009			
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product				
4	NDC:49348-529- 08	1 in 1 CARTON	06/27/2003	12/31/2021		
4		48 in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA075232	06/27/2003			

## **Labeler -** Strategic Sourcing Services LLC (116956644)

Revised: 11/2022 Strategic Sourcing Services LLC