

**XTRA GUARD (PART A)- lactic acid solution**  
**AgroChem USA, LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**XTRA GUARD (Part A)**

XTRA GUARD (Part A)

CIO2 Teat Dip Part A

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**

**NOT FOR HUMAN USE.**

**AVOID FREEZING.**

**KEEP OUT OF DIRECT SUNLIGHT.**

**CAUSES SERIOUS EYE IRRITATION.**

**FOR FARM AND INDUSTRIAL USE ONLY.**

**PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.**

**FIRST AID / RESPONSE:**

**IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.**

**STORAGE: Store locked up.**

**DISPOSAL: Dispose of contents/ container to an approved waste disposal plant**

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**NDC: 17307-3018-3**

**NET CONTENTS: 55 GALLON (208.2L)**

**LOT:**

**EXP. DATE:**

**ACTIVE INGREDIENT: Lactic Acid 3.0%**

**Use XTRA Guard effectively to reduce the spread of mastitis causing organisms.**

**Recommended for post milking applications.**

**DIRECTIONS:**

**IMPORTANT:** MUST BE MIXED PRIOR TO USE. Mix PART A & PART B in equal parts. Mix by volume. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA GUARD solution after 24 hours.

**General Use**

**Once mixed according to directions , do not mix product with any other teat dips, or other product.**

**Use a teat dip cup to apply. Clean dip cup just prior to use.**

**If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.**

**If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.**

**If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.**

**This product can also be foamed or sprayed using recommended equipment.**

**Pre Dipping Use**

**Perform forestripping and inspect teats for clinical mastitis.**

**Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels**

**Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.**

**Allow product to stand for 30 seconds.**

**Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.**

**Post Dipping Use**

**Immediately after each milking, use product.**

**Dip each teat full-length into the teat dip cup containing product. Allow to**

air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

Manufactured By:

AgroChem USA, LLC 26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

## XTRA GUARD (PART A)

ClO<sub>2</sub>  
TEAT DIP  
PART A

### CAUTION

NOT FOR HUMAN USE.  
AVOID FREEZING.  
KEEP OUT OF DIRECT SUNLIGHT.  
CAUSES SERIOUS EYE IRRITATION.  
KEEP OUT OF REACH OF CHILDREN.  
FOR FARM AND INDUSTRIAL USE ONLY.



**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

**FIRST AID/ RESPONSE:**

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/ container to an approved waste disposal plant.

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 (CN205494)

ACTIVE INGREDIENT: Lactic Acid 3.0%

NDC: 17307-3018-3

NET CONTENTS: 55 GALLONS (208.2L)

LOT:  
EXP:



Use XTRA Guard effectively to reduce the spread of mastitis causing organisms. Recommended for post milking applications.

#### DIRECTIONS:

**IMPORTANT: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts. CREATES CHLORINE DIOXIDE.** Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA Guard after 24 hours.

#### General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product. Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product.

Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment.

#### Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip.

Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

#### Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

#### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

Manufactured By: AgroChem USA, LLC  
26 Freedom Way  
Saratoga Springs, NY 12866  
Phone: 518-226-4850



## XTRA GUARD (PART A)

lactic acid solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17307-3018
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	0.03 kg in 1 kg

### Inactive Ingredients

Ingredient Name	Strength
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3018-1	19 kg in 1 CONTAINER		
2	NDC:17307-3018-2	56.8 kg in 1 DRUM		
3	NDC:17307-3018-3	208.2 kg in 1 DRUM		
4	NDC:17307-3018-4	1003.1 kg in 1 CONTAINER		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/01/2023	

**Labeler** - AgroChem USA, LLC (118336116)

**Registrant** - AgroChem USA, LLC (118336116)

### Establishment

Name	Address	ID/FEI	Business Operations
AgroChem USA, LLC		118336116	api manufacture, manufacture, analysis, label, pack

Revised: 12/2023

AgroChem USA, LLC