HABITROL NICOTINE POLACRILEX 4 MG COATED CINNAMON- nicotine polacrilex gum, chewing Dr. Reddy's Laboratories Inc.

Nicotine Polacrilex 4 mg Cinnamon Coated USP Gum

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex (equal to 4 mg nicotine)

Purpose

Stop smoking aid

Use

 reduces withdrawal symptoms, including nicotine craving associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth,teeth, or jaw problem occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- oral blistering occurs
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up. use 4 mg nicotine gum according to the following 12 week schedule

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use a least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health provider.

Other information

- each piece contains: calcium 115 mg, sodium 11 mg
- store at 20 25°C (68 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, butylated hydroxytoluene, carnauba wax, FD&C red #40 lake, gum base, hydroxypropyl cellulose, hypromellose, natural and artificial, cinnamon flavors, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

Questions or comments?

Call toll-free 1-800-585-8682 weekdays (8:00 am - 8: 00 pm EST)

Principal Display Panel

*Compare to the active ingredient in Nicorette® Gum

Habitrol®

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

STOP SMOKING AID

Gum

4 MG

Sugar-free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP

If you smoke your first cigarette <u>MORE THAN</u> 30 MINUTES after waking up use Habitrol 2 mg Gum.

Coated Cinnamon Flavor

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivate to quit
- 2. **Use Enough** Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks
- 3. **Use long enough** Use Nicotine Polacrilex Gum for the full 12 weeks
- 4. Use with a Support Program as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- Not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

^{*}This product is not manufactured or distributed by Haleon group of companies

distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

DISTRIBUTED BY:

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Package Label

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- 3. Use Long Enough Use Nicotine Polacrilex Gum for the full 12 weeks.
- Use With a Support Program as directed in the enclosed User's Guide.







starting at corner

NDC 55741-001-80

*Compare to the active ingredient in Nicorette® Gum

Habitrol **Nicotine Polacrilex Gum** USP, 4 mg (nicotine)

STOP SMOKING AID

Gum



Sugar-free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Habitrol 2 mg Gum.

80 PIECES 4 mg EACH



Coated Cinnamon Flavor

Habitrol®

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

STOP SMOKING AID



und autoput builz est If you are under 18 years of age, ask, a boolsh before use If you are under the control med by the company as before using the pools, med the value bland before before therebox and other important information. It is not a begon and a label to use the proof of the control med by the control media after webling up, If you entities your that department them there is no median up, and the control median after webling up,

erough rhochre to make children and pals sick. Wrap used phoose of gurn in paper and throw away in the teash is case of owndose, get medical help or contact a freison Control Center (1-500-2521-1522) right away. Keep out of reach of children and pals. Pieces of nicotine gum may have в Алифрии од ви вувиђе изверои (апер ве одусту), ризарији от изву)

teachest bigs bes, ass

4 mg EACH 08

if you smoke your first algerette MORE THAN 30 MINUTES FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES WE WENTED WITHIN 30 MINUTES WITHIN 30 MIN

after waking up, use Habitrol 2 mg Gum.



"This product is not manufactured or distributed by Haleon group of companies distributor of Micovotor" Gum.

(35) holi-free 1-**800-565-6662** weekdays (8:00 am - 8:00 pm EST)

bicarbonale, aodum carbonale, aucralose, fait, fitanium dioxide, xyflol

batolyon ingeredients sessis as extentens prisessim, betalente prisessim, betalenten betalenten betalenten betalen bet Step use and sak a doctor if m mouth, touth, or jaw pratierns occur m insgular hearbest or palpitations occur m Existing prescription medicine for depression or estima. Your prescription does may need to be adjusted. and a property of the larger o PLD-8620C FC009488 (411-89) 3.52-02 (8 axos m er calcium 115 mg, sodium 11 mg Drup funyous dos superio-usu a Bush w Other Information 53 Ask a declor or pharmacist before use if you are pudia, bayon po yeeb yuuu surayyudi pay po kon, yeegyy cana buoxigeo m to not use more than 24 phoess a day.

If you leet you need to use the gum for a 03/5059 m high blood pressure, not controlled with medication. Micotine can increase blood pressure, If you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another within may cause you hoozen, heerflourin, neuses or other side effects. Dr. Reddy's Laboratories, Inc. Princeton, NJ 08540 Pearl decise, recent heart attack, or irregular heartbeak. Nicotine can increase your heart rate. THE USTRIBUTED BY: M is socium-restricted diet m is socium-restricted diet ■ to improve your chances of quilding, use at least 9 pieces per day for the first 6 soud a fluwout bus Alector usung usewhad if Alect maff, selghat til Datu glwods mug odt werls an emitten elegt uer filler, utlage gelevente niged, aveng et egent ele mer Maring gelennen GD, Luckold emog as legen elle to been filter associag skill taleger an ellen, to, prog enchooling of grinweits erolled esskuning it volt kinds to the bro ob an B occider gum to a medicine and must be used a certain way to get the best SID 4 HORIZ \$100 Z DE L 410 g ponz medices withdrawal symptoms, including nicotine crawing, associated with Anna acoud i St of Ot saleeM B of T saleeM Alcotine polacifiex (squal to 4 ing nicetine). m y hon emoto hom. Uses cigaraste within 30 minutes of washed up, use 4 mg michine gum exceeding to the following 12 week acheolule: Active ingredient (in each chewing piece) Purpose gund Lacts (continued)



ваитине јо битени и

York, pearl uster

гринты

Europus Busselt

Drug Facts

HABITROL Nicotine Polacrilex Gum Coated Cinnamon Flavor

HABITROL NICOTINE POLACRILEX 4 MG COATED CINNAMON

nicotine polacrilex gum, chewing

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:55741-001

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) **NICOTINE** 4 mg

Inactive Ingredients Ingredient Name Strength **ACACIA** (UNII: 5C5403N260) ACESULFAME POTASSIUM (UNII: 230V73Q5G9) **BUTYLATED HYDROXYTOLUENE** (UNII: 1P9D0Z171K) CARNAUBA WAX (UNII: R12CBM0EIZ) FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T) HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) **SODIUM BICARBONATE** (UNII: 8MDF5V39QO) **SODIUM CARBONATE** (UNII: 45P3261C7T) **SUCRALOSE** (UNII: 96K6UQ3ZD4) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) XYLITOL (UNII: VCQ006KQ1E)

Product Characteristics							
Color	pink (Dark pink)	Score	no score				
Shape	SQUARE	Size	14mm				
Flavor	CINNAMON	Imprint Code					
Contains							

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:55741- 001-80	8 in 1 CARTON	07/28/2025			
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product				
2	NDC:55741- 001-15	15 in 1 CARTON	07/28/2025			
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA079219	07/28/2025			

Labeler - Dr. Reddy's Laboratories Inc. (802315887)

Revised: 7/2025 Dr. Reddy's Laboratories Inc.