

**UMBRELLA SUNSCREEN- octinoxate titanium dioxide lotion**  
**CA-BOTANA INTERNATIONAL**

-----  
**6236 Umbrella Sunscreen Lotion**

For external use only. Do not use on damaged or broken skin. When using this product keep out of eyes. Rinse with water to remove. Stop use and ask a doctor if rash occurs. Keep out of reach of children. If product is swallowed, get medical help or contact a poison control center right away.

Helps prevent sunburn. If used as directed with other sun protection measured decreases the risk of skin cancer and early skin aging caused by the sun. Apply liberally 15 minutes before sun exposure. Use a water resistant sunscreen if swimming or sweating. Reapply: at least 2 hours. Children under 6 months: Ask a doctor. Sun protection measurements. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m - 2 p.m. Wear long-sleeve shirts, pants, hats, and sunglasses. Protect this product from excessive heat and direct sun

ALOE VERA LEAF, ALUMINUM STARCH OCTENYLSUCCINATE, ASCORBIC ACID, ASCORBYL PALMITATE, BORON NITRIDE, CETOSTEARYL ALCOHOL, CETEARYL ISONONANOATE, CITRIC ACID MONOHYDRATE, CAPRYLIC/CAPRIC MONO/DIGLYCERIDES, ETHYLHEXYLGLYCERIN, GLYCERIN, SUNFLOWER OIL, ISOMALT, LECITHIN SOYBEAN, MACADAMIA OIL, NIACIN, PANTHENOL, PHENOXYETHANOL, LECITHIN SOYBEAN, POTASSIUM SORBATE, PROPANEDIOL, ROSA MOSCHATA OIL, ROSEMARY, SODIUM BENZOATE, HYALURONATE SODIUM, TETRASODIUM IMINODISUCCINATE, TOCOPHEROL, GRAPE SEED OIL, WATER

Sunscreen

Titanium Dioxide, Octinoxate

For external use only.

Avoid contact with eyes.

Keep out of reach of children.

Do not apply to open wounds.

STOP USE AND ask a doctor if condition worsens or symptoms persist for more than seven days, discontinue use of the product.

Helps prevent sunburn. If used as directed with other sun protection measured decreases the risk of skin cancer and early skin aging caused by the sun. Apply liberally 15 minutes before sun exposure. Use a water resistant sunscreen if swimming or sweating. Reapply: at least 2 hours. Children under 6 months: Ask a doctor. Sun protection measurements. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m - 2 p.m. Wear long-sleeve shirts, pants, hats, and sunglasses. Protect this product from excessive heat and direct sun

Package label principal display panel




DOCTOR D.  
SCHWAB®

UMBRELLA SUNSCREEN

BROAD SPECTRUM  
SPF 40

ENVIRONMENTAL PROTECTION

CRÈME SOLAIRE HYDRATANTE  
HUMECTANTE PROTECCIÓN SOLAR



Drug Facts	
Active ingredients	Purpose
Titanium Dioxide 6%.....	Sunscreen
Octinoxate 7.5%.....	Sunscreen
<b>Uses</b> • helps prevent sunburn • if used as directed with other sun protection measured (see <b>Directions</b> ), decreases the risk of skin cancer and early skin aging caused by the sun.	
<b>Warnings</b> • <b>For external use only</b>	
<b>When using this product</b> keep out of eyes. Rinse with water to remove.	
<b>Do not use</b> on damaged or broken skin.	
<b>Stop use and ask a doctor</b> if rash occurs.	
<b>Keep out of reach of children.</b> If product is swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> • apply liberally 15 minutes before sun exposure • use a water resistant sunscreen if swimming or sweating • reapply at least every 2 hours • children under 6 months: Ask a doctor • <b>Sun Protection Measures</b> • Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeve shirts, pants, hats, and sunglasses.	
<b>Inactive ingredients:</b> Aloe barbadensis Extract, Aluminum Starch Octenylsuccinate, Ascorbic Acid, Ascorbyl Palmitate, Boron Nitride, Camellia sinensis (Tea) Leaf Extract, Caprylic/Capric Triglycerides, Cetearyl Alcohol, Cetearyl Isononanoate, Cetyl Alcohol, Citric Acid, Citrus sinensis (Orange) Oil, Coco Caprylate, Emulsifying Wax, Ethylhexylglycerin, Glycerin, Helianthus annuus (Sunflower) Seed Oil, Isomalt, Lecithin, Macadamia ternifolia Seed Oil, Niacin, Phenoxyethanol, Phosphatidylcholine, Potassium Sorbate, Propanediol (Zemea), Rosa canina (Rosehip) Fruit Oil, Rosmarinus officinalis leaf (Rosemary) extract CO2, Sodium Benzoate, Sodium Hyaluronate, Tetrasodium EDTA, Tocopherol, Vitis vinifera (Grape) Fruit Cell Extract, Water (Aqua), Xanthan Gum.	
<b>Other information</b> • protect this product from excessive heat and direct sun.	

NDC#: 35192-010-02  
REF #6236

DOCTOR D. SCHWAB®  
Manufactured by CA BOTANA  
San Diego, CA 92121, USA  
Made in USA  
[www.DoctorSchwabCA.com](http://www.DoctorSchwabCA.com)



08120666236

**UMBRELLA SUNSCREEN**

octinoxate titanium dioxide lotion

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:35192-010
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	6.8 g in 113.4 g
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	8.5 g in 113.4 g

## Inactive Ingredients

Ingredient Name	Strength
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>ALUMINUM STARCH OCTENYLSUCCINATE</b> (UNII: I9PJ006294)	
<b>ASCORBIC ACID</b> (UNII: PQ6CK8PD0R)	
<b>ASCORBYL PALMITATE</b> (UNII: QN83US2B0N)	
<b>BORON NITRIDE</b> (UNII: 2U4T60A6YD)	
<b>CETOSTEARYL ALCOHOL</b> (UNII: 2DMT128M1S)	
<b>CETEARYL ISONONANOATE</b> (UNII: P5O01U99NI)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>CAPRYLIC/CAPRIC MONO/DIGLYCERIDES</b> (UNII: U72Q2I8C85)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SUNFLOWER OIL</b> (UNII: 3W1JG795YI)	
<b>ISOMALT</b> (UNII: S870P5502W)	
<b>SOYBEAN OIL</b> (UNII: 241ATL177A)	
<b>MACADAMIA OIL</b> (UNII: 515610SU8C)	
<b>NIACIN</b> (UNII: 2679MF687A)	
<b>PANTHENOL</b> (UNII: WW9CM0067Z)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>PROPANEDIOL</b> (UNII: 5965N8W85T)	
<b>ROSA MOSCHATA OIL</b> (UNII: J99W255AWF)	
<b>ROSEMARY</b> (UNII: IJ67X351P9)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>ETHYLHEXYLGLYCERIN</b> (UNII: 147D247K3P)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>TETRASODIUM IMINODISUCCINATE</b> (UNII: GYS41J2635)	
<b>TOCOPHEROL</b> (UNII: R0ZB2556P8)	
<b>GRAPE SEED OIL</b> (UNII: 930MLC8XGG)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35192-010-02	113.4 g in 1 TUBE; Type 0: Not a Combination Product	11/09/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	06/01/2012	

---

**Labeler** - CA-BOTANA INTERNATIONAL (106276728)

**Registrant** - RODOLFO UGELSTAD (106276728)

**Establishment**

Name	Address	ID/FEI	Business Operations
CA-BOTANA INTERNATIONAL		106276728	manufacture(35192-010)

Revised: 12/2024

CA-BOTANA INTERNATIONAL