

XTORO - finafloxacin suspension
Fonseca Biosciences, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use XTORO safely and effectively. See full prescribing information for XTORO.

**XTORO (finafloxacin otic suspension) 0.3%,
for topical otic use**
Initial U.S. Approval: 2014

----- **INDICATIONS AND USAGE** -----

XTORO® is a quinolone antimicrobial indicated for the treatment of acute otitis externa (AOE) caused by susceptible strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*. (1)

----- **DOSAGE AND ADMINISTRATION** -----

Instill four drops in the affected ear(s) twice daily for seven days. For patients requiring use of an otowick, the initial dose can be doubled (to 8 drops), followed by 4 drops instilled into the affected ear twice daily for seven days. (2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Otic Suspension: finafloxacin 0.3% (3)

----- **CONTRAINDICATIONS** -----

None. (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Prolonged use of this product may lead to overgrowth of nonsusceptible organisms. Discontinue use if this occurs. (5.1)
- Allergic reactions may occur in patients with a history of hypersensitivity to finafloxacin, to other quinolones, or to any of the components in this medication. Discontinue use if this occurs. (5.2)

----- **ADVERSE REACTIONS** -----

The most common adverse reactions occurring in 1% of patients with XTORO were ear pruritus and nausea. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Fonseca Biosciences, LLC at 1-877-436-6732 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 3/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Growth of Resistant Organisms with Prolonged Use

5.2 Allergic Reactions

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

XTORO® is indicated for the treatment of acute otitis externa (AOE) with or without an otowick, caused by susceptible strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in patients age 1 year and older.

2 DOSAGE AND ADMINISTRATION

Instill four drops into the affected ear(s) twice daily for seven days. For patients requiring use of an otowick, the initial dose can be doubled (to 8 drops), followed by 4 drops instilled into the affected ear twice daily for seven days.

Important administration instructions include:

- Warm the suspension by holding the bottle in the hand for one or two minutes prior to dosing in order to avoid dizziness which may result from the instillation of a cold suspension. Shake bottle well before use.
- Lie with the affected ear upward, instill the drops, and maintain the position for 60 seconds to facilitate penetration of the drops into the ear canal.
- Repeat if necessary for the opposite ear.

3 DOSAGE FORMS AND STRENGTHS

Otic Suspension: Each mL of XTORO contains finafloxacin 0.3% (equivalent to 3 mg finafloxacin).

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Growth of Resistant Organisms with Prolonged Use

As with other antibacterial preparations, prolonged use of XTORO may lead to overgrowth of nonsusceptible organisms, including yeast and fungi. If this occurs, discontinue use and institute alternative therapy.

5.2 Allergic Reactions

Allergic reactions to XTORO may occur in patients with a history of hypersensitivity to fluroxacin, to other quinolones, or to any of the components in this medication. If this occurs, discontinue use and institute alternative therapy.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

A total of 618 patients were treated with XTORO in two Phase 3 clinical trials. The most frequently reported adverse reactions of those exposed to XTORO occurring at an incidence of 1% included ear pruritus and nausea.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate or well-controlled studies with XTORO in pregnant women. Fluroxacin was shown to be teratogenic in rabbits and rats following oral administration. Neural tube defects and skeletal anomalies in both species, and limb anomalies in rabbits, were observed at exposures estimated to be at least 1300 times the maximum human systemic exposure following topical otic administration of 0.3% fluroxacin. Because animal studies are not always predictive of human responses, XTORO should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Data

Animal Data

In rabbit embryofetal studies, maternal toxicity was not observed at oral doses up to 9 mg/kg/day (estimated 8000 times the maximum human systemic exposure [0.234 ng/mL] following topical otic administration with 0.3% fluroxacin). Fetal toxicity was observed at the lowest dose tested, 1 mg/kg/day (estimated 1300 times the maximum human systemic exposure following topical otic administration with 0.3% fluroxacin), and included exencephaly, enlarged fontanel, spina bifida, phocomelia, paw hyperflexure, missing lumbar vertebra, missing lumbar arch, and sternbra fusion.

In a rat embryofetal study, no adverse maternal toxicity was observed at oral doses up to 100 mg/kg/day (estimated 60,000 times the maximum human systemic exposure following topical otic administration with 0.3% fluroxacin). The developmental no observed adverse effect level (NOAEL) was 30 mg/kg (estimated 22,000 times the maximum human systemic exposure following topical otic administration with 0.3% fluroxacin). Exencephaly was observed in one fetus at 100 mg/kg. At 500 mg/kg, additional developmental toxicities were observed including increased preimplantation loss, decreased fetal weight, decreased placental weight, increased incidence of non-ossified sternebrae, and delayed ossifications in the sternebrae, xiphisternum, sacral arches and metacarpals.

8.2 Lactation

Fluroxacin has been identified in the milk of nursing rats following oral administration. The human systemic concentration of XTORO following topical otic treatment is low [see *Clinical Pharmacology (12.3)*]. It is not known whether topical otic administration could result in sufficient systemic absorption to produce detectable quantities in the human breast milk. Caution should be exercised when fluroxacin is administered to a nursing mother.

8.4 Pediatric Use


The safety and efficacy of XTORO in infants below one year of age have not been established. The safety and efficacy of XTORO in treating acute otitis externa in pediatric patients one year or older have been demonstrated in adequate and well controlled clinical trials [see *Clinical Studies (14)*].

8.5 Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

11 DESCRIPTION

XTORO (fluroxacin otic suspension) 0.3% is a quinolone antimicrobial. Its chemical name is (-)-8-cyano-1-cyclopropyl-6-fluoro-7-[(4a*S*,7a*S*)-hexahydropyrrolo[3,4-*b*]-1,4-oxazin-6(2*H*)-yl]- 4-oxo-1,4-dihydroquinoline-3-carboxylic acid (CAS number 209342-40-5). Its structural formula is:



Finafloxacin has a molecular weight of 398.4. Finafloxacin is a white to yellow powder or crystals that is slightly soluble in water (0.125 mg/mL).

XTORO (finafloxacin otic suspension), 0.3% is supplied as a sterile, preserved, aqueous suspension for topical otic use. It has a pH of approximately 6.0 and an osmolality of approximately 290 mOsm/kg.

XTORO contains **Active ingredient:** finafloxacin, 0.3%. **Preservative:** benzalkonium chloride (0.005%). **Inactive ingredients include:** sodium chloride, hydroxyethylcellulose, tyloxapol, magnesium chloride, and purified water. May contain hydrochloric acid and/or sodium hydroxide to adjust pH.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Finafloxacin is a fluoroquinolone antimicrobial [see *Microbiology (12.4)*].

12.3 Pharmacokinetics

Finafloxacin plasma concentrations were evaluated following single or repeated ototopical doses of XTORO (finafloxacin otic suspension) 0.3%. In healthy subjects administered 4 drops in each ear twice daily for seven days, quantifiable finafloxacin concentrations were observed in 2 of 14 subjects; and these concentrations were just above the quantitation limit (0.05 ng/mL). Similarly, in AOE patients administered a single dose of 4 or 8 drops in each ear, quantifiable finafloxacin concentrations of up to 0.234 ng/mL were observed in plasma samples from 2 of 36 AOE patients.

12.4 Microbiology

Finafloxacin belongs to the fluoroquinolone class of antibacterials which involves the inhibition of bacterial type II topoisomerase enzymes, DNA gyrase and topoisomerase IV, which are required for bacterial DNA replication, transcription, repair and recombination.

Finafloxacin has been shown to be active against most isolates of the following bacteria, both in vitro and clinical studies as described in the INDICATIONS and USAGE section of the package insert for XTORO

Pseudomonas aeruginosa
Staphylococcus aureus

Mechanism of Resistance

Resistance to fluoroquinolones occurs primarily by mutations in the chromosomal DNA that encode for DNA gyrase and DNA topoisomerase enzymes, decreased outer membrane permeability or drug efflux mechanisms. *In vitro* resistance to fleroxacin due to spontaneous mutation is rare.

Cross Resistance

Cross-resistance has been observed between fleroxacin and other fluoroquinolones. No cross-resistance has been observed between fleroxacin and other classes of antibacterial agents.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity

Animal studies have not been conducted to determine the carcinogenic potential of fleroxacin.

Mutagenesis

Floxacin was shown to be genotoxic and clastogenic *in vitro*, with and without metabolic activation, and *in vivo*. In a bacterial reverse mutation assay, fleroxacin was positive in only one strain (TA102). Fleroxacin was positive in mammalian cell culture assays: mouse lymphoma cell forward mutation assays, a mutagenicity assay in V79 Chinese hamster lung cells, and a micronucleus test in V79 cells. Fleroxacin was clastogenic in mouse micronucleus studies.

Impairment of fertility

An oral rat fertility study detected a NOAEL for male and female fertility of 100 mg/kg/day (estimated 60,000 times the maximum human systemic exposure following topical otic administration with 0.3% fleroxacin). At 500 mg/kg/day, males were completely infertile, presumably due to low sperm count and sperm immobility.

General toxicity studies in rats have confirmed sperm toxicity following oral and intravenous dosing. Following intravenous dosing, the NOAEL for sperm toxicity was 30 mg/kg/day (150,000 times the maximum human exposure following topical otic administration with 0.3% fleroxacin).

14 CLINICAL STUDIES

In two randomized multicenter, vehicle controlled clinical trials, XTORO dosed four drops twice daily for 7 days was superior to its vehicle for both clinical and microbiological outcomes as well as in time to cessation of ear pain in patients with acute otitis externa (AOE).

Among 560 patients (161 with an otowick) that were pathogen positive (baseline microbiological specimen that contained *Staphylococcus aureus* and/or *Pseudomonas*

aeruginosa), clinical cure on Day 11 was 71% in XTORO versus 37% in Vehicle. Among 1234 patients who received study treatment (Intent to Treat population (ITT)), aged 6 months to 85 years, clinical cures were 71% for XTORO and 50% in Vehicle.

Clinical Cures^a at Day 11 (Pathogen Positive Subset and ITT)

	Study 1			Study 2		
	XTORO	Vehicle	XTORO vs. Vehicle Difference (95% CI)	XTORO	Vehicle	XTORO vs. Vehicle Difference (95% CI)
Pathogen + Subset	104/145 (71.7%)	46/138 (33.3%)	38.4% (27.6%, 49.1%)	101/147 (68.7%)	52/130 (40.0%)	28.7% (17.4%, 40.0%)
ITT	245/344 (71.2%)	173/342 (50.6%)	20.6% (13.5%, 27.8%)	194/274 (70.8%)	134/274 (48.9%)	21.9% (13.9%, 29.9%)

^a A clinical cure was attained if the sum of the numerical scores of the 3 signs and symptoms of AOE (tenderness, erythema, and edema) was 0 at Day 11 (TOC).

The median time to cessation of ear pain in pathogen positive patients treated with XTORO was 3.5 days compared to 6.8 days in Vehicle. The median time to cessation of ear pain in ITT patients treated with XTORO was 3.5 days compared to 5.3 days in Vehicle.

Median Time (in Days) to Cessation of Ear Pain (Pathogen Positive Subset and ITT)

	Study 1			Study 2		
	XTORO	Vehicle	XTORO vs. Vehicle Difference (95% CI)	XTORO	Vehicle	XTORO vs. Vehicle Difference (95% CI)
Pathogen + Subset	4.0	7.0	-3.0 (-5.0, -0.8)	3.0	6.5	-3.6 (-5.0, -2.0)
ITT	4.0	5.0	1.0 (-2.0, -0.5)	3.0	6.5	-2.2 (-3.0, -1.0)

Among the pathogen positive patients, microbiological success (eradication of all baseline organisms) was achieved on Day 11 in 67% in XTORO versus 13% in the Vehicle treated patients.

Microbiological Success^b at Day 11 (Pathogen Positive Subset)

	Study 1			Study 2		
	XTORO	Vehicle	XTORO vs. Vehicle Difference (95% CI)	XTORO	Vehicle	XTORO vs. Vehicle Difference (95% CI)
			52.0%			

Pathogen + Subset	97/145 (66.9%)	18/138 (13.0%)	55.9% (44.4%, 63.4%)	97/147 (66.0%)	15/130 (11.5%)	54.4% (45.0%, 63.9%)
^b Microbiological success was attained if all pre-therapy bacteria were absent from the exit otic specimen. The presence of fungi and/or yeast was not considered in the determination of microbiological success.						

In clinically cured pathogen positive patients, XTORO demonstrated eradication rates of 89% in both *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Vehicle eradication rates were 33% for *Staphylococcus aureus* and 20% for *Pseudomonas aeruginosa*.

16 HOW SUPPLIED/STORAGE AND HANDLING

XTORO (flaxloxacina otic suspension) 0.3% is a sterile, preserved, aqueous, otic suspension supplied in an opaque plastic bottle with a controlled drop tip and a white cap:

5 mL fill in a 5 mL bottle (NDC 83709-001-01)

Storage and Handling

Store at 2°C to 25°C (36°F to 77°F). Do not freeze.

17 PATIENT COUNSELING INFORMATION

Allergic Reactions

Advise patients that if a rash or allergic reaction occurs, they should discontinue the use of the product immediately and contact their physician.

Warm the Bottle in Hands Before Use

Advise patients or caregivers that prior to administration of XTORO, they should warm the bottle by holding it in their hands for one or two minutes to avoid dizziness which may result from the instillation of a cold solution.

For Use with an Otowick

Advise patients that following instillation of 8 drops at the time of otowick insertion, they should continue with the lower dose of 4 drops administered twice daily for seven days.

U.S. Patents Nos: 8,536,167; 9,119,859; 9,504,691; 9,993,483

©2026 Fonseca Biosciences, LLC

Manufactured For:

Fonseca Biosciences, LLC

2807 Long Beach Blvd Ste 1A

Long Beach Township, New Jersey 08008 USA

1-877-436-6732

medinfo@fonsecabio.com

PATIENT INFORMATION
XTORO (ex tore' oh)

(finafloxacin otic suspension) 0.3%
for topical otic use

What is XTORO?

XTORO is a prescription medicine that is used to treat bacterial infections of the external ear canal, with or without an otowick in people 1 year of age and older. It is not known if XTORO is safe and effective in children under 1 year of age.

Before using XTORO, tell your healthcare provider about all of your medical conditions, including if you:

- allergic to any antibacterial including fluoroquinolones or to any of the ingredients in XTORO. See the end of this Patient Information leaflet for a complete list of the ingredients in XTORO.
- are pregnant or plan to become pregnant. It is not known if XTORO will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if XTORO can pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with XTORO.

Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use XTORO?

- XTORO is for use in the ear only. **Do not use XTORO in the eyes or mouth.**
- Use XTORO exactly as your healthcare provider tells you to take it.
- Read the **Instructions for Use** that comes with XTORO.

What are the possible side effects of XTORO?

XTORO may cause serious side effects, including:

- **Growth of yeast or fungi.** XTORO may cause growth of yeast or fungi during long term use. Stop using XTORO if you develop a growth of yeast or fungi.
- **Allergic reactions.** Stop using XTORO and call your healthcare provider or go to the nearest emergency room right away if you have any of the following signs or symptoms of an allergic:
 - rash
 - hives
 - swelling of your face, lips, mouth, or tongue
 - itching
 - trouble breathing
 - dizziness, fast heartbeat, or pounding in your chest.

The most common side effects of XTORO include:

- itching in the ear
- nausea

Tell your healthcare provider if you have any side effects that bothers you or does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store XTORO?

- Store XTORO between 36°F to 77°F (2°C to 25°C).

- Do not freeze XTORO.

Keep XTORO and all medicines out of the reach of children.

General Information about the safe and effective use of XTORO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use XTORO for a condition for which it was not prescribed. Do not give XTORO to other people, even if they have the same symptoms that you have. It may harm them. This Patient Information leaflet summarizes the most important information about XTORO. You can ask your pharmacist or healthcare provider for information about XTORO that is written for healthcare professionals.

What are the ingredients in XTORO?

Active ingredients: finafloxacin, 0.3%

Inactive ingredients: sodium chloride, hydroxyethylcellulose, tyloxapol, magnesium chloride, and purified water. Hydrochloric acid and sodium hydroxide may be added to adjust pH.

U.S. Patent Nos: 8,536,167; 9,119,859; 9,504,691; 9,993,483

©2026 Fonseca Biosciences, LLC

Manufactured For: Fonseca Biosciences, LLC 2807 Long Beach Blvd Ste 1A Long Beach Township, New Jersey 08008 USA

1-877-436-6732

medinfo@fonsecabio.com

This Patient Information has been approved by the U.S. Food and Drug Administration
Revised: 3/2026

PATIENT INSTRUCTIONS FOR USE

XTORO (ex tore' oh)

(finafloxacin otic suspension) 0.3%

for topical otic use

Important information you need to know before using XTORO.

- **XTORO is for use in the ear only.** Do not use XTORO in the eyes or mouth.
- Shake XTORO well before use.
- **Do not** touch the ear, fingers or any other surfaces with the tip of the bottle. This could contaminate the drops.

How should I use XTORO?

1. Wash your hands

Wash your hands with soap and water.

2. Warm and shake the bottle

Hold the bottle in your hands for 1 to 2 minutes to warm XTORO (see Figure 1). Using cold drops may cause dizziness. Shake the bottle of XTORO well before use.



Figure 1

3. Add the drops

Lie down on your side with the affected ear facing up (see Figure 2).




Figure 2

Squeeze the bottle and put **4 drops** of XTORO into the affected ear (see Figure 3). If otowick is in place, put 4 drops of XTORO into the affected ear at the surface of the otowick. **Do not touch** the ear, fingers or any other surfaces **with the tip of the bottle** as it could contaminate the drops.




Figure 3

4. Pull ear lobe

Gently pull the outer ear lobe upward and backward (see Figure 4). This will allow the ear drops to flow down into the ear canal.




Figure 4

5. Stay on your side

Stay on your side with the affected ear facing up for at least 60 seconds. Repeat Steps 3-5 for the other ear if both ears are infected.

How should I store XTORO?

- Store XTORO between 36°F to 77°F (2°C to 25°C).
- Do not freeze XTORO.

Keep XTORO and all medicines out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration
Revised: 3/2026

PRINCIPAL DISPLAY PANEL - 5 mL


NDC 83709-001-01


Xtoro® (finafloxacin otic suspension) 0.3%

® Fonseca Biosciences, LLC

STERILE

5 mL





PRINCIPAL DISPLAY PANEL- 1.5 mL SAMPLE

NDC 83709-001-90

Xtoro® (finafloxacin otic suspension) 0.3%

® Fonseca Biosciences, LLC

STERILE

1.5 mL



3600940 01


SAMPLE. NOT FOR SALE.

Each mL contains: Active: finafloxacin 3 mg
Storage: Store at 2°C - 25°C (36°F - 77°F)
Do Not Freeze. Dosage: see Prescribing Information.
Manufactured for: Fonseca Biosciences, LLC
Long Beach Twp, NJ 08008 USA
Product of Spain. © 2026 Fonseca Biosciences



NDC 83709-001-90
XTORO[®]
(finafloxacin otic suspension) 0.3%
STERILE 1.5 mL  Rx Only

LOT
EXP



SAMPLE. NOT FOR SALE.

Rx Only

NDC 83709-001-90

XTORO[®]
(finafloxacin otic suspension) 0.3%

Dosage: see Prescribing Information.

Storage: Store at 2°C-25°C (36°F-77°F) **Do Not Freeze.**



**FOR TOPICAL
USE IN THE EAR**

Manufactured for:
Fonseca Biosciences, LLC
Long Beach Twp, NJ 08008 USA

LOT
EXP

FONSECA
BIOSCIENCES
3601084 01



Sterile
1.5 mL

PRINCIPAL DISPLAY PANEL - 5 mL SAMPLE


NDC 83709-001-91

Xtoro® (finafloxacin otic suspension) 0.3%

® Fonseca Biosciences, LLC

STERILE

5 mL



XTORO

finafloxacin suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83709-001
Route of Administration	AURICULAR (OTIC)		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Finafloxacin (UNII: D26OSN9Q4R) (Finafloxacin - UNII:D26OSN9Q4R)	Finafloxacin	3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Benzalkonium Chloride (UNII: F5UM2KM3W7)	
Sodium Chloride (UNII: 451W47IQ8X)	
Hydroxyethyl Cellulose (1500 Mpa.s At 1%) (UNII: L605B5892V)	
Tyloxapol (UNII: Y27PUL9H56)	
Magnesium Chloride (UNII: 02F3473H9O)	
Water (UNII: 059QF0KO0R)	
Hydrochloric Acid (UNII: QTT17582CB)	
Sodium Hydroxide (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83709-001-01	1 in 1 CARTON	05/01/2026	
1		5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
2	NDC:83709-001-91	1 in 1 CARTON	05/01/2026	
2		5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
3	NDC:83709-001-90	1 in 1 CARTON	05/01/2026	
3		1 in 1 POUCH		
3		1.5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA206307	05/01/2026	

Labeler - Fonseca Biosciences, LLC (118929241)**Registrant** - Fonseca Biosciences, LLC (118929241)

Establishment

Name	Address	ID/FEI	Business Operations
Siegfried El Masnou, S.A.		462029349	manufacture(83709-001)

Revised: 3/2026

Fonseca Biosciences, LLC